

## Intermediate disturbance by off-road vehicles promotes endangered pioneer cryptogam species of acid inland dunes

### Moderate Störungen durch Geländefahrzeuge fördern gefährdete Pionier-Kryptogamenarten in sauren Binnendünen

Piotr T. Zaniewski<sup>1,\*</sup> , Łukasz Kozub<sup>2</sup>  & Małgorzata Wierzbicka<sup>2</sup>

<sup>1</sup>Department of Forest Botany, Institute of Forest Science, Warsaw University of Life Sciences,  
ul. Nowoursynowska 159, 02-776, Warsaw, Poland;

<sup>2</sup>Institute of Environmental Biology, Faculty of Biology, University of Warsaw, ul. Żwirki i Wigury 101,  
02-089 Warsaw, Poland

\*Corresponding author, e-mail: lukasz.kozub@biol.uw.edu.pl

#### Abstract

Psammophilous grassland communities of inland dunes often occurring together with common juniper scrub are among the most threatened habitat types in Central Europe. Once they were related to disturbance caused by traditional agriculture and forest management (sheep grazing, burning, litter raking and overexploitation of forests). Currently, after the above-mentioned drivers have disappeared, those communities are subjected to secondary succession leading to loss of their typical biodiversity, an important component of which are cryptogams (bryophytes and lichens). In the presented study we ask whether the recently increasing off-road activity, concentrated in dune areas, can maintain such habitats and their biodiversity or instead leads to an even faster deterioration.

To answer this question, we studied the cryptogam diversity in an off-road disturbed inland dune environment of a seldom used military training area located in Central Poland. A set of vegetation samples was collected, together with a range of measured or estimated variables including soil parameters, off-road disturbance intensity and factors related to ecological succession, like common juniper scrub cover.

The obtained results suggest that highest off-road intensity led to a transformation of the habitat into its very initial stage with active dune processes, while within undisturbed patches the succession led to the development of initial pine forest. The highest number of lichen and bryophyte species occurred within not or only slightly disturbed habitats. However, lichens preferred less shrub cover than bryophytes. Nevertheless, pioneer, typical psammophilous grassland species, which also include some threatened lichen species, benefited from small to medium disturbance intensity. Diverse and spatially complex off-road activity increases the number of microhabitats and can thus be a positive factor promoting the coexistence of all ecological groups of cryptogams connected with the acid inland dune environment. In our opinion, off-road activity may to some extent be used as an inexpensive nature conservation tool in order to maintain open, cryptogam-rich dune habitats.

**Keywords:** bryophytes, disturbance, *Juniperus communis*, lichens, military training site, psammophilous grassland, *Stereocaulon*

**Erweiterte deutsche Zusammenfassung am Ende des Artikels**

## 1. Introduction

Grasslands on acid inland dunes comprise an important habitat for the conservation of biodiversity in Europe and are protected as Natura 2000 habitats (EUROPEAN COMMISSION 2013). In nemoral and boreal vegetation zones, they are principally semi-natural habitat types, created as a result of man-made deforestation and long-lasting intensive human-induced disturbance (including high grazing pressure) (SCHWABE et al. 2013, DENGLER et al. 2020a). Their area significantly decreased in the 20<sup>th</sup> century, primarily due to the abandonment of traditional land-use practices and subsequent intentional afforestation as well as natural succession (WOLFF et al. 2017, DENGLER et al. 2020b). Habitat patches dominated by cryptogam vegetation (the so-called “lichen steppes”) are of particularly high conservation value as they are the main habitat for many rare and threatened lichens (KETNER-OOSTRA et al. 2012). In Central Europe open inland dunes are also often occupied by common juniper, *Juniperus communis* (BARKMAN 1985). Nowadays this species is also in increasing danger of extinction in Europe, mainly due to habitat loss and low natural regeneration (VERHEYEN et al. 2009, BROOME et al. 2017).

Although military training sites are to be found all over the world, they have still been insufficiently explored. Scientific research on the development of biological diversity under conditions of intensive exploitation by military manoeuvres has been undertaken on a somewhat larger scale since the 1990s, when as a result of political changes those areas became more accessible to civilians. Military training grounds are specific sites where intensive disturbance caused by human activities dramatically changes the environment (MILCHUNAS 2000) and the heterogeneity of the landscape is higher than that of its surroundings (VANDERPOORTEN et al. 2005). Surprisingly, in times of abandonment of traditional agricultural land-use practices, it turned out that many ecosystems and species that were once widespread and related to agricultural landscapes found their only refuge within military training areas (WARREN & BÜTTNER 2008, JENTSCH et al. 2009). This may be due to the diverse and heterogeneous habitat properties of military training areas, where newly created microsites such as exposed soil, blast craters, tyre tracks of heavy vehicles, trenches and burned parts can become valuable habitats, e.g. psammophilous grasslands and heathlands, accommodating a large number of species (IUCN 1996, ROMAHN 1998, MILCHUNAS et al. 2000, WARREN et al. 2007, GUGNACKA-FIEDOR & ADAMSKA 2010, KRAWCZYK et al. 2019 etc.). Good light conditions coupled with suitably intense landscape disturbance processes like those occurring in military areas are particularly important for the conservation of endangered bryophytes and lichens, especially those related to otherwise disappearing pioneer conditions (MOTIEJUNAITE 1996, VANDERPOORTEN et al. 2005, DINGOVÁ 2010, GUGNACKA-FIEDOR & ADAMSKA 2010, ADAMSKA & DEPTUŁA 2015, KRAWCZYK et al. 2019).

The use of off-road vehicles in dune systems is considered a negative factor in many countries (e.g. RICKARD et al. 1994, KUTIEL et al. 2000). In some parts of Central and Eastern Europe, this problem has only recently arisen, and former military training areas are becoming heavily utilised by off-road amateurs. There have only been a few studies on the relationship between off-road impact and cryptogam biology, which focused mainly on the recovery time of cryptobiotic soil crusts after disturbance (BELNAP 1993, LALLEY & VILES 2008). There is little information about the relationship between off-road disturbance and cryptogam richness in changing, dynamic habitats. Some valuable bryophytes are known to be connected with a disturbed environment (VANDERPOORTEN et al. 2005), and there is a high probability that lichens may also benefit from it.

Due to political changes in the world and changes in military doctrine resulting from technological progress, some military facilities were closed, e.g. in Poland after 1989 (IUCN 1996). This poses a threat to valuable disturbance-dependent ecosystems and species (JENTSCH et al. 2009). Off-road activity very often appears in abandoned military training areas as their landscape heterogeneity and often unregulated ownership status attracts both legal and illegal off-road activities. In the presented study we wanted to check whether off-road activities can successfully replace the former military disturbance in an abandoned training area and maintain the threatened cryptogam biodiversity in its acid sandy grasslands. To achieve this, we tried to define the main ecological factors influencing cryptogam communities in the disturbed, changing environment of an open acid inland dune area of a partly abandoned military training area including, among other environmental variables, off-road disturbance intensity.


## 2. Study area

The research was carried out within a military training area called Grochalskie Piachy (ca. 250 ha), located in Central Poland (52°24'15"N, 20°37'05"E), 22 km northwest of the capital city, Warsaw. The site is nowadays part of the Kampinos National Park. The area lies within the pre-glacial Vistula river valley on deep river sands remodelled by the aeolian process in the early Holocene and is part of one of the largest and best-developed dune belts in Europe (ZGORZELSKI & PAWŁOWSKA 2003). The mean annual temperature of the site is 7.7 °C, the mean annual rainfall 546.9 mm, the mean annual temperature amplitude 24.6 °C, the mean daily amplitude 9.1 °C and the length of the growing period 185 days (ANDRZEJEWSKA 2003). The terrain of the military range is rugged, with many small depressions and hills, but altitude differences are small (77–83 m a.s.l.). Nowadays dune processes are still present in large parts of the site. However, only small dune forms (sand ripples and low hillocks) are observed. The open areas of the military range were created on purpose by deforestation (as foreground of a fortress) in the late nineteenth and early twentieth century, and until the last decade of the 20<sup>th</sup> century, they were used only for military purposes. The military range is composed of three open sand fields with active dune processes triggered by military activities. The open areas (ca. 100 ha) are predominantly covered with *Corynephorion canescentis* Klika 1931 vegetation (MATUSZKIEWICZ 2015). Nowadays the northern part of the site is still under heavy disturbance caused by regular demolition of old unexploded bombs from World War II. The remnant part, where the study was conducted, is not used for military purposes, and no other forms of management have been applied in recent decades. The whole non-forested area of the military training area is utilised by enthusiasts of off-road vehicles such as four-wheel drive cars, all-terrain vehicles (ATVs), motorcycles and motocross vehicles.

## 3. Material and methods

### 3.1 Sampling methods

The fieldwork was carried out in the growing season of 2010. In order to minimise the risk of sampling the same habitat path more than once, the following procedure was applied: The central and southern part of the military range was evenly covered with a 25 m × 25 m grid. Within the squares, only evenly disturbed habitat patches were sampled. No more than one sample per square was taken (Fig. 1). Squares with no suitable habitat patches were omitted. Each sample was a 64 m<sup>2</sup> square plot.




**Fig. 1.** Location of sampling plots within the Grochalskie Piachy military training area.

**Abb. 1.** Lage der Untersuchungsflächen auf dem militärischen Übungsgelände Grochalskie Piachy.

The relatively large plot size was chosen in order to obtain averaged results as within these highly dynamic habitats, disturbance intensity and shrub cover vary. The cover of trees and shrubs, of common juniper in the shrub layer and of open substrate and litter were visually estimated with a precision of 5%. Vascular plants in the herb layer, terrestrial bryophytes, liverworts and lichens were sampled using a modified (BARKMAN et al. 1964) BRAUN-BLANQUET (1928) scale (original vegetation data provided in Supplement E1 and environmental variables in Supplement E2). Samples of problematic species were collected for laboratory analyses. To account for natural habitat variability originating from topography, the inclination of each plot was measured with the use of a hand inclinometer. In each sampling plot a set of 25 evenly distributed sampling points (2 m apart from each other) was selected. The visible off-road disturbance and dune processes were noted as present or absent in each of these sampling points. As off-road disturbance we classified any sign indicating that the area had been driven on during the growing season, usually visible as linear tyre tracks. This included situations when the soil crust was compressed or even broken as well as situations when a tyre track appeared within a completely bare patch of sand. Dune processes representing naturally occurring disturbance were identified by visible sand ripples and/or partially buried plants of *Corynephorus canescens*. Hillocks and grooves made by insects were rather rare and not taken into account. No other natural disturbances were recorded within the study plots. From the 9 sampling points in the centre (Fig. 2), mixed soil samples were collected from a depth of 5–10 cm.

### 3.2 Species determination and laboratory analyses

Cryptogam species were determined using standard morphological and anatomical methods: for bryophytes according to SMITH (2004), for lichens according to SMITH et al. (2009) and for liverworts according to DAMSHOLT (2002). Thin Layer Chromatography (TLC) was also used for the determination of lichen chemical compounds in the *Cladonia chlorophaea* and *C. coccifera* groups according to ORANGE et al. (2001). Sieving analysis of the soil samples was performed with mesh sizes of 1 mm (very coarse sand and gravel), 0.5 mm (coarse sand), 0.25 mm (medium sand) and 0.1 mm (fine sand),



**Fig. 2.** Method of data collection within a 64-m<sup>2</sup> sampling plot 1) sampling points located 2 m apart from each other, 2) 25 points used for the assessment of disturbance and dune processes intensity, 3) 9 central points used for the soil sample collection.

**Abb. 2.** Methode der Datenerfassung innerhalb einer 64-m<sup>2</sup>-Fläche 1) 2 m voneinander entfernte Untersuchungsflächen; 2) Störungsintensität und Dünenprozesse wurden an allen 25 Punkten beurteilt; 3) an den mittleren 9 Punkten wurden Bodenproben genommen.

the remaining fractions representing very fine sand, silt and clay. For chemical analysis the soil samples were cleaned using a 1-mm sieve. Conductivity and pH were determined with an electronic pH/conductivity meter in a suspension of 1 part of sand and 5 parts of water by volume after standing for 24 hours. The total organic matter content was measured as loss on ignition at 550 °C.

### 3.3 Cryptogam groups and nomenclature

Valuable cryptogam species were assessed according to Polish national red lists (OCHYRA 1992, CIEŚLIŃSKI et al. 2006) and lists of species protected by national law. Ecological groups of cryptogams were described according to KETNER-OOSTRA et al. (2012) and slightly modified to adapt them to local climatic conditions according to succession models of FALIŃSKI et al. (1993). Recorded lichens were assigned to four groups: pioneer, humicolous, humicolous/aero-hygrophytes and aero-hygrophytes (KETNER-OOSTRA et al. 2012). The nomenclature of vascular plants follows MIREK et al. (2002), liverworts DAMSHOLT (2002), bryophytes SMITH (2004) and lichens SMITH et al. (2009). We decided to use *Cladonia chlorophaea* s.l. (including recognised *C. chlorophaea*, *C. merochlorophaea* and *C. grayi*), *Cladonia coccifera* s.l. (including recognised *C. diversa* and *C. pleurota*), *Cladonia macilenta* s.l. (including recognised *C. macilenta* and *C. floerkeana*) and *Hypnum cupressiforme* s.l. Abbreviations of cryptogam species names used in the presented figures are explained in Table 1. Species names of vascular plants are abbreviated using the first three letters of the genus name and the first four letters of the species epitheton.

### 3.4 Statistical analysis

The values obtained for the intensity of off-road disturbance and dune processes (ranging from 0 to 25, cf. section 3.1) were recalculated into percentages and used as continuous variables in the analysis. After that, a pairwise correlation analysis of all independent variables was conducted using Pearson's linear correlation coefficient. In cases where  $|r| > 0.7$ , one of the strongly correlated variables was removed from further analyses until no such cases were left. Firstly, a very strong correlation ( $r = 0.99$ ) appeared between total shrub cover and cover of common juniper in the shrub layer. Only the first variable was used in further analyses, but it can be assumed that they can be used interchangeably. We also removed from further analyses bare soil cover and intensity of dune processes, which were correlated to disturbance intensity (probably because disturbance creates open sand and triggers dune processes), litter cover, which was correlated to shrub cover, and soil pH, which was negatively correlated to organic matter content. Also, middle soil fractions were removed as they were correlated either with the largest or the smallest soil separates (Table 2). The set of values representing the spatial arrangement of sampling plots ( $x, y, xy, x^2, y^2, x^3, y^3, x^2y$  and  $xy^2$ ) was calculated according to BORCARD et al. (1992) on the basis of their coordinates in the PUWG 1992 metric coordinate system. These were used

**Table 1.** Cryptogam species recorded within 47 samples with information about their ecological status (KETNER-OOSTRA et al. [2012], slightly modified according to FALIŃSKI et al. [1993]), and their conservation status (CIEŚLIŃSKI et al. 2006). Abbreviations: pio = pioneer, hum = humicolous, aero-hyg = aero-hygrophyte, epi = epigeic epixylite

**Tabelle 1.** In den 47 Vegetationsaufnahmen vorkommende Kryptogamenarten, mit Informationen über ihr ökologisches Verhalten (KETNER-OOSTRA et al. [2012], leicht modifiziert nach FALIŃSKI et al. [1993]), sowie ihren Rote Liste und Schutzstatus (CIEŚLIŃSKI et al. 2006).

| Species name   | Abbreviation | Ecological status | Red List / National law protection status | Frequency (%) |
|--|--------------|-------------------|---|---------------|
| <i>Ceratodon purpureus</i>                           | Cer_pur      | pio               | -/-                                       | 47            |
| <i>Cetraria aculeata</i>                             | Cet_acu      | pio               | -/+                                       | 79            |
| <i>Cladonia cervicornis</i> ssp. <i>verticillata</i> | Cl_cerv      | pio               | -/-                                       | 45            |
| <i>Cladonia foliacea</i>                             | Cl_foli      | pio               | -/-                                       | 2             |
| <i>Cladonia mitis</i>                                | Cl_miti      | pio               | -/+                                       | 85            |
| <i>Placynthiella uliginosa</i>                       | Pla_ulig     | pio               | -/-                                       | 36            |
| <i>Polytrichum piliferum</i>                         | Pol_pili     | pio               | -/-                                       | 85            |
| <i>Racomitrium canescens</i>                         | Rac_cane     | pio               | -/-                                       | 9             |
| <i>Stereocaulon condensatum</i>                      | Ste_cond     | pio               | VU/+                                      | 15            |
| <i>Trapeliopsis granulosa</i>                        | Tra_gran     | pio               | -/-                                       | 40            |
| <i>Baeomyces rufus</i>                               | Bae_rufu     | hum               | -/-                                       | 2             |
| <i>Brachythecium albicans</i>                        | Bra_albi     | hum               | -/-                                       | 4             |
| <i>Cephaloziella divaricata</i>                      | Cep_diva     | hum               | -/-                                       | 9             |
| <i>Cladonia chlorophaea</i> s.lat.                   | Cl_chlo      | hum               | -/-                                       | 19            |
| <i>Cladonia coccifera</i> s.lat.                     | Cl_cocc      | hum               | -/-                                       | 36            |
| <i>Cladonia fimbriata</i>                            | Cl_fimb      | hum               | -/-                                       | 19            |
| <i>Cladonia furcata</i>                              | Cl_furc      | hum               | -/-                                       | 4             |
| <i>Cladonia glauca</i>                               | Cl_glau      | hum               | -/-                                       | 21            |
| <i>Cladonia macilenta</i> s.lat.                     | Cl_maci      | hum               | -/-                                       | 40            |
| <i>Cladonia phyllophora</i>                          | Cl_phyl      | hum               | -/-                                       | 45            |
| <i>Cladonia subulata</i>                             | Cl_subu      | hum               | -/-                                       | 57            |
| <i>Placynthiella oligotropha</i>                     | Pla_olig     | hum               | -/-                                       | 11            |
| <i>Polytrichum juniperinum</i>                       | Pol_juni     | hum               | -/-                                       | 2             |
| <i>Stereocaulon incrustatum</i>                      | Ste_incr     | hum               | EN/+                                      | 4             |
| <i>Cetraria islandica</i>                            | Cet_isla     | hum / aero-hyg    | VU/+                                      | 6             |
| <i>Cladonia cornuta</i>                              | Cl_corn      | hum / aero-hyg    | -/-                                       | 9             |
| <i>Cladonia deformis</i>                             | Cl_defo      | hum / aero-hyg    | -/-                                       | 23            |
| <i>Cladonia squamosa</i>                             | Cl_squa      | hum / aero-hyg    | -/-                                       | 2             |
| <i>Cladonia uncialis</i>                             | Cl_unci      | hum / aero-hyg    | -/-                                       | 47            |
| <i>Sciuro-hypnum oedipodium</i>                      | Sci_oedi     | hum / aero-hyg    | -/-                                       | 11            |
| <i>Pohlia nutans</i>                                 | Poh_nuta     | hum / aero-hyg    | -/-                                       | 11            |
| <i>Cladonia arbuscula</i>                            | Cl_arbu      | aero-hyg          | -/+                                       | 2             |
| <i>Cladonia crispata</i>                             | Cl_cris      | aero-hyg          | -/-                                       | 2             |
| <i>Cladonia gracilis</i>                             | Cl_grac      | aero-hyg          | -/-                                       | 19            |
| <i>Dicranum polysetum</i>                            | Dic_poly     | aero-hyg          | -/+                                       | 23            |
| <i>Dicranum scoparium</i>                            | Dic_scop     | aero-hyg          | -/+                                       | 17            |
| <i>Hylocomium splendens</i>                          | Hyl_sple     | aero-hyg          | -/+                                       | 2             |
| <i>Hypnum cupressiforme</i> s.lat.                   | Hyp_cupr     | aero-hyg          | -/-                                       | 13            |
| <i>Lophocolea heterophylla</i>                       | Lop_hete     | aero-hyg          | -/-                                       | 2             |
| <i>Pleurozium schreberi</i>                          | Ple_schr     | aero-hyg          | -/+                                       | 26            |
| <i>Placynthiella dasaea</i>                          | Pla_dase     | epi               | -/-                                       | 4             |
| <i>Hypogymnia physodes</i>                           | Hyp_phys     | epi               | -/-                                       | 2             |

**Table 2.** Summary of all environmental variables recorded within the study plots. Variables included in the pCCAs are indicated with a “+”.

**Tabelle 2.** Zusammenfassung aller innerhalb der Untersuchungsflächen gemessenen Umweltvariablen. Diejenigen, die nach der Entfernung der stark korrelierten Variablen in die beiden pCCA einbezogen wurden, sind mit “+” gekennzeichnet.

| Variable  | Abbreviation   | Unit                | Mean  | SD   | Median | Min   | Max   | Predictor |
|---|----------------|---------------------|-------|------|--------|-------|-------|-----------|
| Inclination                                       | INCL           | °                   | 2.66  | 2.70 | 3      | 0     | 11    | +         |
| Cover of tree layer                               | A_COV          | %                   | 1     | 5    | 0      | 0     | 30    | +         |
| Cover of shrub layer                              | B_COV          | %                   | 21    | 25   | 10     | 0     | 100   | +         |
| Common juniper cover in shrub layer               | JC_B           | %                   | 21    | 25   | 10     | 0     | 100   | -         |
| Share of plot area with visible dune processes    | DUNE_INT       | %                   | 29    | 41   | 0      | 0     | 100   | -         |
| Share of plot area disturbed by off-road vehicles | DIST_INT       | %                   | 57    | 35   | 64     | 0     | 100   | +         |
| Cover of visible open substrate                   | bare_soil      | %                   | 51    | 35   | 50     | 0     | 100   | -         |
| Cover of litter                                   | Litter         | %                   | 8     | 12   | 3      | 0     | 45    | -         |
| pH of upper soil horizon                          | pH             | -                   | 5.32  | 0.44 | 5.40   | 4.05  | 6.04  | -         |
| Electrical conductivity of upper soil horizon     | EC             | µS*cm <sup>-1</sup> | 35.5  | 26.7 | 27.9   | 15.6  | 142.6 | +         |
| Weight loss on ignition of upper soil horizon     | ORGANIC        | %                   | 0.67  | 0.61 | 0.54   | 0.14  | 3.13  | +         |
| Share of soil fractions > 1 mm                    | fr_coarse      | %                   | 0.50  | 0.49 | 0.33   | 0.02  | 2.30  | +         |
| Share of soil fractions 0.5–1 mm                  | fr_coarse_sand | %                   | 8.68  | 6.04 | 6.34   | 1.65  | 29.62 | -         |
| Share of soil fractions 0.25–0.5 mm               | fr_med_sand    | %                   | 55.35 | 6.83 | 56.50  | 20.83 | 67.20 | -         |
| Share of soil fractions 0.1–0.25 mm               | fr_fine_sand   | %                   | 34.07 | 9.82 | 33.61  | 19.51 | 76.47 | -         |
| Share of soil fractions < 0.1 mm                  | fr_fine        | %                   | 1.40  | 0.69 | 1.14   | 0.42  | 3.39  | +         |

as covariates in all of the performed multivariate analyses to remove the variance related only to the spatial arrangement of plots. In order to identify the most important ( $p < 0.05$ ) environmental variables (Table 2) influencing the analysed ground vegetation, we performed a partial Canonical Correspondence Analysis (pCCA) with manual forward selection of explanatory variables (1) for both herb and cryptogam layers, which were combined and transformed arithmetically according to TÜXEN & ELLENBERG (1937), and (2) for presence-absence data of cryptogams only. The significance of the species-environment relations was checked with a Monte Carlo test (499 runs). After the selection of the most relevant variables, a set of Generalised Additive Models (GAM) for calculated biodiversity and ecological indices of cryptogams (Table 3) was calculated together with XY(Z) plots according to recommended procedures (ŠMILAUER & LEPS 2014). They showed the response of the cryptogam species groups to disturbance intensity and shrub cover. The relevance of these gradients was supported by the results of the partial CCA. All analyses were performed using the software packages R and CANOCO 5 with Canodraw for Windows (ŠMILAUER & LEPS 2014, R CORE TEAM 2019).

## 4. Results

### 4.1 Cryptogam richness and ecological groups

A total of 29 vascular plant, 12 moss, 2 liverwort and 32 lichen species were recorded in the samples. The cryptogam species represented all ecological groups occurring on dune habitats (KETNER-OOSTRA et al. 2012). In the study 10 pioneer, 14 humicolous, 7 humicolous/aero-hygrophyte and 9 aero-hygrophyte species were found. One species was considered as an epigeic epiphyte and one as an epigeic epixylic species. Ten cryptogam species

**Table 3.** Summary of dependent variables used in the analysis (protection of species according to Polish law, red-listing according to CIEŚLIŃSKI et al. [2006], lichen ecological groups according to KETNER-OOSTRA et al. [2012] with slight modifications based on work by FALIŃSKI [1993]).

**Tabelle 3.** Zusammenfassung der abhängigen Variablen, die in der Analyse verwendet wurden (Schutzstatus nach polnischem Recht, Rote Liste nach CIEŚLIŃSKI et al. [2006], Flechtenökologische Gruppen nach KETNER-OOSTRA et al. [2012] mit leichten Modifikationen basierend auf Arbeiten von FALIŃSKI [1993]).

| Variable   | Abbreviation        | Mean | SD   | Median | Min | Max |
|--|---------------------|------|------|--------|-----|-----|
| No. of valuable (protected, red listed) lichen species | lich_val            | 1.91 | 0.86 | 2      | 0   | 3   |
| No. of bryophyte species                               | n_moss              | 2.60 | 2.13 | 2      | 0   | 9   |
| No. of lichen species                                  | n_lichen            | 6.77 | 4.34 | 7      | 0   | 15  |
| No. of all cryptogams (bryophytes, lichens)            | n_cryptogam_species | 9.36 | 5.70 | 9      | 0   | 20  |
| No. of pioneer species                                 | Pioneer n           | 4.43 | 2.14 | 5      | 0   | 8   |
| No. of humicolous species                              | Hum n               | 2.74 | 2.31 | 3      | 0   | 7   |
| No. of humicolous/aero-hygrophytes                     | Hum-Aero n          | 1.09 | 1.08 | 1      | 0   | 4   |
| No. of aero-hygrophytes                                | Aero n              | 1.06 | 1.62 | 0      | 0   | 6   |

protected by national law and three epigeic lichen species red-listed in Poland were found. Two of them, *Stereocaulon condensatum* and *S. incrustatum*, have the highest conservation status (Table 1). Within the study plots we recorded up to 20 cryptogam species (9.36 on average) and up to 15 species of lichens (6.77 on average) (Table 3).

#### 4.2 Main factors influencing the cryptogam community structure

The partial Canonical Correspondence Analysis performed for the combined herb and cryptogam layers revealed the disturbance intensity, shrub and tree cover, conductivity and gravel content to be relevant ( $p < 0.05$ ) variables influencing the community structure (Fig. 3, Table 4). The influence of soil organic matter content as well as fine soil particles content proved not to be relevant ( $p > 0.05$ ) in the applied model. The second partial Canonical Correspondence Analysis conducted only for presence-absence data in the cryptogam layer also indicated the disturbance intensity and tree and shrub cover as well as the share of the smallest fractions in the soil to be significant factors in the model (Fig. 4). In both cases the tree (first pCCA1:  $r = -0.6272$ , second pCCA1:  $r = 0.6561$ ) and shrub (first pCCA1  $r = -0.6865$ , second pCCA1  $r = 0.8101$ ) cover vectors were correlated with the first ordination axis whereas the disturbance intensity turned out to be unrelated to the first but connected with the second most important ordination axis (first pCCA2:  $r = 0.8690$ , second pCCA2:  $r = -0.7022$ ). This indicates that off-road disturbance and shrub density are the most important factors influencing cryptogams within the study area.

#### 4.3 Cryptogam biodiversity and response of ecological groups to disturbance level and shrub cover


The GAM fitting shrub cover and disturbance intensity to biodiversity indices and ecological groups of cryptogams produced significant results ( $p < 0.05$ ) throughout (Table 5). In the case of valuable lichen and pioneer cryptogam species numbers, the responses to disturbance intensity were strongly non-linear (non-linearity test  $p < 0.01$ ) (Table 5). According to XY(Z) plot analyses, the highest numbers of cryptogam species occurred in not or only slightly disturbed patches with medium shrub cover. The lowest numbers of species were



**Table 4.** Results of the Canonical Correspondence Analysis of the uncorrelated variables.


**Tabelle 4.** Ergebnisse der kanonischen Korrespondenzanalyse der unabhängigen Variablen.

| Variable        | Explains (%) | Contribution (%) | <i>pseudo-F</i> | <i>p</i>     |
|-----------------|--------------|------------------|-----------------|--------------|
| EC              | 11.6         | 25.8             | 5               | <b>0.002</b> |
| DIST_INT        | 11           | 24.5             | 5.3             | <b>0.002</b> |
| fr_skeleton_sum | 6.5          | 14.4             | 3.3             | <b>0.002</b> |
| INCL            | 4.1          | 9.2              | 2.2             | <b>0.024</b> |
| B_COV           | 3.9          | 8.6              | 2.1             | <b>0.02</b>  |
| A_COV           | 3.7          | 8.2              | 2               | 0.068        |
| fr_silt_clay    | 2.2          | 4.9              | 1.2             | 0.282        |
| ORGANIC         | 2            | 4.5              | 1.1             | 0.29         |



**Fig. 3.** Result of the partial Canonical Correspondence Analysis for combined herb and cryptogam layers. Only significant ( $p < 0.05$ ) factors influencing the species data are shown (B\_COV – Cover of shrub layer, DIST\_INT – Share of plot area disturbed by off-road vehicles, EC – Electrical conductivity of upper soil horizon, Fr\_coarse – Share of soil fractions larger than 1 mm, INCL – inclination). Abbreviations of cryptogam species names are explained in Table 1. Species names of vascular plants are abbreviated using the first three letters of the genus name and the first four letters of the species epithet.

**Abb. 3.** Ergebnis der kanonischen Korrespondenzanalyse (CCA) für Gefäßpflanzen und Kryptogamen. Nur signifikante ( $p < 0,05$ ) Faktoren werden gezeigt. Abkürzungen der Kryptogamenarten sind in Tabelle 1 erläutert. Abkürzungen der Gefäßpflanzenarten bestehen aus den ersten drei Buchstaben des Gattungsnamens und den ersten vier Buchstaben des Artnamens.



**Fig. 4.** Result of the partial Canonical Correspondence Analysis for the cryptogam layer using binominal (presence/absence) species data. Only significant ( $p < 0.05$ ) factors influencing the species data are shown (A\_COV – Cover of tree layer, B\_COV – Cover of shrub layer, DIST\_INT – Share of plot area disturbed by off-road vehicles, Fr\_fine – Share of soil fractions smaller than 0.1 mm). Abbreviations of cryptogam species names are explained in Table 1.

**Abb. 4.** Ergebnis der partiellen kanonischen Korrespondenzanalyse (pCCA) für die Kryptogamen mit binominaler Behandlung der Artendaten (Anwesenheit/Abwesenheit). Nur signifikante ( $p < 0,05$ ) Faktoren werden gezeigt. Abkürzungen der Kryptogamenarten sind in Tabelle 1 erläutert.


noted in sites with the highest disturbance levels and no shrub cover (Fig. 5a). Lichen richness was also highest in undisturbed patches, although with lower shrub cover (Fig. 5b). Bryophyte richness, on the other hand, clearly increased with increasing shrub cover, and at the same time showed a slight preference towards somewhat disturbed areas (Fig. 5c). The highest numbers of valuable lichen species occurred when the anthropogenic disturbance levels were medium and the shrub cover was below 40% (Fig. 5d).

The results of both multivariate analyses revealed that very high off-road intensity leads to the development of an initial stage of psammophilous grassland with active dune processes and low species diversity (very few species except for *Corynephorus canescens*) (Fig. 3 and 4). Medium disturbance is related with more species-rich pioneer communities with

**Tabelle 5.** Ergebnisse des verallgemeinerten Additivmodells, zur Untersuchung des Einflusses der Gehölzdeckung und Störungsintensität auf Biodiversitätsindizes und ökologische Kryptogamengruppen.


**Table 5.** Results of the Generalised Additive Model (GAM) fitting shrub cover and disturbance intensity to biodiversity indices and ecological groups of cryptogams.

| Index               | <i>F</i> | <i>p</i>  | Tests of non-linearity of response ( <i>p</i> ) |                |
|---------------------|----------|-----------|---|----------------|
|                     |          |           | B_COV   | DIST_INT       |
| lich_val            | 5.7      | 0.00091   | 0.27183   | <b>0.00409</b> |
| n_moss              | 32.1     | < 0.00001 | 0.10504   | 0.06103        |
| n_lichen            | 18.5     | < 0.00001 | <b>0.02309</b>                                  | 0.08864        |
| n_cryptogam_species | 27.1     | < 0.00001 | <b>0.01658</b>                                  | <b>0.03812</b> |
| Pioneer n           | 15.0     | < 0.00001 | <b>0.00255</b>                                  | <b>0.00013</b> |
| Hum n               | 13.5     | < 0.00001 | 0.11844   | 0.47264        |
| Hum-Aero n          | 9.8      | 0.00001   | 0.28905   | 0.29838        |
| Aero n              | 20.5     | < 0.00001 | 0.47838   | 0.09156        |



**Fig. 5.** XY(Z) plots showing the responses of cryptogams to disturbance intensity and shrub cover (GAM). **a)** total number of cryptogam species, **b)** total number of lichen species, **c)** total number of bryophytes, **d)** total number of valuable lichen species (protected and threatened according to CIEŚLIŃSKI et al. 2006).

**Abb. 5.** XY(Z)-Diagramme zum Einfluss der Störungsintensität und Gehölzdeckung (GAM) auf **a)** Gesamtartenzahl der Kryptogamen, **b)** Gesamtartenzahl der Flechten, **c)** Gesamtartenzahl der Moose, **d)** Gesamtartenzahl der Flechtenarten mit hohem Naturschutzwert.



**Fig. 6.** XY(Z) plots showing the responses of main ecological cryptogam groups (KETNER-OOSTRA et al. 2012, slightly modified according to succession models of FALIŃSKI et al. 1993) to disturbance intensity and shrub cover (GAM). **a)** total number of pioneer species, **b)** total number of humicolous species, **c)** total number of humicolous/aero-hygrophyte species, **d)** total number of aero-hygrophyte species.

**Abb. 6.** XY(Z)-Diagramme zum Einfluss der Störungsintensität und Gehölzdeckung (GAM) auf **a)** Gesamtzahl der Pionierarten, **b)** Gesamtzahl der humicolen Arten, **c)** Gesamtzahl der humicolen/aerohygrophytischen Arten, **d)** Gesamtzahl der aerohygrophytischen Arten.

*C. canescens* and *Spergula vernalis*, characterised by the presence of many pioneer cryptogams (Fig. 6a). The maximum pioneer species numbers occurred in patches of low or medium disturbance with low to medium shrub cover (5–45%). A decreasing impact of disturbance led to the gradual encroachment of humicolous species (Fig. 6b). The maximum numbers of cryptogam species belonging to this group were recorded within not or only slightly disturbed patches with medium shrub cover (15–60%). Humicolous/aero-hygrophytes (Fig. 6c) preferred undisturbed patches with high shrub cover (around 60%). The number of aero-hygrophytes increased gradually with a decrease in disturbance intensity and an increase of shrub cover, reaching its maximum where the shrub cover exceeded 70% (Fig. 6d).

## 5. Discussion

### 5.1 The interaction of ecological succession and off-road disturbance influences the cryptogam community structure

Military disturbance increases non-forest area (MILCHUNAS et al. 2000), reduces the cover of the herb layer and increases the exposed surface of the soil (SHAW & DIERSING 1990, PROSSER et al. 2000). The intensity of military use is an important factor competing with ecological succession. Excessive use of a training area can lead to a situation where the disturbance processes occur faster than the regeneration processes (HIRST et al. 2000, MILCHUNAS et al. 2000). On the other hand, some military training areas where disturbing human activities have ended suffer from rapid succession and require active management in order to save their valuable habitats (PRACH et al. 2000, JENTSCH et al. 2009). In our study partial Canonical Correspondence Analyses for the ground layer revealed that off-road disturbance and variables connected with ecological succession represented by shrub cover are the main independent factors explaining the structure of the whole dataset (Fig. 3). These are typical factors also described as influencing the vascular vegetation of other military training areas (e.g. HIRST et al. 2000, MILCHUNAS et al. 2000, PRACH et al. 2000). If only the presence of cryptogams was considered, the same factors turned out to be the most important (Fig. 4). This indicates that cryptogams respond to environmental factors similarly to vascular plants. What is more important, vectors representing both factors turned out to be perpendicular, suggesting that they are independent. It implicates that disturbance can influence the community at different successional stages, from an initial *Corynephorus* stand to dense *Juniperus* scrubland. Additionally, the observed high correlation between the presence of dune processes and the disturbance intensity rather than the successional stage suggests that off-road disturbance can trigger natural geomorphological processes typical for active inland dune habitats. In the presented study the increase of conductivity was also found to be a relevant factor following succession. It was probably connected with the rising concentrations of H<sup>+</sup> ions caused by the presence of organic acids in humus-enriched soils (Fig. 3). The presented relationships remain relevant within the whole dataset despite some heterogeneity of soils exhibited by variations in the dust, gravel or organic matter contents.

### 5.2 Habitat heterogeneity and intermediate disturbance promote cryptogam diversity

Landscape heterogeneity significantly influences the diversity of grassland ecosystems (JANIŠOVÁ et al. 2014, ZULKA et al. 2014). It is already known to positively correlate with the diversity of bryophytes and other organisms on military training sites (VANDERPOORTEN et al. 2005, DENGLER et al. 2020a). The presence of juniper scrub increases the small-scale heterogeneity of grasslands, mainly by providing valuable microhabitats for many terricolous cryptogam species (FALIŃSKI et al. 1993). It enhances their diversity primarily because it creates niches by increasing both shade and soil organic matter, which benefits late successional species in patches dominated by early successional ones (KALAIPOS & MÁZSA 2001, BOCH et al. 2016). Army vehicles are already known to create bare ground microhabitats (SHAW & DIERSING 1990, PROSSER et al. 2000). In the presented study the bare ground appeared to be positively correlated with off-road disturbance. It can be assumed that the interaction of these two factors increases the small-scale heterogeneity of the research site. Thirty-two terrestrial lichen species (including rare and endangered ones) recorded only within the sampled plots is a number fully comparable with Western European sites rich in

epigeic lichens, e.g. *Hulshorsterzand* in the Netherlands (KETNER-OOSTRA et al. 2012). This high gamma (landscape level) diversity of the studied area may be related to the landscape heterogeneity created by unevenly distributed juniper scrub and off-road disturbance.

At plot level (i.e. alpha) diversity, the highest richness of valuable lichens and pioneer species was found under moderate levels of disturbance (Fig 5d and 6a). High disturbance intensity increased the bare soil area and decreased the number of cryptogam species (Fig. 5). Cryptogams are rather scarce in habitats directly affected by the dune process, and in such habitats only *Corynephorus canescens* remains relatively vital (KETNER-OOSTRA et al. 2012), an observation that was also made in our study area. On the other hand, it is known that pioneer species can only occur in areas with bare soil patches available for colonisation (COLE 1990, BELNAP 1993, LALLEY & VILES 2008). Ecological succession leads to the gradual disappearance of psammophilous grassland communities and the formation of initial pine forest communities (FALIŃSKI 1993). Within undisturbed habitat patches with high shrub cover, the vanishing of pioneer species and an increase in other ecological groups of cryptogams were registered (Fig. 6). In the presented study the presence of juniper scrub enhanced the species number of all registered groups of cryptogams (Fig. 5). The richness optima of ecological groups (Fig. 6) were connected consecutively with rising shrub cover. The aero-hygrophytes, which are already forest species, were the last to appear when the habitat was no longer open.

The most important factor maintaining high lichen species numbers was variable disturbance intensity connected with high microhabitat diversity representing all of the main successional stages. Within moderately disturbed patches disturbance and regeneration processes were probably somewhat balanced. This ensured the continuation of ecological niches and short-distance colonisation as the most sufficient lichen dispersal occurs within less than a few metres (HEINKEN 1999). One can assume that off-road disturbance additionally helped to move cryptogam fragments and inoculate new niches. Thus, many new habitat patches could be quickly colonised. The observed importance of moderate levels of disturbance for the maintenance of high species richness and conservation of some valuable species agrees well with the “intermediate disturbance hypothesis” (CONNELL 1978, WILKINSON 1999).

### **5.3 Can off-road vehicle use maintain the high terrestrial cryptogam diversity of military training areas?**

The conservation of endangered species in operating military training areas can generate many conflicts (JENNI et al. 2012), and it is difficult to balance the necessary military use, not always legal off-road use and conservation of natural values at one location, even within the borders of a National Park. Acid inland dunes are temporary habitats of high conservation value and need management to persist. One recently applied management practice of open acid dune habitats is tree and shrub removal and grazing management to maintain open landscape structures. The restoration of more altered habitats requires expensive topsoil removal (KETNER-OOSTRA et al. 2012). According to the results of the presented study, off-road disturbance may also be used as an inexpensive alternative to these activities. Spatially diverse off-road disturbance of low to medium intensity may be used in habitat management, whereas intensive off-road use may be implemented as a tool for the restoration of the initial stage of the habitat with active dune processes.

## 6. Conclusions

Increasing shrub cover related to successional processes and off-road activity were the two most important factors influencing cryptogam richness and community composition on acid inland dune habitats in the studied military training area. Pioneer cryptogam species and valuable lichens benefited from low and medium off-road disturbance. At the same time, other species were instead associated with higher successional stages found in undisturbed patches with rising juniper scrub cover. We interpret the growing intensity of disturbance as a factor leading to the redevelopment of the initial stage of the habitat, where dune processes can also occur. Diverse off-road activity locally reversed successional progress and thus was a positive factor promoting the coexistence of all ecological groups of cryptogams connected with the acid inland dune environment. In our opinion, off-road vehicle activity may to some extent be used as an inexpensive nature conservation tool in order to maintain epigeic, open-habitat cryptogam species. However, we are aware of the limitations of our study, in which we were not able to fully disentangle the impact of off-road disturbance from the naturally occurring one, especially as they seem to be closely related. Thus, we think that our results should be verified in a system in which both types of disturbance can be better controlled (e.g. in an experimental study).

### Erweiterte deutsche Zusammenfassung

**Einleitung** – Auf Binnendünen wachsende, psammophile Pflanzengesellschaften mit Wachholder (*Juniperus communis*) gehören zu den am meisten bedrohten Lebensraumtypen in Mitteleuropa. Einst standen sie im Zusammenhang mit Störungen, die durch die traditionelle Land- und Forstwirtschaft verursacht wurden (Schafweiden, Brennen, Streunutzung, Abholzen von Wäldern). Wird die Nutzung aufgegeben, sind diese Vegetationstypen einer sekundären Sukzession unterworfen. Diese führt zum Verlust der charakteristischen Artenkombination und der typischen Biodiversität. Ein wichtiger Bestandteil dieser typischen Vegetation sind Kryptogamen (Moose und Flechten; WOLFF et al. 2017, DENGLER et al. 2020a). In dieser Studie untersuchten wir, ob die in letzter Zeit zunehmende Off-Road-Aktivität mit Geländefahrzeugen in Dünengebieten solche Lebensräume und ihre Biodiversität erhalten kann oder stattdessen zur weiteren Verschlechterung der Habitatqualität führt. Um diese Frage zu beantworten, untersuchten wir die Kryptogamendiversität in einer durch Off-Road-Aktivitäten gestörten Dünenlandschaft eines selten genutzten militärischen Übungsgeländes im Binnenland Zentralpolens.

**Untersuchungsgebiet** – Die Untersuchung wurde auf einem militärischen Übungsgelände namens Grochalskie Piachy (ca. 250 ha) in Zentralpolen (52°24'15"N 20°37'05"E), 22 km nordwestlich der Hauptstadt Warschau, durchgeführt. Das Gebiet liegt im vorzeitlichen Weichseltal auf tiefem Flusssand, der durch den äolischen Prozess im frühen Holozän umgestaltet wurde. Es ist Teil eines der größten und am besten entwickelten Dünengürtel in Europa (ZGORZELSKI & PAWŁOWSKA 2003) und entstand durch Abholzung im späten 19. und 20. Jahrhundert. Das Gebiet umfasst drei offene Sandfelder mit aktiven Dünenprozessen, die durch militärische Aktivitäten mobilisiert wurden. Die offenen Flächen (ca. 100 ha) sind überwiegend mit Pflanzengesellschaften des *Corynephorion canescentis* Klika 1931 bedeckt. Der nördliche Teil des Geländes ist noch immer durch die regelmäßige Neutralisierung alter, nicht explodierter Bomben aus dem Zweiten Weltkrieg stark gestört. Der restliche Teil des Gebiets, in dem die Studie durchgeführt wurde, ist heute nicht mehr militärisch genutzt. Das gesamte nicht bewaldete Gebiet des Geländes wird momentan Geländefahrzeugliebhabern genutzt.

**Material und Methoden** – Wir erstellten 47 Vegetationsaufnahmen und erfassten eine Reihe von gemessenen oder geschätzten Variablen, wie Bodenparameter und Störungsintensität sowie Faktoren die im Zusammenhang mit der Sukzession stehen, wie die Gehölzdeckung. Zur Auswahl der unabhängigen erklärenden Variablen verwendeten wir Pearson-Korrelationen. Mittels partieller Kanonischer

Korrelationen (CCAs) ermittelten wir die wichtigsten Bodenvegetation-beeinflussenden Variablen. Anhand von GAM-Modellen untersuchten wir den Einfluss der Störungsintensität und der Gehölzbedeckung auf die Kryptogamendiversität und weitere ökologische Kryptogamen-Indizes.

**Ergebnisse** – Die Struktur der psammophilen Kryptogamengemeinschaften wurde vor allem durch Störungen von Geländefahrzeugen und der Gehölzbedeckung beeinflusst (Abb. 3, 4). Die höchste Störungsintensität durch Geländefahrzeuge führte zu einer Transformation des Lebensraums zum Initialstadium mit aktiven Dünenprozessen, in ungestörten Bereichen entwickelte sich hingegen ein Kiefernwald im Laufe der Sukzession. Die höchste Anzahl von Flechten- und Moosarten trat innerhalb nicht oder nur leicht gestörter Lebensräume auf. Flechten bevorzugten jedoch eine geringere Gehölzbedeckung als Moose (Abb. 5). Dennoch profitierten Pionier- und typische Arten psammophiler Vegetationstypen, zu denen auch einige bedrohte Flechtenarten gehören, von einer geringen bis mittleren Störungsintensität (Abb. 6).

**Diskussion** – Vielfältige und räumlich komplexe Off-Road-Aktivitäten mit Geländefahrzeugen zusammen mit einer variablen Gehölzbedeckung waren positive Faktoren, die die Koexistenz aller ökologischen Kryptogamengruppen der sauren Binnendünen förderten, womöglich aufgrund der erhöhten Lebensraumheterogenität und Anzahl Mikrohabitate. Auf moderat gestörten Flächen waren die Störungs- und Regenerationsprozesse wahrscheinlich eher ausgeglichen, was sich positiv auf den Erhalt wertvoller Flechten- und Pionierartengruppen auswirkte. Die Ergebnisse stützen die "mittlere Störungshypothese" (CONNELL 1978, WILKINSON 1999). Unserer Meinung nach können Off-Road-Aktivitäten mit Geländefahrzeugen bis zu einem gewissen Grad als ein kostengünstiges Naturschutzinstrument eingesetzt werden, um offene, Kryptogamen-reiche Dünenlebensräume zu erhalten.

## Acknowledgements

We would like to thank Paweł Wojdał and Aleksandra Grabowska for their assistance in the fieldwork. We are also thankful to Aiko Huckauf for very thorough linguistic corrections.


## Author contribution


PZ – fieldwork, laboratory work, literature survey, statistics, writing,

LK – fieldwork, laboratory work, literature survey, statistics, writing,

MW – laboratory work, literature survey, writing

## ORCIDiDs

Piotr Zaniewski  <https://orcid.org/0000-0002-0792-9854>

Łukasz Kozub  <https://orcid.org/0000-0002-6591-8045>

## Supplements

**Additional supporting information may be found in the online version of this article.**

**Zusätzliche unterstützende Information ist in der Online-Version dieses Artikels zu finden.**

**Supplement E1.** Original vegetation data used in the analyses.

**Anhang E1.** Ursprüngliche Vegetationsdaten, die in den Analysen verwendet wurden.

**Supplement E2.** Raw environmental data used in the analyses.

**Anhang E2.** In den Analysen verwendete Rohdaten der Umweltvariablen.



## References

- ADAMSKA, E. & DEPTUŁA, M. (2015): Materials for biota of lichens and lichenicolous fungi in the military area near Toruń, Poland. – *Ecol. Quest.* 21: 45–53.
- ANDRZEJEWSKA, A. (2003): Klimat. – In: ANDRZEJEWSKI, R. (Ed.): *Kampinoski Park Narodowy*. Vol. 1: 41–68. *Kampinoski Park Narodowy, Izabelin*.
- BARKMAN, J.J., DOING, H. & SEGAL, S. (1964): Kritische Bemerkungen und Vorschläge zur quantitativen Vegetationsanalyse. – *Acta Bot. Neerl.* 13: 394–419.
- BARKMAN, J.J., (1985): Geographical variation in associations of juniper scrub in the central European plain. – *Vegetatio* 59: 67–71.
- BELNAP, J. (1993): Recovery rates of cryptobiotic crusts: inoculant use and assessment methods. – *Great Basin Nat.* 53: 89–95.
- BOCH, S., PRATI, D., SCHÖNING, I. & FISCHER M. (2016): Lichen species richness is highest in non-intensively used grasslands promoting suitable microhabitats and low vascular plant competition. – *Biodivers. Conserv.* 25: 225–38.
- BORCARD, D., LEGENDRE, P. & DRAPEAU, P. (1992): Partialling out the spatial component of ecological variation. – *Ecology* 73: 1045–1055.
- BRAUN-BLANQUET, J. (1928): *Pflanzensoziologie. Grundzüge der Vegetationskunde*. – *Biologische Studienbücher* 7, Springer, Berlin: 330 pp.
- BROOME, A., LONG, D., WARD, L.K. & PARK K.J. (2017): Promoting natural regeneration for the restoration of *Juniperus communis*: a synthesis of knowledge and evidence for conservation practitioners. – *Appl. Veg. Sci.* 20: 397–409.
- CHOJNACKA, J., CYZMAN, W., NIENARTOWICZ, A. & DEPTUŁA, M. (2010): Variability of structure and directions in the development of heaths and psammophilous grasslands within the artillery range near Toruń. – *Ecol. Quest.* 12: 87–125.
- CIEŚLIŃSKI, S., CZYŻEWSKA, K. & FABISZEWSKI, J. (2006): Red list of lichens in Poland. In: MIREK Z., ZARZYCKI K., WOJEWODA W. & SZELĄG Z. (Eds): *Red list of plants and fungi in Poland*: 71–90. Polish Academy of Science, Kraków.
- COLE, D.N. (1990): Trampling disturbance and recovery of cryptogamic soil crusts in Grand Canyon National Park. – *Great Basin Nat.* 50: 321–325.
- CONNELL, J.H. (1978): Diversity in tropical rain forests and coral reefs. – *Science* 199: 1302–1310.
- DAMSHOLT, K. (2002): *Illustrated Flora of the Nordic liverworts and hornworts*. Nordic Bryological Society, Lund, Sweden: 837 pp.
- DEGLER, J., BIURRUN, I., BOCH, S., DEMBICZ, I. & TÖRÖK, P. (2020a): Grasslands of the Palaearctic biogeographic realm: introduction and synthesis. – In: GOLDSTEIN, M.I., DELLASALA, D.A. & DI PAOLO, D.A. (Eds.): *Encyclopedia of the world's biomes. Vol. 3: Forests – trees of life. Grasslands and shrublands – sea of plants*: 617–637. Elsevier, Amsterdam.
- DEGLER, J., BIRGE, T., BIRUNN, H.H., RAŠOMAVIČIUS, V., RÜSIŃA, S. & SICKEL, H. (2020b): Grasslands of Northern Europe and the Baltic States. – In: GOLDSTEIN, M.I., DELLASALA, D.A. & DI PAOLO, D.A. (Eds.): *Encyclopedia of the world's biomes. Vol. 3: Forests – trees of life. Grasslands and shrublands – sea of plants*: 689–702. Elsevier, Amsterdam.
- DINGOVÁ, A. (2010): Vplyv manažmentových opatrení na diverzitu terestrických lišajníkov vybraných dopadových plôch vo vojenskom výcvikovom priestore Záhorie (The impact of the management actions on diversity of lichens in Military Training Area Záhorie [in Slovak]). – *Vedecký Obzor – Scientific Horizon* 2: 17–26.
- EUROPEAN COMMISSION (2013): *Interpretation manual of European Union habitats*. – EUR 28. April 2013. DG Environment. *Nature ENV B.3*: 144 pp.
- FALIŃSKI, J.B., CIEŚLIŃSKI, S. & CZYŻEWSKA, K. (1993): *Dynamic-floristic atlas of Jelónka*. – *Phytocoenosis* 5, Suppl. *Cart. Geobot.* 3, Warszawa-Białowieża: 1–139.
- FITTER, H. & JENNINGS, R.D. (1975): The effects of sheep grazing on the growth and survival of seedling junipers (*Juniperus communis* L.). – *J. Appl. Ecol.* 12: 637–642.
- GARCIA, D., ZAMORA, R., HÓDAR, J.A. & GÓMEZ, J.M. (1999): Age structure of *Juniperus communis* L. in the Iberian peninsula: Conservation of remnant populations in Mediterranean mountains. – *Biol. Conserv.* 87: 215–220.
- GUGNACKA-FIEDOR, W. & ADAMSKA, E. (2010): The preservation state of the flora and vegetation of the artillery range near the city of Toruń. – *Ecol. Quest.* 12: 75–86.

- HIRST, R.A., PYWELL, R.F. & PUTWAIN, P.D. (2000): Assessing habitat disturbance using an historical perspective: The case of Salisbury Plain military training area. – *J. Environ. Manag.* 60: 181–193.
- IUCN (1996): Tanks and Thyme – Biodiversity in Former Soviet Military Areas in Central Europe. – IUCN, Gland, Switzerland, and Cambridge, UK: 136. pp.
- JANIŠOVÁ, M., MICHALCOVÁ, D., BACARO, G. & GHISLA, A. (2014): Landscape effects on diversity of semi-natural grasslands. – *Agr. Ecosys. Environ.* 182: 47–58.
- JENNI, G.D.L., PETERSON, M.N., CUBBAGE, F.W. & JAMESON, J.K. (2012): Assessing biodiversity conservation conflict on military installations. – *Biol. Conserv.* 153: 127–133.
- JENTSCH, A., FRIEDRICH, S., STEINLEIN, T., BEYSCHLAG, W. & NEZADAL W. (2009): Assessing conservation action for substituting of missing dynamics on former military training areas in Central Europe. – *Restor. Ecol.* 17: 107–116.
- JUŠKIEWICZ, B. & ENDLER, Z. (2000): Fitocenozy *Diantho-Armerietum* na byłym poligonie wojskowym Muszaki Jagorzewo (Równina Mazurska) (Phytocenoses of *Diantho-Armerietum* within the former military training area Muszaki Jagorzewo [Masurian Plain])[in Polish]. – *Fragm. Flor. Geobot. Pol.* 7: 159–165.
- KALAPOŠ, T. & MÁZSA, K. (2001): Juniper shade enables terricolous lichens and mosses to maintain high photochemical efficiency in a semiarid temperate grassland. – *Photosynthetica* 39: 263–268.
- KASARI, L., GAZOL, A., KALWIJ, H.M. & HELM, A. (2013): Low scrub cover in alvar grasslands increases small-scale diversity by promoting the occurrence of generalist species. – *Tuexenia* 33: 293–308.
- KETNER-OOSTRA, R., APTROOT, A., JUNGERIUS, P.D. & SÝKORA, K.V. (2012): Vegetation succession and habitat restoration in Dutch lichen-rich inland drift sand. – *Tuexenia* 32: 245–268.
- KRAWCZYK, R., ZUBEL, R. & KOMSTA, Ł. (2019): Military training areas and vegetation – the effect of explosion craters on species diversity along a moisture gradient. – *Pol. J. Ecol.* 67: 194–205.
- KUTIEL, P., EDEN, E. & ZHEVELEV, Y. (2000): Effect of experimental trampling and off-road motorcycle traffic on soil and vegetation of stabilized coastal dunes, Israel. – *Environ. Conserv.* 27: 14–23.
- LALLEY, J.S. & VILES, H.A. (2008): Recovery of lichen-dominated soil crusts in a hyper-arid desert. – *Biodivers. Conserv.* 17: 1–20.
- MATUSZKIEWICZ, J.M. (2015): Roślinność rzeczywista pasów wydmowych Kampinoskiego Parku Narodowego. Mapa fitosocjologiczna, – Kampinoski Park Narodowy (Actual vegetation of the dune belts of Kampinos National Park. – Phytosociological Map)[in Polish]. – Izabelin.
- MILCHUNAS, D.G. (2000): Plant community structure in relation to long-term disturbance by mechanized military maneuvers in a semiarid region. – *Environ. Manag.* 25: 525–539.
- MIREK, Z., PIĘKOŚ-MIRKOWA, H., ZAJĄC, A. & ZAJĄC, M. (2002): Flowering plants and pteridophytes of Poland. A checklist. – W. Szafer Institute of Botany, Polish Academy of Science, Kraków: 442 pp.
- MOTIEJUNAITE, J. (1996): Mycological and lichenological investigations in the former soviet military forestries. Lichens and allied fungi. – *Bot. Lith.* 2: 343–364.
- OCHYRA, R. (1992): Czerwona lista mchów zagrożonych w Polsce (Red list of threatened mosses in Poland). – In: ZARZYCKI, K., WOJEWODA, W. & HEINRICH, Z. (Eds.): Lista roślin zagrożonych w Polsce (List of threatened plants in Poland) [in Polish with English summary]: 79–86. Polska Akademia Nauk, Instytut Botaniki im. W. Szafera, Kraków.
- ORANGE, A., JAMES, P.W. & WHITE, F.J. (2001): Microchemical methods for the identification of lichens. – British Lichen Society, London: 101 pp.
- PRACH, K., BUFKOWÁ, I., ZEMEK, F., HEŘMAN, M. & MAŠKOVÁ, Z. (2000): Grassland vegetation in the former military area Dobrá Voda, the Šumava National Park. – *Silva Gabreta* 5: 101–112.
- PROSSER, C.W., SEDIVEC, K.K. & BARKER, W.T. (2000): Tracked Vehicle effects on Vegetation and Soil Characteristics. – *J. Range Manag.* 53: 666–670.
- R CORE TEAM (2019): R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. – URL: <http://www.r-project.org/index.html>.
- RICKARD, C.A., MCLACHLAN, A. & KERLEY, G.I.H. (1994): The effects of vehicular and pedestrian traffic on dune vegetation in South Africa. – *Ocean Coast. Manag.* 23: 225–247.
- ROMAHN, K.S. (1998): Die Vegetation der Krempfer und Nordoer Heide – Vegetationskundliche Untersuchungen auf einem Standortübungsplatz der Bundeswehr. – *Mitt. Arbeitsgem. Geobot. Schlesw.-Holst. Hambg.* 54: 1–92.

- SCHWABE, A., SÜSS, K. & STORM, C. (2013): What are the long-term effects of livestock grazing in steppic sandy grassland with high conservation value? Results from a 12-year field study. – *Tuexenia* 33: 189–212.
- SHAW, R.B. & DIERSING, V.E. (1990): Tracked vehicle impacts on vegetation at the Pinon Canyon maneuver site, Colorado, USA. – *J. Environ. Qual.* 19: 234–243.
- SMITH, A.J.E. (2004): *The Moss Flora of Britain and Ireland*. – Cambridge University Press, Cambridge: 1012 pp.
- SMITH, C.W., APTROOT, A., COPPINS, B.J., FLETCHER, A., GILBERT, O.L., JAMES, P.W. & WOLSELEY, P.A. (2009): *The lichens of Great Britain and Ireland*. – British Lichen Society. London: 1046 pp.
- ŠMILAUER, P. & LEPŠ, J. (2014): *Multivariate analysis of ecological data using CANOCO 5*. – Cambridge University Press, Cambridge: 362 pp.
- TÜXEN R. & ELLENBERG H. (1937): Der systematische und ökologische Gruppenwert. Ein Beitrag zur Begriffsbildung und Methodik der Pflanzensoziologie. – *Mitt. Florist.-Soziol. Arbeitsgem.* 3: 171–184.
- VANDERPOORTEN, A., SOTIAUX, A. & ENGELS, P. (2005): A GIS-based survey for the conservation of bryophytes at the landscape scale. – *Biol. Conserv.* 121: 189–194.
- VERHEYEN, K., SCHREURS, K., VANHOLEN, B. & HERMY, M. (2005): Intensive management fails to promote recruitment in the last large population of *Juniperus communis* (L.) in Flanders (Belgium). – *Biol. Conserv.* 124: 113–121.
- VERHEYEN, K., ADRIAENSSENS, S., GRUWEZ, R., MICHALCZYK, M., WARD, L.K., ROSSEEL, Y., VAN DEN BROECK, A. & GARCIA, D. (2009): *Juniperus communis*: victim of the combined action of climate warming and nitrogen deposition? – *Plant Biol.* 11: 49–59.
- WARD, L.K. (1982): The conservation of juniper: longevity and old age. – *J. Appl. Ecol.* 19: 917–928.
- WARREN, S.D., HOLBROOK, S.W., DALE, D.A., WHELAN, N.L., ELYN, M., GRIMM, W. & JENTSCH, A. (2007): Biodiversity and the heterogenous disturbance regime on military training lands. – *Restor. Ecol.* 15: 606–612.
- WARREN, S.D. & BÜTTNER, R. (2008): Active military training areas as refugia for disturbance-dependent endangered insects. – *J. Insect Conserv.* 12: 671–676.
- WILKINSON, D.M. (1999): The disturbing history of intermediate disturbance. – *Oikos* 84: 145–147.
- WOLFF, A., GILHAUS, K., HÖLZEL, N. & SCHNEIDER S. (2017): Status and restoration potential of heathlands and sand grasslands in the southwest of Luxembourg. – *Tuexenia* 37: 179–200.
- ZGORZELSKI, M. & PAWŁOWSKA, T. (2003): Geomorfologia. – In: ANDRZEJEWSKI, R. (Ed.) *Kampinoski Park Narodowy*, Vol. 1: 87–95. *Kampinoski Park Narodowy*, Izabelin.
- ZULKA, K.P., ABENSPERG-TRAUN, M., MILASOWSZKY, N. ... ZECHMEISTER, H. (2014): Species richness in dry grassland patches of eastern Austria: A multi-taxon study on the role of local, landscape and habitat quality variables. – *Agr. Ecosyst. Environ.* 182: 25–36.





# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Tuexenia - Mitteilungen der Floristisch-soziologischen Arbeitsgemeinschaft](#)

Jahr/Year: 2020

Band/Volume: [NS\\_40](#)

Autor(en)/Author(s): Zaniewski Piotr T., Kozub Lukasz [Åukasz], Wierzbicka Malgorzata [MaÅ,gorzata]

Artikel/Article: [Intermediate disturbance by off-road vehicles promotes endangered pioneer cryptogam species of acid inland dunes Moderate Störungen durch Geländefahrzeuge fördern gefährdete Pionier-Kryptogamenarten in sauren Binnendünen 479-497](#)