

Thymi bulgarici.

Auctore: **Iv. K. Urumov** (Sofia).

Thymus pannonicus ALL. var. *Hackelianus* (OP.). — In graminosis, et in collinis aridis prope Stanimaka, distr. Plovdiv (det. RONNIGER).

Th. Marschallianus WILLD. — In pratis prope Sofia, Trnovo, Kazanlak, ad pagum Bela slatina, Klementino, Bučino, m. Rhodope, in collibus aridis prope opp. Pleven (det. RONN.). — *ssp. citratus* OPIZ. — In pratis et in graminosis ad pagum Klementino, Bivolan, distr. Pleven. — *ssp. Wondracekianus* OPIZ. — In graminosis siccis prope oppidum Pleven (det. RONN.). — *ssp. Radoslavovii* RONN. et URUM. — In lapidosis aridis ad oppida Pleven et Trnovo. Mit feinen Stieldrüsen an den Blütenstielen und Kelchen, sonst wie bei *ssp. Wondracekianus* OPIZ. — *ssp. arenicolus* H. BR. — In pascuis et in graminosis ad opp. Pleven, ad pagum Rakovski (det. RONN.). — var. *Ljulinensis* (DEG. et URUM.) RONN. — In graminosis siccis m. Ljulin planina distr. Sofia. — var. *bulgaricus* (DOM. et PODP.) RONN. — In graminosis ad opp. Tatarpazardžik, Rhodope, pr. Banki.

Th. rumelicus DEG. — Ad pagum Papazlii distr. Plovdiv, m. Rhodope.

Th. brachyodon BORB. — In graminosis ad pagum Suhindol, distr. Trnovo.

Th. montanus W. K. — In pratis et in graminosis ad opp. Sofia, Pleven, Lukovit, vicum Banki, Boruštica, Karaš. — var. *amplificatus* SCHUR. — Ad opp. Karlovo et Kalofer distr. Plovdiv. — var. *micranthus* (WIERZB.) HEUFF. — In graminosis ad pagum Banki, Gornja banja et in m. Ljulin planina, distr. Sofia. — *ssp. danubialis* (SIMK.) RONN. — In graminosis siccis ad pagum Plačkovica, Boškovei, Krastec, distr. Trnovo; p. Boruštica non procul ab opp. Kazanlik (det. RONN.)

Th. noricus RONN. — In pratis et in graminosis ad pagum Plačkovica, distr. Trnovo.

Th. Borbásii H. BR. — Ad oppidum Stanimaka, distr. Plovdiv.

Th. heterotrichus GRIS. var. *leucanthus* DEG. et URUM. — In m. Rilo et Osogovo.

Th. Toševii VEL. — In graminosis m. Sakar planina, Rilo, Stara-planina, Stanimaka, Kotel, Klisura. — *ssp. Degenii* (H. BR.) RONN. — In graminosis m. Rilo, Rhodope, Pirin (det. RONN. et DEGEN). — *ssp. Raikovii* RONN. et URUM. — In herbisidiosis saxosis m. Pirin. Kelch ebenso klein, wie bei *Th. Toševii* VEL., Blattform ebenfalls wie bei diesem, aber obere Kelchzähne nur

sehr kurz papillös, wie bei *Th. glaucus* FRIV. Blätter beiderseits spärlich behaart. — *var. longifrons* RONN. — In graminosis siccis m. Rhodope. — *var. cinerascens* (VEL.) RONN. — Ad pedem m. Balkan prope opp. Karlovo, m. Rilo, Čepelare, Trnovo-Seimen, Brestovica, distr. Plovdiv. — *var. hirtiformis* VEL. — In graminosis prope opp. Vasiliko, distr. Burgas (det. RONN.)

Th. macedonicus (DEG. et URUM.) RONN. [*Th. Kosteleckyanus* var. *maced.* DEG. et URUM. in Journ. Acad. bulg. XV. (1923.):189]. — In pascuis Čepelarski Karlak m. Rhodope, prope Kistendil, ad pagum Kostinbrod non procul ab oppido Sofia.

Th. doiranensis RONN. In graminosis siccis prope Doiran.

Th. gracilis VEL. (Reliquiae Mrkvickianae. 1922:25). — In graminosis et in collinis aridis prope oppidum Varna.

Th. moesiacus VEL. — In graminosis aridis m. Rhodope, ad pagum Suhindol et Panlikene, distr. Trnovo.

Th. Callieri BORBÁS *typicus*. — In pratis et in graminosis prope oppidum Trnovo, Varna, Lom, Kazanlak, Nova-Zagora, in m. Rhodope. — *var. microcalyx* DEG. et URUM. — In graminosis siccis prope oppidum Trnovo. — *var. Urumovii* VEL. — In pratis et in graminosis prope opp. Eski Džumaja, Trnovo, Popovo, pagum Aboba, Papazlii, Osikovica.

Th. zygioides GRIS. — In graminosis et in collinis aridis prope Stanimaka, Varna, Haskovo, pagum Šeitan küpri.

Th. longidentatus (DEG. et URUM. pro var. *Th. zygioidis* in Magy. Bot. Lapok 1920:20). — In graminosis et in collinis siccis prope Pazardžik, Stanimaka, distr. Plovdiv.

Th. Kümmerleanus DEG. — In lapidosis aridis m. Pirin (det. RONN.)

Th. Aivalii H. BR. — In graminosis et in collinis aridis prope opp. Malko Trnovo et pagum Čausovo, distr. Plovdiv.

Th. glaucus FRIV. — In lapidosis aridis ad pagum Kavaklii distr. Plovdiv. — *var. hellesponticus* DEG. et URUM. — In lapidosis aridis Thraciae merid.-orientalis, ad Rodosto.

Th. Kernerii BORB. — In graminosis m. Balkan, Rhodope et Rilo (det. RONN.). *var. epitrichus* BORB. — In pascuis m. Balkan (det. RONN.)

Th. praecox OP. — In graminosis prope opp. Trnovo, Ruse, Stanimaka, pagum Tekira, Svoje, Papazlii, Rila.

Th. spathulatus OPIZ. — In pascuis m. Rhodope, Kostenski Balkan (det. DEGEN).

Th. glabrescens WILLD. — In pratis ad pagum Hisar, distr. Plovdiv (det. RONN.) — *var. stenophyllus* OPIZ. — Ad pag. Rakovski, Borisovgrad, distr. Plovdiv.

Th. austriacus BERNH. — In saxosis ad Čauški Balkan m. Rhodope, m. Belašica (det. RONN.)

Th. albanus H. BR. — In graminosis m. Rilo (det. RONN.).

— *f. gracilior* RONN. — In pascuis et in graminosis m. Vitoša et Rhodope (det. RONN.)

Th. thracicus VEL. — In graminosis siccis et in collinis prope opp. Sofia, Plovdiv, Loveč, Stanimaka.

Th. Stribrnyi VEL. — In herbis saxosis m. Konevo planina distr. Kistendil.

Th. euxinus (RONN.) — Ad oppidum Seozopol, distr. Burgas.

Th. longidens VEL. — In graminosis siccis m. Rhodope, Gabrovski Balkan, Alibotuš, Kozek planina. — *f. trnovensis* DEG. et URUM. — In lapidosis prope opp. Trnovo.

Th. alšariensis RONN. — In collinis aridis m. Rhodope ad pag. Čausevo.

Th. ocheus HELDR. et SART. — In m. Rhodope, Pirin et Alibotuš (det. RONN.).

Th. Jankae ČEL. — In graminosis m. Stara planina, Rhodope, Rui, Rilo et ad opp. Šumen. (det. DEGEN). — *ssp. Škorpilii* VEL. — In graminosis siccis ad pagum Madara distr. Šumen. — *var. hirsutior* FORMÁNEK. — In pratis ad m. Belašica, — *var. subacicularis* BORB. — In pratis ad pagum Dragoman distr. Sofia.

Th. Kapelae BORB. — In pascuis m. Gabrovski Balkan (det. RONN.). — *f. glabrior* NEIČEV. — In m. Gabrovski Balkan.

Th. balcanus BORB. — In m. Stara planina, Vitoša, Rilo, Rhodope, Pirin et Osogovo (det. RONN.). — *f. suprapilosus* DEG. In pascuis m. Etropolski Balkan (det. DEGEN). — *var. albifrons* VEL. Ad pagum Osengovo. — *ssp. glabrilateratus* (BORB.) RONN. — In pascuis m. Trojanski Balkan (det. RONN.).

Th. Nikolovii DEG. et URUM. — In Thracia meridionali ad oppidum Rodosto.

Th. boeoticus H. BR. — In herbis saxosis m. Rhodope.

Th. corthiaticus ADAMOVIĆ. — In graminosis siccis prope oppidum Gorna džumaia, Nevrekop (det. RONN.)

Th. atticus ČEL. — In pratis et in graminosis prope Haskovo, Jambok, Trnovo-Sejmen, Vitos.

Th. comptus FRIV. — In graminosis aridis prope Plovdiv, Nevrekop, m. Rhodope.

Th. acicularis W K. — In graminosis et in collinis aridis ad pagum Trojaci Macedoniae (det. RONN.).

Th. Boissieri HAL. — In herbis, saxosis regionis superioris m. Pirin. — *var. ciliatopubescens* HAL. — In rupestribus regionis alpinae m. Trojanski Balkan.

Th. pulvinatus ČEL. — In pascuis m. Stara planina. — *var. pirinicus* VEL. — In graminosis lapidosis m. Pirin.

Th. Neičeffii DEG. et URUM. — In graminosis lapidosis m. Pirin et Parnar-Dagh.

Th. carnosulus VEL. — In arenosis et in saxosis prope opp. Varna, Razgrad, Provadia, Šumen, Gebedže.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Ungarische Botanische Blätter](#)

Jahr/Year: 1926

Band/Volume: [25](#)

Autor(en)/Author(s): Urumoff I.K.

Artikel/Article: [Thymi bulgarici 227-229](#)