

KEYWORD-REGISTER

A

abundance	268, 310	cambium	76
activity	296	canopy structure	768
adaptation	70	<i>Capsella bursa-pastoris</i>	82
aerated sewage ponds	652	carabid beetles	98
agricultural	132	<i>Carabidae</i>	126
- areas	632	cereals	98
- fields	98	changes	
- landscape	424, 600, 718	- in agricultural landuse	434
- landuse	132	- in the weed flora	364
- soil	282, 310	climatic factors	82
agriculture	88, 554, 566	clonal growth	54
agro-ecosystems	514	<i>Collembola</i>	262, 268, 302, 310
<i>Alauda arvensis</i>	216	colonization	108
angiosperms	82	- strategies	404
<i>Anguilla anguilla</i>	694	community	
annual population cycle	88	- bird breeding	216, 632
annual rate of increase	88	- herbaceous edge	336
anthropogenic habitat conditions	616	- structure	644
<i>Apoidea</i>	638	- tall herb	768
aquatic biotops	670	- trampling	616
arable land	216, 268	competition	498
<i>Arrhenatherion</i>	442	consequences of intensification	424
artificial snow	746, 758	conservation	
Athabasca Glacier (Canada)	330	- measurements	44
Austria	116, 126	- measures	222
available soil water	386	- nature	566
		- site assessment	116
		- soil	718, 736

B

banks and dikes of small brooks	410	consolidation of farmland	210
biochemical nitrogen cycle	536	consumption	192
biological evaluation	126	contamination of the drinking water	536
biomass	296, 302, 478	conventional farming	498
- increment	54	crop rotation	514
- microbial	330	<i>Cruciferae</i>	82
biomonitoring system	670	cultivated	
biotope		- land	528
- losses	434	- plants	498
- quality	164	<i>Curculionidae</i>	116
bird		cutting regimes	410

- breeding community	216, 632	cycle	
- meadow	132	- annual population	88
<i>Brassicaceae</i>	70	- data	290
breeding bird community	216, 632	<i>Cygnus olor</i>	226

Bremen	108, 144	D	
brown hare	88, 566	data-bases	236
bumblebee	144	decomposition	276
Bundesministerium für Umwelt, Natur- schutz und Reaktorsicherheit	710	demographic analyses	44
Burgenland	116, 126	denitrification	652

C

calcareous fen	356	- dependent sources of regulation	184
<i>Calthion palustris</i>	488	- dependent fertility	226

different soils	268	fertility	170
Diptera larvae	282	fertilization	498, 506
distinct taste bud damages	682	fertilizers	178
distribution	200, 592, 662	field	254
- of species within the population	514	- agricultural	98
- patterns	374, 616	- clearance	316
- strategies	20	- margins	460
diversity	316	- ridges	450
- difference	316	field-track vegetation	600
dominance	336	fish	
dominant species	616	- ponds	698
dry grassland	116	- production	698
Dümmer	554	flavonoids	82
dwarf shrubs	768	flower-visiting insects	144
 E		forest	
ecological		- decline	76
- and geographical differentiation	616	- ecosystem	178
ecosystem		- edge	336
- forest	178	fragmentation	200
- grassland	768	freezing	60
eco-physiology	324	frequency	164, 336
ecological farming	498	 G	
edge communities	336	<i>Galium aparine</i>	60
eel	694	geographical	
efficiency	282	- and ecological differentiation	616
<i>Enchytraeidae</i>	192	- distribution	592
<i>Enchytraeus</i>	290	Geomorphologische Karte (GMK)	736
endangering	662	germination	30, 506
Endosulfan	302	- behaviour	70
environmental		GMK (Geomorphologische Karte)	736
- impact statement	126	grasshoppers	108
- monitoring	236	grassland	254, 442
- pollution	544	- dry	116
<i>Epilobium fleischeri</i>	30	- ecosystems	768
<i>Equisetum hyemale</i>	44	- fallows	470
erosion	398, 726, 746	- open	108
establishment	30	- moisture indicators	386
euvoecious	164	- wet	470
eutrophication	450, 566	grazing	152
evaluation	710	green areas	418
- biological	126	ground water	
extensive agricultural landuse	132	- regime	470
 F		- table	398
<i>Fagus sylvatica</i>	76	growth	
fallow land	528	- clonal	54
farming	498	- form	30, 44, 54, 450
- conventional	498	grue bugs	108
- ecological	498	 H	
- methods	424	habitat selection	222
feeding	222	<i>Halimione portulacoides</i>	54
<i>Felis silvestris</i>	200	heavy metal	310, 566
fen hay meadows	346	- pollution	624
fen-meadows	434	hedge	116
fens	478	herbaceous edge communities	336

hibernation sites	170	
home range utilisation	206	
hop	302	
husbandry	544	
 I		
ichthyofauna	662, 670	
impact statement	126	
impoverishment		
- in the nitrogen supply	606	
inclination	768	
incubation		
- in the field	528	
indicator values	386	
industrialised agriculture	554	
influence of climatic factors	82	
insecticide	170, 302	
insects	108	
- flower-visiting	144	
integration science	10	
intensification		
- consequences	424	
intensive livestock husbandry	544	
intraspecific variation	82	
inventarization	710	
 K		
key factors	184	
Kinzig	670	
 L		
Lahn	694	
land		
- cultivated	528	
land use		
- agricultural	132, 434	
- changes	20	
- conflicts	554	
- extensive agricultural	132	
landscape		
- agricultural	424, 600, 718	
- design	398	
- ecology	434	
- preservation	442	
landuse-changes	20	
- agricultural	434	
leaching	528	
leaf		
- inclination	768	
- litter	192	
- miner	184	
leafhoppers	108	
Lepidoptera	638	
<i>Lepus europaeus</i>	88, 566	
life		
- cycle data	290	
- forms	450	
- history strategy	226	
litter		
- consumption	192	
- meadow	356	
- bags	276	
lixiviation	536	
loach	682	
long distance spread	524	
long-term studies	418	
<i>Lumbricidae</i>	178, 644	
lumbricids	296	
 M		
macrophytic plants	710	
maintenance coefficient	324	
manure	528	
mapping	670	
marl	374	
maximal carbon uptake rate	324	
meadow		
- birds	132	
- fen	434	
- fen hay	346	
- litter	356	
- subalpine	758	
mechanical loads	262	
meltwater runoff	746	
<i>Mesenchytraeus glandulosus</i>	192	
metabolic quotient	324, 330	
metropolitan area	20	
microbial		
- biomass	330	
- respiration-to-biomass	330	
mineralisation		
- nitrogen	488, 528	
- rate of nitrogen	606	
modification of flavonoid		
accumulation	82	
moisture indicators on grassland	386	
<i>Molinia caerulea</i>	356	
<i>Molinio-Arrhenatheretea</i>	442	
monitoring		
- system	682	
mowing	356	
museums	236	
mute swan	226	
mycorrhizal infection	514	
 N		
N-fertilization	514	
natural		
- diversity	398	
- populations	70	

nature			
- conservation	566	- of natural soil erosion	736
- conservation in urban greens	418	predators	170
- reserve	592	predatory mites	262
neophytes	20	preference for hedges or trees	632
nitrate lixiviation	536	preservation of landscape	442
nitrification	652	protection	
nitrogen		- plant	210
- delivery	528	- soil	398
- input	606	- white stork	222
- mineralisation	488	protein data	290
- mineralisation	528	<i>Psammotettix putoni</i>	152
- mineralisation rate	606		
- removal	652		
- residue	528		
- supply	606		
<i>Noemacheilus barbatulus</i>	682	R	
North Limburg	544	radio-tracking	206
nutrient	346	realized annual rate of increase	88
- enrichment	478	recovery of aquatic biotops	670
- output	478	reproductive effort	54
O		reproductivity	670
occurrence	210	<i>Rhinanthus serotinus</i>	144
Odonata	164	<i>Rhynchaenus populi</i>	184
open grassland	108	river marsh	144
ordination	346	riverine marshland	132
P		road shoulder	616
PAR distribution	768	roadside	606
<i>Passer montanus</i>	210	- management	624
patterns	374, 616	- plant communities	624
perimeter of glacier	30	- soils	624
phenology	600	- structures	632
photosynthesis	60	- vegetation	566, 592, 600, 606
phytosociology	460	- verges	566, 638
plant		<i>Robinia pseudacacia</i>	116
- available nutrients	374	root	
- communities	364, 592	- bud	30
- cultivated	498	- growth pattern	398
- distribution	478	rotating disc	652
- macrophytic	710	Rotmoos Ferner (Austria)	330
- protection	210	runoff	726
- viruses	524		
<i>Platynus dorsalis</i>	170	S	
<i>Poecilus</i>	98	Saarland (West-Germany)	200
politics	718	salt	566
- science	236	<i>Saltatoria</i>	116
pollution	544, 624	saltmarsh	152
population	200	saprophytic bacteria	652
- analysis	88	<i>Schoenus ferrugineus</i>	356
- annual cycle	88	science politics	236
- biology	10	<i>Sciurus vulgaris</i>	206
- dynamics	184	seed	
- regulation	226	- bank	404, 410
- structure	30, 152	- mixture	398
- types	226	- rain	404
		sewage sludge	310
		significance of removal	170
		ski	
		- run	726
		- tourism	726

skiing-area	758	
skin papillomas	694	
soil	268, 524	
- agricultural	282, 310	
- available water	386	
- biomass	254	
- characteristics	498	
- chemistry	178	
- columns	296	
- compaction	262, 296	
- conservation	718, 736	
- cultivation	246, 276	
- erosion	736	
- fauna	246, 276	
- microbial biomass	324	
- oxygen concentration	746	
- protection	398	
- respiration	254	
- root zone	330	
- type	246	
- washing and flotation method	282	
<i>Solidago canadensis</i>	20	
South Oldenburg	544	
species	616	
- decline	434	
- distribution	670	
- diversity	268	
- richness	460	
specific leaf	192	
spider-communities	316	
spiders	126	
spore density	514	
standardized sample method	336	
<i>Stellario-Carpinetum</i>	374	
straw decomposition	276	
subalpine meadows	758	
sublethal water pollution	682	
succession	316	
sugar beets	170	
surface waters	524	
Switzerland	718, 758	
synsystematic classification	592	
<i>Syrphidae</i>	638	
		U
ultrastructural features		290
Universal Soil Loss Equation (USLE)		736
urban		
- environment		98
- greens		108, 418
<i>Urtica dioica</i>		336
USLE (Universal Soil Loss Equation)		736
		V
VA-mycorrhizae		514
Vechte-lowland		662
vegetation		
- change		478, 488
- development		424
- history		424
		W
waste water utilization		698
water		
- courses		710
- drinking...contamination		536
- ground...regime		470
- ground...table		398
- movement		296
- pollution		682, 694
- surface		524
weed		498
- communities		460
- species		506
- flora		364
West Lower Saxony		662
Westallgäu		478
wet grassland		470
white stork		222
windbreak		116
winter annuals		60
		X
xylem		76

T

tall herb community	768
taste bud damage	682
taxonomy	236, 290
terrestrial vegetation	450
thorn savannah	20
torrent events	726
trampling communities	616
tree sparrow	210
<i>Tribulus terrestris</i>	20
trickling filter	652
<i>Trifolio-Agrimonietum</i>	336

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Verhandlungen der Gesellschaft für Ökologie](#)

Jahr/Year: 1990

Band/Volume: [19_2_1990](#)

Autor(en)/Author(s): unbekannt

Artikel/Article: [Keyword-Register 781-785](#)