

# CO<sub>2</sub>-fixation in poplar leaves treated with ozone

Thomas Drogies\* & Robert Guderian

## Synopsis

Five weeks old cuttings of *Populus nigra* L. cv. Loenen were continuously exposed to 40 nL\*L<sup>-1</sup> (10.00–17.00 h) and 12 nL\*L<sup>-1</sup> ozone (17.00–10.00 h) for three weeks in a growth cabinet. After exposure the enzymes of CO<sub>2</sub>-fixation i.e. the photosynthetic ribulose-1,5-bisphosphate carboxylase-/oxygenase (Rubisco) and the anapleurotic phosphoenolpyruvate carboxylase (PEPcase) were analysed. Older leaves, fully expanded prior exposure, exhibited a decrease in total activity (by 50 %) and quantity (by 70 %) of Rubisco; PEPcase activity was doubled. In younger leaves, which were developed during ozone treatment, activity of both enzymes was still unchanged. Rubisco content, by contrast, was decreased by 30 %. The observation made in older leaves indicates that a higher percentage of the newly fixed carbon is used to support catabolic metabolism in the ozone stressed poplar.

CO<sub>2</sub>-fixation, ozone, PEPcase, *Populus nigra* L., Rubisco.

## 1. Introduction

In previous open-top-chamber-experiments the ozone-sensitive *Populus nigra* L. cv. Loenen exhibited a decrease in net photosynthesis (BALLACH & al. 1992) and alterations of carbohydrate levels (BÜCKER & BALLACH 1992) under the influence of ozone, especially in older leaves. These observations suggest an impact of ozone on the CO<sub>2</sub>-fixation, and thus, plant productivity. Key enzymes in this process are Rubisco and /or PEPcase. Rubisco catalyses the first irreversible step of CO<sub>2</sub>-fixation during photosynthesis leading to production of carbohydrates. In C<sub>3</sub>-plants, PEPcase is involved in anapleurotic reactions such as replenishment oxalacetate for tricarboxylic acid cycle (LATZKO & KELLY 1983). In this study the impact of ozone on both enzymes is investigated.

## 2. Material and methods

### 2.1 Plant material and exposure

Cloned cuttings of *Populus nigra* L. cv. Loenen purchased from Algemene Keuringsdienst voor Boomkwekerijgewassen (NAKB), The Hague/NL, were planted in 1 L pots filled with an unfertilized peat/sand mixture (17:3, v/v). Cultivation and following fumigation was performed in a phytotron (Härens). The day/night-regime was 16h/8h with a light intensity of 420 μmol\*m<sup>-2</sup>\*s<sup>-1</sup>. Relative humidity was between 75 % and 45 %, temperature by 23°C and 16°C.

After five weeks of cultivation plants were exposed for three weeks. The ozone concentration was by 40 (±10) nL\*L<sup>-1</sup> from 10.00–17.00 h and 12 (±5) nL\*L<sup>-1</sup> from 17.00–10.00 h. The controls were supplied with charcoal (Purafil) filtered air.

Fertilizing was started three weeks after implanting of the cuttings by adding 100 mg of micronutrient (Hortrilon) to each plant, followed weekly by 0.5 g of N-P-K-fertilizer (15-10-15; Hakaphos blau).

For analyses samples were taken from leaves fully expanded prior fumigation, termed older leaves, and leaves developed during exposure, termed younger leaves.

### 2.2 Analyses

Extraction of enzymes was performed according to DANN & PELL (1989). Total activity of Rubisco was measured spectrophotometrically as described by LILLEY & WALKER (1974), the activity of PEPcase according to ASHTON (1990). The amount of Rubisco was determined by gel electrophoresis as described by KEYS & PARRY (1990) and chlorophyll content was measured according to ARNON (1949).

## 3. Results

After three weeks of exposure to low ozone concentrations, the older leaves exhibited a decrease in Rubisco activity and quantity by 50 % and 70 %, respectively (fig. 1). The activity of PEPcase was doubled (fig. 1).

The younger leaves did not achieve the Rubisco content observed in leaves of plants kept in filtered air.

\* Corresponding author.




Fig. 1  
Impact of ozone on total activity and quantity of Rubisco, on PEPcase activity and chlorophyll content in leaves of cv. Loenen.   
□ control ▨ ozone.   
Values are  $\bar{x}$  (n=6); SD.

Rubisco- and PEPcase activity remained unchanged. All these alterations occurred before injuries were visible, only a slight decrease in chlorophyll content was detected (fig. 1).

#### 4. Discussion

It can be concluded, that the alterations in photosynthesis and carbohydrate levels are also caused by the reduction in quantity and activity of Rubisco. The finding in younger leaves suggests a diminished synthesis of Rubisco protein under the influence of ozone.

In protease assays with artificial substrate no increased activity was found (data not shown). Therefore an increased degradation of Rubisco is not very probable. Alterations in status of Rubisco after exposure to ozone has been found before in wheat (LEHNHERR & al. 1987), potato (DANN & PELL 1989), radish (PELL & al. 1992) and the hybrid *Populus maximowizii x trichocarpa* (PELL & al. 1992).

Beyond this, increase of PEPcase-activity suggest that newly fixed  $\text{CO}_2$  is used for feeding catabolic metabolism. Increase of PEPcase has been reported for spruce (TIETZ & WILD 1991).

Decrease in Rubisco quantity and activity is also assumed to contribute to normal leaf senescence (WITTENBACH & al. 1980) and premature senescence induced by air pollutants (LANDRY & PELL 1993). Premature foliar senescence due to air pollutants has also been described for cv. Loenen (BALLACH & al. 1988).

### Acknowledgements

This research was supported by a grant from Gesamtverband des deutschen Steinkohlenbergbaus, Essen. The skilful technical assistance of G. Friesewinkel and C. Kosch is gratefully acknowledged.

### References

- ARNON, D. J., 1949: Copper enzymes in isolated chloroplasts. Polyphenoloxidase in *Beta vulgaris*. – *Plant Phys.* 24: 1–15.
- ASHTON, A.R., BURNELL, J.N., FURBANK, R.T., JENKINS, J.L.D. & M.D. HATCH, 1990: Enzymes of C4 Photosynthesis. – In: P. J. LEA (Vol. ed.): *Enzymes of primary metabolism*: 40–73. – In: P.M. DEY & J.B. HARBORNE (Ser. ed.): *Methods in Plant Biochemistry*, vol. 3. – Academic Press, London/Toronto.
- BALLACH, H.J., MOOI, J. & R. WITTIG, 1992: Premature aging in *Populus nigra* L. after exposure to air pollutants. – *Angew. Botanik* 66: 14–20.
- BALLACH, H.J., MOOI, J. & J. BÜCKER, 1988: Injury of populus nigra Loenen cultivars caused by flue gas from an atmospheric fluidised bed boiler (AFBB). – *Acta Biol. Benrodis* 1: 69–79.
- BÜCKER, J. & H.J. BALLACH, 1992: Alterations in carbohydrate levels in leaves of *Populus* due to ambient air pollution. – *Phys. Plant.* 86: 512–517.
- DANN, M.S. & E.J. PELL, 1989: Decline of activity and quantity of ribulose bisphosphate carboxylase/oxygenase and net photosynthesis in ozone-treated potato foliage. – *Plant Phys.* 91: 427–432.
- KEYS, A.J. & M.A.J. PARRY, 1990: Rubisco and carbonic anhydrase. In: P.J. LEA (Vol. ed.): *Enzymes of primary metabolism*: 1–13. – In: P.M. DEY & J.B. HARBORNE (Ser. ed.): *Methods in Plant Biochemistry*, vol. 3. – Academic Press, London/Toronto.
- LANDRY, L.G. & E.J. PELL, 1993: Modification of rubisco and altered proteolytic activity in O<sub>3</sub>-stressed hybrid Poplar (*Populus maximowizii* x *trichocarpa*). – *Plant Phys.* 101: 1355–1362.
- LATZKO, E. & G.J. KELLY, 1983: The many-faceted function of phosphoenolpyruvate carboxylase in C3 plants. – *Phys. Veg.*, 21: 805–815.
- LEHNHERR, B., GRANDJEAN, A., MÄCHLER, F. & J. FUHRER, 1987: The effect of ozone in ambient air on ribulosebisphosphate carboxylase/oxygenase activity decreases photosynthesis and grain yield in wheat. – *J. Plant Phys.*, 130: 189–200.
- LILLEY, R.McC. & D.A. WALKER, 1974: An improved spectrophotometric assay for ribulosebisphosphate carboxylase. – *Biochim. Biophys. Acta* 358: 226–229.
- PELL, E.J., ECKHARDT, N.A. & A.J. ENYEDI, 1992: Timing of ozone stress and resulting status of ribulose bisphosphate carboxylase/oxygenase and associated net photosynthesis. – *New Phytol.* 120: 397–405.
- WITTENBACH, V.A., ACKERSON, R.C., GIAQUINTA, R.T. & R.R. HERBERT, 1980: Changes in photosynthesis, ribulose bisphosphate carboxylase, proteolytic activity and ultrastructure of soybean leaves during senescence. – *Crop Sci.* 20: 225–231.

### Adresse

R. Guderian  
Th. Drogies  
Institut für angewandte Botanik, FB 9.4  
UNI/GHS Essen  
Universitätsstraße 5  
45117 Essen  
Fax: (0201) 183- 2529

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Verhandlungen der Gesellschaft für Ökologie](#)

Jahr/Year: 1995

Band/Volume: [24\\_1995](#)

Autor(en)/Author(s): Guderian Robert, Drogies Thomas

Artikel/Article: [C02-fixation in poplar leaves treated with ozone 271-273](#)