

Some hoverfly records from Turkey (Diptera, Syrphidae)

Menno Reemer and John T. Smit

Reemer, M.; Smit, J.T. (2007): Some hoverfly records from Turkey (Diptera, Syrphidae). – Volucella 8, 135-146. Stuttgart.

In this paper 84 species of hoverflies are recorded from Turkey, based on material collected in several parts of the country between 1991 and 2003. As far as the authors could establish, 13 of these species have not been previously recorded for Turkey. These are: *Anasimyia transfuga*, *Cheilosia portschinskiana*, *C. schnabli*, *Chrysogaster mediterraneus*, *C. simplex*, *Eupeodes nuba*, *Merodon aeneus*, *Mesembrius peregrinus*, *Neoascia annexa*, *N. interrupta*, *Pipiza festiva*, *Sphegina sibirica* and *Sphiximorpha subsessilis*. Brief comments are made on some of the species.

Key words: Turkey, new records, Syrphidae

Zusammenfassung

Nachweise von 84 Schwebfliegenarten aus der Türkei werden vorgelegt, gefangen in verschiedenen Landesteilen zwischen 1991 und 2003. 13 der nachgewiesenen Arten waren vorher aus der Türkei nicht bekannt: *Anasimyia transfuga*, *Cheilosia portschinskiana*, *C. schnabli*, *Chrysogaster mediterraneus*, *C. simplex*, *Eupeodes nuba*, *Merodon aeneus*, *Mesembrius peregrinus*, *Neoascia annexa*, *N. interrupta*, *Pipiza festiva*, *Sphegina sibirica* und *Sphiximorpha subsessilis*.

Introduction

At the crossroads between Africa, Asia and Europe, Turkey hosts a very interesting fauna with mixed elements derived from each of these three continents. Despite this, and despite the recent publication of several papers by Turkish authors, the syrphid fauna of Turkey is still relatively poorly known. This paper presents some records of Syrphidae collected in Turkey during the past couple of years. Although the records originate from various parts of the country, they will by no means provide a complete image of the fauna. This is merely a compilation of records collected by the authors and a few other Dutch entomologists, during a couple of short visits to Turkey. With this paper we hope to make a useful, albeit modest, contribution to the knowledge of the Syrphidae of this large, diverse and interesting country.

Material and methods

Most of the species recorded were collected in 2000 by the authors in the province of Muğla in southwestern Turkey. Additional specimens were been collected in other parts of Turkey by Kees Goudsmits, Vincent Kalkman, Herman Teunissen and Theo Zeegers in the years 1991, 2001, 2002 and 2003.

Most identifications were carried out by the authors, except for the *Merodon* specimens, which were identified by Willem Hurkmans (Zwolle, Netherlands), and specimens of *Eupeodes*, which were identified by Libor Mazánek (Olomouc, Czech Republic). All collected specimens are deposited in the collections of the authors, except for a small number of specimens in the collection of Kees Goudsmits (Doorn, the Netherlands).

To check whether a species had already been recorded for Turkey, the following literature has been examined: Aktas & Saribiyik (1996, 2001), Candemir & Kara (2003), Claussen & Hayat (1997a, b), Coe (1966), Dirickx (1994), Hayat (1997), Hayat & Alaoğlu (1990a, b), Hayat & Claussen (1997), Hurkmans (1987, 1988, 1993), Hurkmans & Hayat (1997), Lindner (1941), Özgür (1986, 1987), Özgür & Saribiyik (2002), Reemer et al. (2005), Saribiyik (1999a, b, 2000a, b, 2001, 2003a, b, c, d, e, 2004), Saribiyik & Aktas (1996), Saribiyik & Hasbenli (1997), Saribiyik & Özgür (2000), Speight & Castella (2003), van Steenis (2000), Toth (1986), Van de Weyer (2000).

Localities

The localities visited are shown in fig. 1, grouped by province. The first two letters of the name of the province are used for the locality codes. For each locality, coordinates (latitude/longitude or UTM grid reference), height above sea level (in metres) and the date(s) of the visit(s) are given. If available, a description of the habitat is given.

Afyon: AF01 24 km SW of Çay, Karamk Gölü and irrigation channels 1 km E of Bulanik köy. 38°30'N 30°55'E. 1000 m asl. 4.vii.2001 (leg. V.J. Kalkman). **AF02** Emir Dagları, 15 km SSE of Emirdag lake at 10 km S of Dereköy. 39°00'N 31°10'E. 1500-1700 m asl. 5.vii.2001 (leg. V.J. Kalkman).

Antalya: AN01 11 km NW of Kalkan, Eşen Çayı near Kinikli. UTM: 35SQA02. 10 m asl. 27.v.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). **AN02** 4 km W of Kalkan, ancient harbour of Patara S of Gelemiş. UTM: 35SQA01. 5 m asl. 27.v.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). **AN03** Lake at Civizli, 20 km N of Akseki. 37°11'N 31°45'E. 11.vii.2001 (leg. V.J. Kalkman).

Artvin: AR01 Coruh N, in the village of Yusufeli. UTM: 37TGF12. 3.vii.2003 (leg. V.J. Kalkman).

Gaziantep: GA01 Durnalik, 20 km W of Gaziantep. 13.v.2002 (leg. K. Goudsmits).

İçel: IC01 Western side of Akgöl near Denizkent, 8 km S of Silifke, Gökusu Delta. 36°17'N 33°56'E. 9.vii.2001 (leg. V.J. Kalkman). Dunes. **IC02** Abanoz, Anamur, Kazancı 1190 m asl. 36°30'N 32°53'E. 10.v.1991 (leg. H. Teunissen).

İsparta: IS01 Kovada Milli Park. 37°40'N 30°50'E. 1.vii.2001 (leg. V.J. Kalkman). Freshwater lake.

Kayseri: KA01 Yahyalı, Ovaçiftliği, Sultanswamp. 1075 m asl. 38°15'N 35°17'E. 7.v.2002 (leg. T. Zeegers).

Muğla: MU01 Köyceğiz, just E of village, near Köyceğiz Gölü. UTM: 35SPA59. 5 m asl. 21.v.2000, 22.v.2000, 24.v.2000, 29.v.2000, 30.v.2000, 3.vi.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). Various habitats: forest of *Liquidamber orientalis* with small streams, marshy grasslands and fields with large, flower-




Fig. 1: Map of Turkey (excluding the islands), with the provinces indicated from which Syrphidae are recorded in this paper. AF: Afyon, AN: Antalya, AR: Artvin, GA: Gaziantep, IC: İçel, IS: Isparta, KA: Kayseri, MU: Muğla, NE: Nevşehir, NI: Niğde, TR: Trabzon, VA: Van.

ing *Euphorbia* spec. **MU02** 1-2 km W of Köyceğiz. UTM: 35SPA49. 5 m asl. 22.v.2000, 30.v.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). Marshy grassland and edges of forest of *Liquidamber orientalis*. **MU03** 3 km NE of Ortaça, along Dalaman Çayı. UTM: 35SPA58. 50 m asl. 23.v.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). **MU04** Muğla 8 km W of Köyceğiz (Muğla), dry rocky slopes 2 km NE of Doğuşbelen. UTM: 35SPA49. 50 m asl. 24.v.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). Dry rocky slopes near pool. **MU05** 4-6 km ENE of Ortaça, river valley of tributary of Dalaman Çayı. UTM: 35SPA67. 50 m asl. 25.v.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). **MU06** Kaunos, near ancient amphitheatre. UTM: 35SPA47. 1-20 m asl. 26.v.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). Dry, rocky hills. **MU07** 10 km NNE of Kestep (=Eşen), Eşen Çayı near bridge E of Alaçatı. UTM: 35SQA14. 25 m asl. 27.v.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). **MU08** Saklıkent, 13 km ENE of Kestep (=Eşen), 25 km ESE of Fethiye. UTM: 35SQA13. 50 m asl. 27.v.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). Trout farm. **MU09** 12 km NNE of Kemer, just N of Ören. UTM: 35SQA17. 100 m asl. 28.v.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). Deciduous forest around trout farm. **MU10** 16 km NNW of Üzümlü, between Arpacık and Karabayır, near Kırkpınar. UTM: 35SPA98. 1350 m asl. 28.v.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). Mountain slopes with streams and marshy area. **MU11** Beyobaşı, E of Köyceğiz Gölü, around trout farm. UTM: 35SPA58. 50 m asl. 28.v.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). **MU12** 6 km W of Köyceğiz, 2 km W of Toparlar. UTM: 35SPA49. 50-100 m asl. 29.v.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). Arboretum with waterfalls. **MU13** Between Sultaniye and Küçükkaraağaç. UTM: 35SPA48. 50 m asl. 30.v.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). **MU14** 5 km W of Sultaniye, near Köyceğiz Gölü. UTM: 35SPA48. 50 m asl. 30.v.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). **MU15** Kartal Gölü, 12 km NE of Ağla. UTM: 35SPB60. 1900 m asl. 31.v.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). Mountain lake with small stream, surrounded by short, grassy vegetation. **MU16** 13 km NNE of Ağla. UTM: 35SPB60. 1600 m asl. 31.v.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). Coniferous forest with some yellow flowers along a stream. **MU17** Gökçeova Gölü, 6 km NE of Ağla. UTM: 35SPB50. 1400 m asl. 31.v.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). Lake with some small streams. **MU18** Just E of Ağla. UTM: 35SPB50. 850 m asl. 1.vi.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). Deciduous forest (mainly *Platanus*) with springs and small streams. **MU19** 7 km SW of Köyceğiz, near bridge SW of Hamitköy, W of Köyceğiz Gölü. 5 m. asl. UTM: 35SPA48. 1.vi.2000 (leg. M. Reemer, J.T. Smit & K. Goudsmits). **MU20** 10 km E of Köyceğiz, spring of rivulet Yavurlakçayı above Beyobaşı, E of Taşdibi, 100

m asl. UTM 35SPA69. 1.vi.2000 (leg. J.T. Smit). **MU21** 5 km S of Köyceğiz, mouth of Yavurlakçayı into Köyceğiz lake, 5 m asl. UTM 35SPA58. 2.vi.2000 (leg. J.T. Smit).

Nevşehir: NE01 Ürgüp, Göreme, 1150 m asl. 38°38'N 34°56'E. 11.v.2002 (leg. T. Zeegers). **NE02** Ürgüp, hilltop 3 km W of Ortahisar, 1450 m asl. 38°38'N 34°56'E. 12.v.2002 (leg. T. Zeegers). **NE03** Ürgüp, hilltop at Ortahisar, 1300 m asl. 38°38'N 34°56'E. 12.v.2002 (leg. T. Zeegers).

Niğde: NI01 Ihlara-Valley, 1250 m. asl. 37°59'N 34°42'E. 15.v.2002 (leg. T. Zeegers). **NI02** Sultanhanı, 1050 m asl. 38°15'N 33°33'E. 17-19.v.2002 (let. T. Zeegers).

Trabzon: TR01 2 km E of Yomra, along the Harmanlı Çayı. UTM: 37TEF73. 24.vi.2003 (leg. V.J. Kalkman). **TR02** 10 km S of Arakli, along the Kara Dere. UTM: 37TEF82. 24.vi.2003 (leg. V.J. Kalkman). **TR03** Uzungöl. 40°37'N 26°50'E. 9.vii.2003 (leg. V.J. Kalkman).

Van: VA01 8 km N of Muradiye, Bendimahi Selalesi. UTM: 38SLJ92. 30.vi.2003 (leg. V.J. Kalkman).

Results

In this paper, 84 species are recorded for Turkey (table 1). For each species, the available literature has been checked for previous records from Turkey. So far as we can ascertain, the following 13 species have not been recorded for Turkey before: *Anasimya transfuga*, *Cheilosia portschinskiana*, *C. schnabli*, *Chrysogaster mediterraneus*, *C. simplex*, *Eupeodes nuba*, *Merodon aeneus*, *Mesembrius peregrinus*, *Neoscia annexa*, *N. interrupta*, *Pipiza festiva*, *Sphegina sibirica* and *Sphiximorpha subsessilis*.

Notes on some species

Cheilosia canicularis: These specimens have been checked for the closely related species *C. griseifacies* Vujić, 1994, *C. himantopus* (Panzer, 1798) and *C. orthotricha* Vujić & Claussen, 1994 using characters provided by Stuke & Claussen (2000), Vujić (1994) and Vujić & Claussen (1994).

Cheilosia portschinskiana: It is possible that some of the records of *C. illustrata* in Saribiyik (2001) belong to this species as well.

Cheilosia schnabli: This species is very similar to *C. impressa* Loew, 1840, but easily recognized by the characters described by Vujić et al. (1998), especially in direct comparison. *C. schnabli* has so far been recorded from the Balkan and the Caucasus. This new record of a male specimen is from a mountain range extending from the Caucasus along the southeastern Black Sea.

Chrysogaster mediterraneus: This appears to be the first record outside the Balkans, from where it has been described by Vujić (1999). It is possible that the specimens of the very similar *C. solstitialis* (Fallén, 1817) reported by Saribiyik (1999b, 2001) also belong to this species.

Merodon velox: At the end of a hot day, a male specimen was observed drinking at a small stream along a roadside.

Neoscia annexa: The specimens have been compared with the description of *N. subannexa* by Claussen & Hayat (1997a), but they did not belong to that species. They have also been compared with specimens of *N. annexa* from Northwest-European countries, to which they were found to be very similar.

Platycheirus scutatus: This specimen has been checked for the closely related *P. aurolateralis* Stubbs, 2002, *P. speighti* Doczkal et al., 2002 and *P. splendidus* Rotheray, 1998.

Riponnensia insignis: A large series of this little known species has been collected near Köyceğiz (prov. Muğla). The specimens were taken from *Euphorbia* flowers on a marshy grassland next to a forest of *Liquidamber orientalis* with small streams. *R. insignis* has previously been recorded for Turkey by Dirickx (1994) (without citing the original reference for this record) and by Özgür (1987). Males can be identified most reliably using the drawings of the genitalia provided by Maibach et al. (1994) in combination with Claussen (1991) and Vujić (1999). The female has not been described yet, but would run to the correct species in the keys of Sack (1932) and Stackelberg (1953).

Sphagina sibirica: Has been recorded from the adjacent Caucasus by Toth (1986).

Sphiximorpha subsessilis: The specimens were checked for other *Sphiximorpha* species, including *S. worelli* (Bradescu, 1972), which has recently been recorded for Turkey by Saribiyik (2003a). We have also compared them with specimens of *S. subsessilis* from other European countries and found that they are very similar in both external morphology and male genitalia.

Discussion

An annotated checklist would be a very useful instrument in the study of the Syrphidae of Turkey. The compilation of a reliable checklist would require a great effort, because of the taxonomic challenge lurking in the fauna of this country, probably still containing many undescribed species. Besides, it would be necessary to re-examine the records of several species which have been published in older literature, in order to establish their true identity. This is required in the case of species complexes containing recently described 'cryptic species', for instance: *Chrysogaster mediterraneus* / *C. solstitialis*, *Myolepta dubia* / *M. trojana*, *Paragus bicolor* / *P. romanicus* and *Platycheirus aurolateralis* / *P. scutatus* / *P. splendidus*.

Large collections of Turkish Syrphidae, containing a lot of unidentified material and undescribed species, are awaiting examination. An important example is the collection of J.A.W. Lucas, now included in the collection of the ZMAN in Amsterdam (the Netherlands). Such important collections should not be ignored when examining the Turkish fauna.

Acknowledgements

We are indebted to Kees Goudsmits, Vincent Kalkman, Herman Teunissen and Theo Zeegers for enabling us to include records from their collecting trips in Turkey. Willem Hurkmans and Libor Mazánek have been helpful in identifying some *Merodon* and *Eupeodes* specimens, for which we are very grateful. We would also like to thank Suleyman Saribiyik for kindly sending us reprints of several useful papers on the Turkish Syrphidae fauna.

References

- Aktas, M.; Saribiyik, S. (1996): Contribution to the Syrphidae fauna of Turkey (Diptera: Syrphidae) II. Milesiinae. – Journal of the Institute of Science and Technology, Gazi University 9, 15-27.
- Aktas, M.; Saribiyik, S. (2001): New records of Syrphinae (Diptera: Syrphidae) from Turkey. – Journal of the Entomological Research Society 31, 41-46.
- Candemir, D.; Kara, K. (2003): Syrphidae (Diptera) fauna in Tokat provinces (Turkey). – Türkiye Entomoloji Dergisi 27, 95-105.
- Claussen, C. (1991): Eine neue *Orthonevra* von Korsika (Diptera, Syrphidae). – Entomofauna 12(16), 205-211.
- Claussen, C.; Hayat, R. (1997a): A new species of the genus *Neoascia* Williston, 1886 (Diptera, Syrphidae) from North-East Turkey. – Studia Dipterologica 4, 215-218.
- Claussen, C.; Hayat, R. (1997b): A new species and new records of the genus *Pipizella* Rondani, 1856 (Diptera, Syrphidae) from Turkey. – Studia Dipterologica 4, 447-452.
- Coe, R.L. (1966): A new *Sericomyia* from Turkey (Diptera: Syrphidae). – Entomologisk Tidskrift 87, 118-120.
- Dirickx, H.G. (1994): Atlas des Diptères syrphides de la région méditerranéenne. – Studiedocumenten van het Koninklijk Belgisch Instituut voor Natuurwetenschappen 75, 1-317.
- Hayat, R.; Alaoğlu, Ö. (1990a): Erzurum yöresi Syrphidae (Diptera) faunası (I). Syrphinae. – Türkiye Entomoloji Dergisi 14, 173-182.
- Hayat, R.; Alaoğlu, Ö. (1990b): Erzurum yöresi Syrphidae (Diptera) faunası (II). Milesiinae. – Türkiye Entomoloji Dergisi 14, 227-234.
- Hayat, R.; Claussen, C. (1997): A new species and new records of the genus *Paragus* Latreille, 1804 from Turkey (Diptera: Syrphidae). – Zoology in the Middle East 14, 99-108.
- Hurkmans, W. (1987): *Merodon telmateia*, a new hoverfly from Turkey (Diptera: Syrphidae). – Entomologische Berichten 47, 119-121.
- Hurkmans, W. (1988): Ethology and ecology of *Merodon* in Turkey (Diptera: Syrphidae). – Entomologische Berichten 48, 107-114.
- Hurkmans, W. (1993): A monograph of *Merodon* (Diptera: Syrphidae). Part 1. – Tijdschrift voor Entomologie 136, 147-234.
- Hurkmans, W.; Hayat, R. (1997): Ethology and ecology of *Merodon* (Diptera, Syrphidae) in Turkey II: descriptions of new species and notes on other syrphid flies. – Dipterists Digest 3, 62-79.
- Lindner, E. (1941): Eine entomologische Studienfahrt nach Anatolien und ihre dipterologischen Ergebnisse. – Jahreshefte des Vereins für Vaterländische Naturkunde in Württemberg 96, 113-136.
- Maibach, A.; Goeldlin de Tiefenau, P.; Speight, M.C.D. (1994): Limites génériques et caractéristiques taxonomiques de plusieurs genres de la tribu des Chrysogasterini (Diptera: Syrphidae). I. Diagnoses génériques et description de *Riponnensia* gen. nov. – Annales de la Société Entomologique de France 30, 217-247.
- Milankov, V.; Vujić, A.; Ludoski, J. (2001): Genetic divergence among cryptic taxa of *Merodon avidus* (Rossi, 1790) (Diptera: Syrphidae). – International Journal of Dipterological Research 12, 15-24.
- Özgür, A.F. (1986): Akdeniz Bölgesi avcı Syrphidae Türleri. – Proceedings of the first Turkish national congress of biological control, Çukurova Üniversitesi Ziraat Fakültesi Bitki Koruma Bölümü, Adana, 293-303.
- Özgür, A.F. (1987): Die Fauna der Familie Syrphidae (Dipt.) von Mittelmeerküstengebiet der Türkei. II. Unterfamilien: Cheilosini, Sphegini, Pelecocerini, Milesiini, Chrysotoxini, Volucellini. – Ziraat Fakültesi Dergisi, Çukurova Üniversitesi, Adana 2, 67-81.
- Özgür, A.F.; Saribiyik, S. (2002): New records of Microdontinae (Diptera: Syrphidae) from Turkey. – Journal of the Entomological Research Society 4, 15-17.
- Reemer, M.; Hauser, M.; Speight, M.C.D. (2005): The genus *Myolepta* Newman in the West-Palaearctic region. – Studia dipterologica 11(2), 553-580.
- Sack, P. (1932): Syrphidae. – Die Fliegen der palaearktischen Region 4, 1-451.
- Saribiyik, S. (1999a): Syrphinae fauna of the West Blacksea Region (Diptera: Syrphidae). – Kastamonu Eğitim Dergisi 7, 185-194.
- Saribiyik, S. (1999b): Milesiinae fauna of the West Blacksea Region (Diptera: Syrphidae). – Kastamonu Eğitim Dergisi 7, 195-204.

- Sarıbiyik, S. (2000a): **Fauna of Syrphidae in Ilgaz and Isik mountains and their vicinity (Diptera-Syrphinae).** – Journal of the Institute of Science and Technology, Gazi University 13, 55-70.
- Sarıbiyik, S. (2000b): **Two new records for the Turkish Milesiinae fauna (Diptera: Syrphidae).** – Bitki Koruma Bülteni 40, 179-181.
- Sarıbiyik, S. (2001): **New records of the subfamily Milesiinae (Diptera: Syrphidae) from Turkey.** – Journal of the Entomological Research Society 3, 43-51.
- Sarıbiyik, S. (2003a): *Sphiximorpha worelli* (Bradescu, 1972), (Diptera: Syrphidae) New record for the fauna of Turkey. – Journal of the Entomological Research Society 5, 13-16.
- Sarıbiyik, S. (2003b): **Fauna of Syrphinae and Milesiinae (Diptera: Syrphidae) around Tuz Lake.** – Kastamonu Education Journal 11, 439-450.
- Sarıbiyik, S. (2003c): **The evaluation of the works on Syrphidae (Diptera) fauna in the western Blacksea region.** – Kastamonu Education Journal 11, 461-466.
- Sarıbiyik, S. (2003d): **Milesiinae fauna of Kahramanmaraş's Andirin town (Diptera: Syrphidae).** – Kastamonu Education Journal 11, 159-164.
- Sarıbiyik, S. (2003e): **Syrphinae fauna of Kahramanmaraş's Andirin town (Diptera: Syrphidae).** – Kastamonu Education Journal 11, 193-198.
- Sarıbiyik, S. (2004): **Three new records for Turkish Milesiinae fauna (Diptera: Syrphidae).** – Journal of the Entomological Research Society 6, 19-22.
- Sarıbiyik, S.; Aktas, M. (1996): **Contributions to the Syrphidae fauna of Turkey (Diptera: Syrphidae) I. Syrphinae.** – Journal of the Institute of Science and Technology, Gazi University 9, 1-13.
- Sarıbiyik, S.; Hasbenli, A. (1997): **Türkiye Syrphidae (Diptera) faunası için yeni kayıtlar.** – Türkiye Entomoloji Dergisi 21, 225-232.
- Sarıbiyik, S.; Özgür, A.F. (2000): **New records of Milesiinae (Diptera: Syrphidae) from Turkey.** – Journal of the Entomological Research Society 2, 5-13.
- Speight, M.C.D.; Castella, E. (2003): **Range and status data for European Syrphidae (Diptera), 2003.** – In: Speight, M.C.D.; Castella, E.; Sarthou, J.P.; Ball, S. (eds.), *Syrph the Net, the database of European Syrphidae* 42, 1-370. Dublin (Syrph the Net publications).
- Stackelberg, A.A. (1953): **Paläarktische vidy roda *Orthoneura* Macq. (Diptera, Syrphidae).** – Entomologiceskoe Obozrenie 33, 342-357.
- Steenis, J. van (2000): **The West-Palaearctic species of *Spilomyia* Meigen (Diptera, Syrphidae).** – Mitteilungen der Schweizerischen Entomologischen Gesellschaft 73, 143-168.
- Stuke, J.-H.; Claussen, C. (2000): ***Cheilosia canicularis* auctt. - ein Artenkomplex.** – Volucella 5, 79-94.
- Toth, S. (1986): **Beiträge zur Kenntnis der Schwebfliegen-Fauna des Südwest-Kaukasus (Diptera: Syrphidae).** – Folia Musei Historico-Naturalis Bakonyiensis 5, 85-98.
- Van de Weyer, G. (2000): **A new syrphid fly species of the *Paragus serratus* complex from Turkey (Diptera: Syrphidae).** – Phegea 28, 149-152.
- Vujić, A. (1994): ***Cheilosia griseifacies*, eine neue Fliegen-Art aus Mitteleuropa (Diptera, Syrphidae).** – Entomofauna 15, 337-344.
- Vujić, A. (1999): **The tribe Chrysogasterini (Diptera: Syrphidae) in the Balkan Peninsula, with the description of three new cryptic species.** – Studia Dipterologica 6, 405-423.
- Vujić, A.; Claussen, C. (1994): ***Cheilosia orthotricha* spec. nov., eine weitere Art aus der Verwandtschaft von *Cheilosia canicularis* aus Mitteleuropa.** – Spixiana 17, 261-267.
- Vujić, A.; Radenković, S.; Stănescu, C. (1998): **New data on hoverflies (Diptera, Syrphidae) in Romania.** – Volucella 3, 63-74.

Addresses of authors:

Menno Reemer & John T. Smit, EIS-Nederland, Postbus 9517, 2300 RA Leiden, The Netherlands. E-mail: reemer@naturalis.nl, smitj@naturalis.nl

Table 1: Hoverfly records from Turkey. The locality codes are explained in the text. In some cases the number and sex of the specimens is given between brackets (f = female, m = male). In the last column references are given to literature in which the species has previously been recorded for Turkey.

	species	localities	previous references for Turkey
1.	<i>Anasimyia transfuga</i> (Linnaeus, 1758)	IS01 (1 f)	
2.	<i>Brachypalpoides lentus</i> (Meigen, 1822)	MU18 (1 m)	Dirickx (1994)
3.	<i>Ceriana conopsoides</i> (Linnaeus, 1758)	MU01, MU02	Dirickx (1994), Saribiyik (2001, 2003b, c)
4.	<i>Ceriana vespiiformis</i> (Latreille, 1809)	MU01, MU02, MU03, MU05, MU14, MU18, MU19	Dirickx (1994)
5.	<i>Chalcosyrphus nemorum</i> (Fabricius, 1805)	MU01	Saribiyik & Özgür (2000)
6.	<i>Cheilosia aerea</i> Dufour, 1848	NI01 (1 m), MU10 (2 f)	Dirickx (1994; as <i>C. zetterstedti</i>) Lindner (1941; as <i>C. zetterstedti</i>)
7.	<i>Cheilosia canicularis</i> (Panzer, 1801)	AR01, TR03	Saribiyik (1999b, 2001)
8.	<i>Cheilosia portschinskiana</i> Stackelberg, 1960	TR03 (1 f)	
9.	<i>Cheilosia schnabli</i> Becker, 1894	TR03 (1 m)	
10.	<i>Cheilosia soror</i> (Zetterstedt, 1843)	MU10	Aktas & Saribiyik (1996; as <i>C. rufipes</i>), Özgür (1987), Saribiyik (1999b, 2001, 2003d; as <i>C. rufipes</i>), Saribiyik (2003b, c)
11.	<i>Chrysogaster mediterraneus</i> Vujić, 1999	MU01 (2 m), MU18 (1 m)	
12.	<i>Chrysogaster simplex</i> Loew, 1843	MU18 (2 m, 2 f)	
13.	<i>Chrysotoxum cautum</i> (Harris, 1776)	IC02 (1 m), MU10 (1 f), UR01 (1 m)	Candemir & Kara (2003), Dirickx (1994), Saribiyik (1999a, 2003b, c)
14.	<i>Chrysotoxum festivum</i> (Linnaeus, 1758)	VA01 (1 m)	Dirickx (1994), Hayat & Alaoğlu (1990a), Hurkmans & Hayat (1997), Saribiyik (1999a, 2000a, 2003b, c), Saribiyik & Aktas (1996)
15.	<i>Chrysotoxum intermedium</i> Meigen, 1822	MU10 (1 f)	Candemir & Kara (2003), Dirickx (1994), Özgür (1987), Saribiyik (1999a, 2000a, 2003b, c, e), Saribiyik & Aktas (1996)
16.	<i>Epistrophe eligans</i> (Harris, 1780)	IC02 (2 m)	Dirickx (1994), Saribiyik (1999a, 2000c), Özgür (1986), Saribiyik & Aktas (1996)

17.	<i>Episyrphus balteatus</i> (De Geer, 1776)	MU01, MU02, MU18	Candemir & Kara (2003), Dirickx (1994), Özgür (1986), Saribiyik (1999a, 2000a, 2003b, c, e), Saribiyik & Aktas (1996)
18.	<i>Eristalinus aeneus</i> (Scopoli, 1763)	MU01, MU03, MU05	Aktas & Saribiyik (1996), Candemir & Kara (2003), Dirickx (1994), Saribiyik (1999b, 2001, 2003b, c, d)
19.	<i>Eristalinus megacephalus</i> (Rossi, 1794)	MU01, MU02, MU03	Aktas & Saribiyik (1996; as <i>E.</i> <i>quinquelineatus</i>), Dirickx (1994)
20.	<i>Eristalinus sepulchralis</i> (Linnaeus, 1758)	AF01, MU01, MU02, MU03, MU04, MU21	Aktas & Saribiyik (1996), Dirickx (1994), Saribiyik (1999b, 2001, 2003b, c, d)
21.	<i>Eristalinus taeniops</i> (Wiedemann, 1818)	MU01, MU03, MU05, MU06, MU11, MU14	Aktas & Saribiyik (1996), Dirickx (1994), Saribiyik (2003d)
22.	<i>Eristalis arbustorum</i> (Linnaeus, 1758)	AF01, MU01, MU02, MU03, MU05, MU10, MU15, MU18	Aktas & Saribiyik (1996), Candemir & Kara (2003), Dirickx (1994), Hayat & Alaoğlu (1990b), Hurkmans & Hayat (1997), Lindner (1941), Saribiyik (1999b, 2001, 2003b, c, d)
23.	<i>Eristalis similis</i> (Fallén, 1817)	MU10	Dirickx (1994; as <i>E. pratorum</i>), Saribiyik (1999b, 2001, 2003c, d; as <i>E. pratorum</i>)
24.	<i>Eristalis tenax</i> (Linnaeus, 1758)	AF01, AF02, AN03, MU01, MU03, MU05, MU09, MU10, MU15, MU16, MU18, MU19, TR03, MU20	Aktas & Saribiyik (1996), Candemir & Kara (2003), Dirickx (1994), Hayat & Alaoğlu (1990b), Hurkmans & Hayat (1997), Lindner (1941), Saribiyik (1999b, 2001, 2003b, c, d)
25.	<i>Eupeodes corollae</i> (Fabricius, 1794)	MU02, MU10, MU15, NE01	Candemir & Kara (2003), Dirickx (1994), Hayat & Alaoğlu (1990a), Lindner (1941), Özgür (1986), Saribiyik (1999a, 2000a, 2003b, c, e), Saribiyik & Aktas (1996)
26.	<i>Eupeodes flaviceps</i> (Rondani, 1857)	NE02 (3 m)	Dirickx (1994)
27.	<i>Eupeodes latifasciatus</i> (Macquart, 1829)	MU01, MU10, MU15	Dirickx (1994), Özgür (1986), Saribiyik (1999a, 2000a, 2003c, e)
28.	<i>Eupeodes luniger</i> (Meigen, 1822)	MU15 (1 m)	Dirickx (1994), Özgür (1986), Saribiyik (1999a, 2000a, 2003c)
29.	<i>Eupeodes nuba</i> (Wiedemann, 1830)	MU21 (1 m)	
30.	<i>Helophilus trivittatus</i> (Fabricius, 1805)	MU01, MU03	Aktas & Saribiyik (1996; as <i>H.</i> <i>parallelus</i>), Dirickx (1994), Saribiyik (1999b, 2001, 2003c, d; as <i>H.</i> <i>parallelus</i>)
31.	<i>Lejogaster tarsata</i> (Meigen, 1822)	AF01, NI02, MU07	Aktas & Saribiyik (1996), Dirickx (1994), Özgür (1987), Saribiyik (1999b, 2001, 2003b, c, d; as <i>L. splendida</i>)

32.	<i>Melanogaster nuda</i> (Macquart, 1829)	MU10, MU16	Dirickx (1994; as <i>Chrysogaster viduata</i>), Özgür (1987; as <i>Chrysogaster viduata</i>)
33.	<i>Melanostoma mellinum</i> (Linnaeus, 1758)	MU01, MU02, MU03, MU05, MU06, MU09, MU10, MU11, MU16, MU17, MU18, MU20, MU21	Candemir & Kara (2003), Dirickx (1994), Hayat & Alaoğlu (1990a), Özgür (1986), Saribiyik (1999a, 2000a, 2003b, c, e), Saribiyik & Aktas (1996)
34.	<i>Melanostoma scalare</i> (Fabricius, 1794)	MU11	Dirickx (1994), Saribiyik (1999a, 2000a, e)
35.	<i>Merodon aeneus</i> Meigen, 1822	MU10 (1 f)	
36.	<i>Merodon albifrons</i> Meigen, 1822	MU10 (1 m)	Dirickx (1994)
37.	<i>Merodon alexeji</i> Paramonov, 1925	MU16	Hurkmans & Hayat (1997)
38.	<i>Merodon avidus</i> Rossi, 1790; form A sensu Milankov et al. 2001	MU15 (5 m), MU16 (3 m), MU17 (1 m, 1 f)	Dirickx (1994), Hurkmans (1988, 1993), Hurkmans & Hayat (1997)
39.	<i>Merodon cf. geniculatus</i> Strobl, 1909	MU03 (1 m)	Dirickx (1994)
40.	<i>Merodon loewi</i> Van der Goot, 1964	NE03 (1 m)	Hurkmans (1988), Saribiyik (2003b, c, d)
41.	<i>Merodon longicornis</i> Sack, 1913	MU04 (1 m)	Hurkmans (1993), Saribiyik (2003b, d)
42.	<i>Merodon minutus</i> Strobl, 1893	MU03 (1 m), MU16 (1 f)	Dirickx (1994)
43.	<i>Merodon spinitarsis</i> Paramonov, 1929	MU03 (1 m)	Dirickx (1994)
44.	<i>Merodon telmateia</i> Hurkmans, 1987	MU03 (9m, 6 f)	Dirickx (1994), Hurkmans (1987)
45.	<i>Merodon velox</i> Loew, 1869	MU03 (1 f), MU05 (1 m)	Dirickx (1994), Hurkmans (1993), Hurkmans & Hayat (1997), Saribiyik (1999b, 2001, 2003b, c, d)
46.	<i>Mesembrius peregrinus</i> (Loew, 1846)	AF01, MU01, MU02	
47.	<i>Microdon devius</i> (Linnaeus, 1758)	TR01 (1)	Dirickx (1994), Özgür & Saribiyik (2002)
48.	<i>Milesia crabroniformis</i> (Fabricius, 1775)	MU01 (1)	Dirickx (1994), Saribiyik (2003d)
49.	<i>Milesia semiluctifera</i> (Villers, 1789)	MU01 (11)	Aktas & Saribiyik (1996), Dirickx (1994), Saribiyik (1999b, 2003c, d)
50.	<i>Myathropa florea</i> (Linnaeus, 1758)	MU01, MU02, MU05, MU10	Aktas & Saribiyik (1996), Candemir & Kara (2003), Dirickx (1994), Saribiyik (1999b, 2001, 2003c, d)
51.	<i>Neoscasia annexa</i> (Meigen, 1822)	MU15 (9 m, 3 f)	

52.	<i>Neoascia interrupta</i> (Meigen, 1822)	KA01 (3 f)	
53.	<i>Neoascia podagrica</i> (Fabricius, 1775)	MU01, MU10, MU16, MU17	Candemir & Kara (2003), Dirickx (1994), Lindner (1941), Özgür (1987), Saribiyik (1999b, 2001, 2003b, c, d)
54.	<i>Orthonevra frontalis</i> (Loew, 1843)	NI02 (1 m), MU17 (8 m, 6 f)	Aktas & Saribiyik (1996), Dirickx (1994), Özgür (1987), Saribiyik (2003b)
55.	<i>Orthonevra nobilis</i> (Fallén, 1817)	MU10 (1 m)	Dirickx (1994), Saribiyik (2001, 2003b, c)
56.	<i>Paragus abrogans</i> Goeldlin de Tiefenau, 1971	MU18 (1m)	Hayat & Alaoğlu (1990a)
57.	<i>Paragus bicolor</i> (Fabricius, 1794)	MU18 (1 m)	Candemir & Kara (2003), Dirickx (1994), Hayat & Alaoğlu (1990a), Özgür (1986), Saribiyik (1999a, 2000a, 2003b, c, e), Saribiyik & Aktas (1996)
58.	<i>Paragus compeditus</i> Wiedemann, 1830	MU01 (1 m)	Dirickx (1994), Hayat & Claussen (1997), Saribiyik (2003b)
59.	<i>Paragus haemorrhous</i> Meigen, 1822	MU01, MU02, MU03, MU07, MU10, MU16, MU18, MU21	Hayat & Claussen (1997)
60.	<i>Paragus pecchiolii</i> Rondani, 1857	MU01, MU02, MU18	Dirickx (1994; as <i>P. majoranae</i>)
61.	<i>Paragus quadrfasciatus</i> Meigen, 1822	AF01, MU05	Candemir & Kara (2003), Saribiyik (2000a, 2003b, c)
62.	<i>Paragus romanicus</i> Stănescu, 1992	MU15 (1 m)	Hayat & Claussen (1997)
63.	<i>Paragus tibialis</i> (Fallén, 1817)	MU01, MU09, NI02	Candemir & Kara (2003), Dirickx (1994), Hayat & Alaoğlu (1990a), Özgür (1986), Saribiyik (1999a, 2000a, 2003b, c, e), Saribiyik & Aktas (1996)
64.	<i>Parhelophilus versicolor</i> (Fabricius, 1794)	AF01, MU01, MU02, MU21	Dirickx (1994)
65.	<i>Pipiza festiva</i> Meigen, 1822	IC02	
66.	<i>Platycheirus fulviventris</i> (Macquart, 1828)	MU01, MU04	Dirickx (1994)
67.	<i>Platycheirus scutatus</i> (Meigen, 1822)	MU18 (1 m)	Saribiyik & Hasbenli (1997), Saribiyik (2000a, c)
68.	<i>Riponnensia insignis</i> (Loew, 1843)	MU01 (17 m, 23 f)	Dirickx (1994), Özgür (1987; as <i>Chrysogaster insignis</i>)
69.	<i>Riponnensia longicornis</i> (Loew, 1843)	MU01 (2 f)	Dirickx (1994), Özgür (1987), Saribiyik (2003b; as <i>Orthonevra longicornis</i>)
70.	<i>Scaeva albomaculata</i> (Macquart, 1842)	GA01 (1 m, 1 f)	Candemir & Kara (2003), Dirickx (1994), Özgür (1986), Hayat & Alaoğlu (1990a), Saribiyik (2000a, 2003b, c, e), Saribiyik & Aktas (1996)

71.	<i>Sphaerophoria rueppelli</i> (Wiedemann, 1830)	MU01, MU03, MU06, MU08, MU11, MU16, MU18	Candemir & Kara (2003), Dirickx (1994), Özgür (1986), Saribiyik (1999a, 2000a, 2003b, c, e)
72.	<i>Sphaerophoria scripta</i> (Linnaeus, 1758)	AR01, IS01, MU01, MU02, MU03, MU04, MU05, MU06, MU09, MU10, Mu11, MU15, MU16, MU17, MU18, MU19, TR03	Candemir & Kara (2003), Dirickx (1994), Hayat & Alaoğlu (1990a), Lindner (1941), Özgür (1986), Saribiyik (1999a, 2000a, 2003b, c, e), Saribiyik & Aktas (1996)
73.	<i>Sphaerophoria turkmenica</i> Bankowska, 1964	GA01 (1 m)	Hayat & Alaoğlu (1990a)
74.	<i>Sphegina sibirica</i> Stackelberg, 1953	TR03	
75.	<i>Sphiximorpha subseasilis</i> (Illiger, 1807)	IC02 (1 f), MU01 (3 m, 2 f)	
76.	<i>Syrirta flaviventris</i> Macquart, 1842	MU01, MU03, MU05, AN02	Dirickx (1994)
77.	<i>Syrirta pipiens</i> (Linnaeus, 1758)	AF01, KA01, MU01, MU02, MU03, MU04, MU05, MU06, MU08, MU09, MU10, Mu11, MU12, MU15, MU16, MU18, MU19	Aktas & Saribiyik (1996), Candemir & Kara (2003), Dirickx (1994), Hayat & Alaoğlu (1990b), Lindner (1941), Özgür (1987), Saribiyik (1999b, 2001, 2003b, c, d)
78.	<i>Volucella bombylans</i> (Linnaeus, 1758)	TR03 (1)	Dirickx (1994), Saribiyik (2001, 2003c)
79.	<i>Volucella pellucens</i> (Linnaeus, 1758)	TR02 (1)	Saribiyik (2001, 2003c)
80.	<i>Volucella zonaria</i> (Poda, 1761)	MU01, MU10, MU13	Aktas & Saribiyik (1996), Hurkmans & Hayat (1997), Özgür (1987), Saribiyik (1999b, 2001, 2003b, c, d)
81.	<i>Xanthogramma pedissequum</i> (Harris, 1776)	MU01, MU10, TR02	Özgür (1986; as <i>X. ornatum</i>), Saribiyik (1999a, 2000a, 2003b, c, e), Saribiyik & Aktas (1996)
82.	<i>Xylota ignava</i> (Panzer, 1798)	AR01 (1 m)	Dirickx (1994), Saribiyik & Özgür (2000)
83.	<i>Xylota segnis</i> (Linnaeus, 1758)	MU09 (1 m)	Dirickx (1994), Özgür (1987), Saribiyik (1999b, 2001, 2003c, d)
84.	<i>Xylota sylvarum</i> (Linnaeus, 1758)	MU01 (1 m, 1 f)	Dirickx (1994), Saribiyik (1999b, 2001, 2003c, d)

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Volucella - Die Schwebfliegen-Zeitschrift](#)

Jahr/Year: 2007

Band/Volume: [8](#)

Autor(en)/Author(s): Reemer Menno, Smit John T.

Artikel/Article: [Some hoverfly records from Turkey \(Diptera, Syrphidae\) 135-146](#)