On the Nomenclature of sundry Historids, including a note on a fourth species of European Dendrophilus.

By George Lewis in Wimbledon.

- 1. Saprinus venustus Er. is a species of Phoxonotus and by the description it apparently differs from P. tuberculatus Mars. and Fryi Lewis, in ten disposition of the tubercles on the thorax and by the arrangement of the elytral striae.
- 2. Saprinus fasciolatus Gebl. 1845, is evidently the same species as S. flexuoso fasciatus Motsch. 1845 and I would suggest that Gebler's name be given the priority. Between fasciolatus Gebl. and ornatus Er. (osiris Mars.) the chief differences lie in the shape of the prosternum, that of the first being truncate in front, with parallel lateral striae; and in the second it is somewhat rounded anteriorly while the striae approach each other in the middle.
- 3. Triballus 4-striatus Woll. = Gnathoncus rotundatus Kugel. The type in the British Museum has been carefully examined by Mr. Waterhouse and myself.
- 4. The Reverend Joh. Alf. Schmidt has lately obtained an example of the true Dendrophilus sulcatus Motsch. from Derbent in Russia and has kindly sent it to me for examination. It is very different to the Dendrophilus noticed by me in the Ann. and Mag. Nat. Hist., June 1885, pag. 467, and there considered to be Motschulsky's species. The last therefore requires a name and I propose to call it Dendrophilus Championi after Mr. G. C. Champion through whose kindness I have received several examples.
- 5. Onthophilus bipartitus Lewis = O. costipennis Mars. This I have been able to decide by the comparison of the types in Paris.
 - 10. August 1886.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Wiener Entomologische Zeitung

Jahr/Year: 1886

Band/Volume: 5

Autor(en)/Author(s): Lewis George

Artikel/Article: On the Nomenclature of sundry Histerids, including a note on a fourth species of European Dendrophilus. 280