

## **Geological substratum, tree vegetation, and floristic diversity of mature upland forest sites in northern New Jersey**

Donald W. Davidson

*Summary:* Data from twenty-nine stands of upland forests in northern New Jersey were used to determine the impact of geological substratum on species diversity of tree plants. Nine different substrata were studied, with more than three stands available on five of the substrata. The nine geological substrata were Kittatinny Limestone (24 species), Martinsburg Shale (22 species), Brunswick Shale (11), Triassic Conglomerate (15), Basalt (24), Longwood Shale (12), High Falls Sandstone (24), Green Pond Conglomerate (11), and Shawangunk Conglomerate (14). Species which are restricted to specific substrata are discussed. These studies encompass 40 tree species. These tree species are in general quite deeply rooted, and thus it would not be likely to have significant tree speciation taking place rapidly on specific substrata/soil types over 10,000 years since the Wisconsin Glaciation.

**Keywords:** geological substrata, tree vegetation, upland forest, forest flora of New Jersey, species diversity, distribution

Northern New Jersey is a region of considerable topographic diversity (700–800 feet in relief) and of extensive geological diversity, with geological substrata that include ten different types. These are Gneiss (Byrum, Pochuck and Losee), Triassic Shale, Triassic Conglomerate, Kittatinny Limestone, Basalt, Martinsburg Shale, Shawangunk Conglomerate, Green Pond Conglomerate, Longwood Shale and High Falls Sandstone. In these studies the field work was conducted from August 1959 until August 1963 with subsequent lab and herbarium studies continuing up until the present time, including the fall of 2004. Voucher specimens are on file in the John W. Thompson Herbarium, University of Wisconsin-Superior. The Geomorphic Provinces covered in this study are: (1) The Ridge and Valley Province, (2) The Highlands Province, and (3) The Piedmont Province. The valleys in these regions have elevations of 130 to 140 meters with a few ridges rising to higher elevations.

Glaciation has had a pronounced influence on northern New Jersey, and north of the Wisconsin Age Terminal Moraine. The surface is a sandy till, which is commonly thinner on ridge tops than on slopes and valley bottoms. Since the ice moved across New Jersey from northeast to southeast, the deposits are more or less of the same materials as the rocks they overlie. Drift older than the Wisconsin occurs south of the Wisconsin Moraine (LEWIS & KUMMEL 1940).

Twenty-nine stands of vegetation in the upland well-drained sites of northern New Jersey were studied and were connected primarily with floristic composition of trees, shrubs and herbs. The herb and shrub papers have been published (DAVIDSON 2004a,b), and thus were gathered by a combination of the Point Quarters Method (COTTAM & CURTIS 1956) and the floristic survey was also conducted within each of the twenty-nine stands studied. Quantitative importance values were computed, but it has been felt that the best approach was floristic presence study computed on each of the ten substrata, and then to compare the presence of the twenty-nine stands on the total number of sties (stands) for each substratum. This has also

D. W. DAVIDSON

been done for the twenty-nine stands of herbs and shrubs (see DAVIDSON 2004a,b). The results are shown in tables 1–5.

Table 1: Total number of forest sites for each substratum type.

Kittatinny Limestone	3
Martinsburg Shale	5
Brunswick Formation	1
Triassic/Border Conglomerate	1
Basalt	5
Longwood Shale	2
Gneiss	(11)
no tree data in this paper	
High Falls Sandstone	8
Green Pond Conglomerate	3
Shawangunk Conglomerate	1
	29

Table 2: Average number of trees per substratum.

Kittatinny Limestone	24
Martinsburg Shale	22
Brunswick Formation	11
Triassic/Border Conglomerate	15
Basalt	24
Longwood Shale	12
Gneiss	
no tree data in this paper	
High Falls Sandstone	24
Green Pond Conglomerate	11
Shawangunk Conglomerate	14

Table 3: Species of trees found on only one substratum.

Kittatinny Limestone	<i>Juglans nigra</i>
Martinsburg Shale	<i>Quercus palustris</i>
Brunswick Formation	no species restricted to this substratum
Triassic/Border Conglomerate	<i>Prunus avium</i>
Basalt	<i>Carpinus caroliniana</i>
Longwood Shale	no species restricted to this substratum
High Falls Sandstone	<i>Betula lutea</i>
Green Pond Conglomerate	no species restricted to this substratum
Shawangunk Conglomerate	<i>Acer pennsylvanica</i>
Franklin Marble	<i>Quercus muhlenbergii</i>
not part of the original substrate study	

## Geological substratum and tree vegetation in northern New Jersey

Table 4: Specifically of species of New Jersey trees from 29 stands, in search of specific tree/substrate relationships (GLEASON &amp; CRONQUIST 1991).

<i>Acer saccharum</i>	Rich to fairly dry woods, especially in calcareous soils
<i>Carya ovata</i>	Rich, moist soil
<i>Fagus grandifolia</i>	Rich, upland soils
<i>Fraxinus americana</i>	Rich, moist soils
<i>Juglans cinerea</i>	Rich, moist soils
<i>Juglans nigra</i>	Rich, moist soils
<i>Liriodendron tulipifera</i>	Rich woods
<i>Quercus muhlenbergii</i>	In good, chiefly calcareous soils (According to PEARSON (1960a,b, 1962), yellow oak is found in New Jersey only on Franklin Marble, but I have it on one site elsewhere)
<i>Tilia americana</i>	Rich woods
<i>Ulmus americana</i>	Usually in moist, fertile soil

Some comments on types of trees found on only one substratum.

## Kittatinny Limestone:

*Juglans nigra* is commonly thought to be found only on limestone in New Jersey (PEARSON 1960a,b, 1962; Davidson, unpubl. observations) so it is expected to be restricted to limestone in New Jersey. *Ulmus* spp. would not be, as it is a combination of the two species *U. rubra* and *U. americana*.

## Martinsburg Shale:

*Quercus palustris* is found only on Martinsburg Shale, but it is somewhat of a weedy species and thus would not be expected to be restricted to Martinsburg Shale by evolution.

## Triassic/Border conglomerate:

*Prunus avium* is a weedy species and an exotic from Europe, and thus would not be expected to be restricted to this substrate in this study.

## Basalt:

*Carpinus caroliniana* is found only on Basalt, and is found widely in other areas in eastern United States, and thus would not be expected to be found only on Basalt in northern New Jersey.

## High Falls Sandstone:

*Betula lutea* is found only on High Falls Sandstone in the study areas, but is found extensively throughout eastern U.S. on other substrata, and thus would not be expected to be restricted to just High Falls Sandstone in this study. It is more of a northern New Jersey species where it occurs.

## Shawangunk Conglomerate:

*Acer pennsylvanica* is found only on Shawangunk Conglomerate in northern New Jersey, but it is found in just one stand in the study region. It may simply be an outlier from the northern forest of adjacent Northeast Pennsylvania and northern New York to the north.

The three substrata with no trees restricted to them were Brunswick Formation, Longwood Shale, and Green Pond Conglomerate.

## D. W. DAVIDSON

Table 5: Trees found on specific substrata.

Kittatinny Limestone	<i>Acer rubrum</i>	<i>Carya tomentosa</i>	<i>Ostrya virginiana</i>	<i>Tsuga canadensis</i>
	<i>Acer saccharum</i>	<i>Cornus florida</i>	<i>Pinus strobus</i>	<i>Ulmus americana</i>
	<i>Betula lenta</i>	<i>Fraxinus americana</i>	<i>Populus grandidentata</i>	<i>Ulmus rubra</i>
	<i>Carya cordiformis</i>	<i>Juglans cinerea</i>	<i>Quercus alba</i>	<i>Quercus velutina</i>
	<i>Carya ovalis-glabra</i>	<i>Juglans nigra</i>	<i>Quercus prinus</i>	<i>Tilia americana</i>
	<i>Carya spp.</i>	<i>Liriodendron tulipifera</i>	<i>Quercus rubra</i>	
Martinsburg Shale	<i>Acer rubrum</i>	<i>Carya spp.</i>	<i>Nyssa sylvatica</i>	<i>Quercus prinus</i>
	<i>Acer saccharum</i>	<i>Carya tomentosa</i>	<i>Ostrya virginiana</i>	<i>Quercus rubra</i>
	<i>Betula lenta</i>	<i>Cornus florida</i>	<i>Quercus alba</i>	<i>Quercus velutina</i>
	<i>Carya cordiformis</i>	<i>Fagus grandifolia</i>	<i>Quercus coccinea</i>	<i>Sassafras albidum</i>
	<i>Carya ovalis-glabra</i>	<i>Fraxinus americana</i>	<i>Quercus palustris</i>	<i>Tsuga canadensis</i>
	<i>Carya ovata</i>	<i>Liriodendron tulipifera</i>		
Brunswick Formation	<i>Acer rubrum</i>	<i>Cornus florida</i>	<i>Quercus alba</i>	<i>Quercus rubra</i>
	<i>Betula lenta</i>	<i>Fagus grandifolia</i>	<i>Quercus coccinea</i>	<i>Quercus velutina</i>
	<i>Carya ovalis-glabra</i>	<i>Fraxinus americana</i>	<i>Quercus prinus</i>	
Triassic/Border Conglomerate	<i>Acer rubrum</i>	<i>Fagus grandifolia</i>	<i>Prunus serotina</i>	<i>Quercus rubra</i>
	<i>Betula lenta</i>	<i>Liriodendron tulipifera</i>	<i>Quercus alba</i>	<i>Quercus velutina</i>
	<i>Betula populifolia</i>	<i>Nyssa sylvatica</i>	<i>Quercus coccinea</i>	<i>Sassafras albidum</i>
	<i>Cornus florida</i>	<i>Prunus avium</i>	<i>Quercus prinus</i>	
Basalt	<i>Acer rubrum</i>	<i>Carya tomentosa</i>	<i>Nyssa sylvatica</i>	<i>Quercus velutina</i>
	<i>Acer saccharum</i>	<i>Cornus florida</i>	<i>Ostrya virginiana</i>	<i>Sassafras albidum</i>
	<i>Betula lenta</i>	<i>Fagus grandifolia</i>	<i>Quercus alba</i>	<i>Tsuga canadensis</i>
	<i>Carpinus caroliniana</i>	<i>Fraxinus americana</i>	<i>Quercus coccinea</i>	<i>Tilia americana</i>
	<i>Carya ovalis-glabra</i>	<i>Juglans cinerea</i>	<i>Quercus prinus</i>	<i>Ulmus americana</i>
	<i>Carya ovata</i>	<i>Liriodendron tulipifera</i>	<i>Quercus rubra</i>	<i>Ulmus rubra</i>
Longwood Shale	<i>Acer rubrum</i>	<i>Cornus florida</i>	<i>Nyssa sylvatica</i>	<i>Quercus rubra</i>
	<i>Acer saccharum</i>	<i>Fagus grandifolia</i>	<i>Quercus coccinea</i>	<i>Quercus velutina</i>
	<i>Betula lenta</i>	<i>Liriodendron tulipifera</i>	<i>Quercus prinus</i>	<i>Sassafras albidum</i>
Gneiss	no data available			
High Falls Sandstone	<i>Acer rubrum</i>	<i>Carya ovata</i>	<i>Ostrya virginiana</i>	<i>Quercus prinus</i>
	<i>Acer saccharum</i>	<i>Cornus florida</i>	<i>Prunus serotina</i>	<i>Quercus rubra</i>
	<i>Amelanchier spp.</i>	<i>Fagus grandifolia</i>	<i>Populus grandidentata</i>	<i>Quercus velutina</i>
	<i>Betula lenta</i>	<i>Fraxinus americana</i>	<i>Pinus rigida</i>	<i>Sassafras albidum</i>
	<i>Betula lutea</i>	<i>Liriodendron tulipifera</i>	<i>Pinus strobus</i>	<i>Tsuga canadensis</i>
	<i>Carya ovalis-glabra</i>	<i>Nyssa sylvatica</i>	<i>Quercus coccinea</i>	
Green Pond Conglomerate	<i>Acer rubrum</i>	<i>Prunus serotina</i>	<i>Quercus prinus</i>	<i>Sassafras albidum</i>
	<i>Betula lenta</i>	<i>Populus grandidentata</i>	<i>Quercus rubra</i>	<i>Tsuga canadensis</i>
	<i>Carya ovalis-glabra</i>	<i>Quercus coccinea</i>	<i>Quercus velutina</i>	
Shawangunk Conglomerate	<i>Acer pennsylvanica</i>	<i>Betula lutea</i>	<i>Prunus serotina</i>	<i>Quercus prinus</i>
	<i>Acer rubrum</i>	<i>Betula populifolia</i>	<i>Pinus rigida</i>	<i>Quercus rubra</i>
	<i>Acer saccharum</i>	<i>Carya ovalis-glabra</i>	<i>Pinus strobus</i>	<i>Quercus velutina</i>
	<i>Amelanchier arborea</i>	<i>Nyssa sylvatica</i>	<i>Quercus alba</i>	<i>Sassafras albidum</i>
	<i>Amelanchier spp.</i>	<i>Ostrya virginiana</i>	<i>Quercus coccinea</i>	<i>Tsuga canadensis</i>
	<i>Betula lenta</i>			

## Acknowledgements

Thanks are expressed to Dr. Mary Balcer for her ongoing support of these studies, and to Sandy Orr, Jon Petersburg and Patser Lynch for their support. These studies were funded by N.S.F. Grant G-7177 (Rutgers University), and by the Board of Regents of the University of Wisconsin System.

## References

- COTTAM, G. & CURTIS, J. T. (1956):** The use of distance measures in phytosociological sampling. – *Ecology* **37**: 451–460.
- DAVIDSON, D. W. (2004a):** Geological substratum, herbaceous vegetation and floristic diversity of mature upland forest sites in northern New Jersey. – *Wulfenia* **11**: 45–55.
- DAVIDSON, D. W. (2004b):** Geological substratum, shrub vegetation, and floristic diversity of mature upland forest sites in northern New Jersey. – *Wulfenia* **11**: 57–61.
- GLEASON, H. A. & CRONQUIST, A. (1991):** Manual of vascular plants of northeastern United States and adjacent Canada. – New York: New York Botanical Garden Press.
- LEWIS, J. V. & KUMMEL, H. B. (1940):** The geology of New Jersey. – New Jersey Dep. Conserv. Development. Series Bull. **50**: 1–203.
- PEARSON, P. R. (1960a):** New Jersey limestone vegetation. – Ph. D. Thesis. Rutgers University, New Brunswick, New Jersey: Univ. Microfilms, Ann Arbor, Michigan.
- PEARSON, P. R. (1960b):** Upland forests on the Kittatinny Limestone and Franklin Marble of northern New Jersey. – *Bull. N. J. Acad. Sci.* **5**: 3–19.
- PEARSON, P. R. (1962):** Increasing importance of sugar maple on two calcareous formations in New Jersey. – *Ecology* **43**: 711–718.

Address of the author:

Prof. Dr. Donald W. Davidson  
Department of Biology and Lake Superior Research Institute  
University of Wisconsin-Superior  
Belknap & Catlin PO Box 2000  
Superior, WI 54880-4500  
U.S.A.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Wulfenia](#)

Jahr/Year: 2005

Band/Volume: [12](#)

Autor(en)/Author(s): Davidson Donald W.

Artikel/Article: [Geological substratum, tree vegetation, and floristic diversity of mature upland forest sites in northern New Jersey 87-91](#)