

Zeitschrift für Malakozoologie.

Herausgegeben

von

Karl Theodor Menke, M. D.

und

Dr. Louis Pfeiffer.

1847.

Vierter Jahrgang.

Juni.

Diagnosen neuer Heliceen.

1. *Helix Besckei* Dunker.

T. anguste umbilicata, orbiculato-convexa, laevigata, lineis impressis, radiantibus notata, nitidissima, corneolutea; spira subconoideo-elevata; anfr. $5\frac{1}{2}$ vix convexiusculi, regulariter accrescentes; apertura parum obliqua, lata, lunaris; perist. simplex, rectum, acutum, margine columellari brevi, verticali, subreflexo. — Diam. maj. 7, min. 6, alt. $2\frac{2}{3}$ mill.

Patria: Brasilia (Bescke).

2. *Helix Tamsiana* Dunker.

T. mediocriter umbilicata, depresso conica, granulata, fusca, acute carinata; spira conoidea, apice obtusiuscula; anfr. 5 subplani, ultimus basi fere planus, antice angulatim deflexus, ad aperturam constrictus; apertura subhorizontalis, auriformis, quadridentata; perist. tenue, marginibus lamina callosa, umbilici partem tegente et medio dentem linguiformem intrantem emittente junctis, supero expanso, unidentato, basali reflexo, dentibus 2 fere aequalibus munito. — Diam. maj. 15, min. 14, alt. $7\frac{1}{2}$ mill.

Patria: Porto Cabello Venezuelae (Dr. Tams).

3. *Helix leucodon* Pfr.

T. anguste umbilicata, depressa, sublenticularis, undique

subtiliter granulata, saturate castanea; spira plane elevata, vertice obtuso; anfr. 5 subplanulati, ultimus carinatus, antice gibbus, ad aperturam subito deflexus, constrictus, subtus scrobiculatus; apertura perobliqua, irregulariter pyriformis; perist. tenue, album, expansum, reflexiusculum, marginibus lamina elevata, flexuosa, medio lamellam erectam, linguiformem emittente junctis, supero dente 1 parvulo, basali 2—3 (altero infra carinam acuto, secundo obtusiore, tertio subobsoleto) munitis. — Diam. maj. 21, min. 19, alt. 10 mill.

Patria?

3. *Bulimus gonochilus* Pfr.

T. subperforata, ovato-conica, tenuis, subtiliter et confertim decussata, diaphana, virens; spira conica, acutiuscula; anfr. 5 vix convexiusculi, ultimus spiram superans; columella superne obsolete plicata; apertura oblonga, oblique truncata; perist. angulatim late expansum, reflexiusculum, tenue, intus candidum, marginibus distantibus. — Long. 17, diam. 9 mill.

Patria . . .

5. *Bulimus Recluzianus* Pfr.

T. subperforata, subfusiformi-oblonga, sublaevigata, lineis confertis spiralibus subtiliter sculpta, nitida, opaca, carnea, strigis latis, irregularibus, castaneis et griseo-violaceis ornata; spira conica, apice obtusiuscula; sutura albo-marginata; anfr. 6 convexiusculi, ultimus spiram aequans; columella tenuis, superne subtorta, leviter arcuata; apertura oblonga, subverticalis; perist. simplex, acutum, margine columellari superne reflexo, subadnato. — Long. 30, diam. 12 mill. Ap. 16 mill. longa, $7\frac{1}{2}$ lata.

Patria?

6. *Bulimus amoenus* Pfr.

T. subperforata, fusiformi-oblonga, striata, opaca, flavescens, roseo-nebulosa, fasciis 4 spadiceis et quinta

basali rubra ornata; spira turrito-conica, apice acutiuscula; anfr. 6 planiusculi, ultimus spiram aequans; columella superne compressa, subtorta, deorsum leviter arcuata; apertura ovali-oblonga, intus concolor; perist. simplex, rectum, margine dextro subrependo, columellari breviter reflexo, subadnato. — Long. 29, diam. 10 mill. Ap. 15 mill. longa, medio 7 lata.

Patria?

7. *Bulimus ringens* Dunk.

T. rimato-subperforata, fusiformis, tenuis, laevigata, pallide cornea, fusco-strigata, vel cornea, epidermide cinerea strigatim obducta; spira turrita, acutiuscula; anfr. 7—7½ planiusculi, ultimus ⅓ longitudinis vix superans; apertura angusta, oblonga, 4 dentata: dente 1 in pariete aperturali, secundo in columella plicata, tertio in basi, quarto in medio marginis dextri; perist. album, expansum. — Long. 18, diam. 5⅓ mill. Ap. 7 mill. longa, 4 lata.

Patria: Brasilia. (Rarissimus teste Bescke.)

8. *Pupa calcarea* Pfr.

T. profunde rimata, subcylindracea, solida, irregulariter striata, nitida, calcarea; spira elongata, apice obtuse conica; anfr. 12 planiusculi, ultimus ascendens, inflatus, antice rugoso-plicatus; columella simplex; apertura semiovalis, intus alba, in fundo fuscula; perist. valde incrassatum, undique expansum, album, marginibus callo nitido junctis, columellari superne sinuato-reflexo. — Long. 38, diam. 12 mill. Ap. c. perist. 14 mill. longa, 12 lata.

Patria . . .

9. *Pupa marmorata* Pfr.

T. rimato-perforata, subcylindraceo-ovata, solida, striatula, nitida, nivea, flammis longitudinalibus castaneis, diaphanis ornata; spira sursum sensim attenuata, apice obtusa; anfr. 10 subplanulati, ultimus antice plicatus; columella obsolete et profunde unidentata; plica parietis

aperturalis profunda, medioeris; apertura semiovalis, intus livido-fusca; perist. incrassatum, reflexum, marginibus superne convergentibus. — Long. 24, diam. vix 10 mill. Ap. 9 mill. longa, 8 lata.

Patria

10. *Balea Sarsii Phil.*

T. sinistrorsa, subperforata, oblongo-turrita, striatula, nitida, olivaceo-cornea, pellucida; spira acutiuscula; anfr. 7 convexi, ultimus basi rotundatus; columella substricta, recedens; apertura oblongo-semiovalis; perist. simplex, expansiusculum, margine sinistro sinuoso-dilatato. — Long. 6, diam. $2\frac{1}{4}$ mill.

Patria: Norvegia. Misit Cl. Sars.

Testaceorum novorum centuria.

Descripsit Dr. A. R. Philippi.

(Continuatio.)

15. *Donax Hanleyana Ph.*

D. testa oblonga, pallida, valde inaequilatera, acute carinata, anterie omnino laevi, versus carinam striis impressis radiantibus sculpta; area postica medio distincte angulata, in parte inferiore lineis elevatis radiantibus quinque granosis, in parte superiore liris elevatis concentricis, striisque radiantibus impressis decussata; margine dentato, in angulo postico spinoso; dentibus lateralibus validis. Long. $16\frac{1}{2}$ ''' ; alt. 10''' ; crass. 7'''.

Patria?

Forma cum D. elongata convenit, sed carina distinctior, acutangula, sculptura areae longe grossior, dentes marginis duplo pauciores, ad summum 50, in angulo an-

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Zeitschrift für Malakozoologie](#)

Jahr/Year: 1847

Band/Volume: [4_1847](#)

Autor(en)/Author(s): diverse

Artikel/Article: [Diagnosen neuer Heliceen 81-84](#)