

Z. Säugetierkunde 54 (1989) 239–242
© 1989 Verlag Paul Parey, Hamburg und Berlin
ISSN 0044-3468

Weight estimation of Spanish ibex, *Capra pyrenaica*, and Chamois, *Rupicapra rupicapra* (Mammalia, Bovidae)

By P. FANDOS, C. R. VIGAL and J. M. FERNANDEZ-LOPEZ

Estación Biológica de Doñana, Sevilla and Unidad de Zoología Aplicada, Alcala de Henares, Madrid

Receipt of Ms. 20. 11. 1987

Abstract

The relationships of body weight to horn length, chest girth and body length were analysed for Spanish ibex, *Capra pyrenaica* (Schinz, 1838), and Chamois, *Rupicapra rupicapra* (L., 1758). Different linear regressions were necessary for each population and for each sex. All relationships were statistically significant ($p < 0.01$). The best predictors of body weight were horn length for male Spanish ibex, body length for female Spanish ibex, and chest girth for both sexes in the chamois.

Introduction

In recent studies of ungulates (GEIST 1966; NIEVERGELT 1966; BUNNELL 1978; CLUTTON-BROCK et al. 1980) great importance has been placed on the relationship between horn length and body size and its ecological, ethological, and evolutionary implications. However, for control or populations management purposes, the relationships between horn length and body weight is also valuable. Nevertheless, since reliable data on body weight is frequently absent in museum specimens or mounted trophies, other linear measurements related to body size, like horn length, have been studied in several ungulates by a large number of authors, among them ROSS (1958), BLOOD and LOVAAS (1966), McEWAN and WOOD (1966), NIEVERGELT (1966), RIDEOUT and WORTHEN (1975), HALL MARTIN (1977), GRAY and SIMPSON (1979), BUNNELL (1980), CLUTTON-BROCK et al. (1980), and POPP (1985).

Although difficulties occur when estimating weight from linear measurements (RIDEOUT and WORTHEN 1975), and sometimes low correlations (BUNNELL 1978), this paper shows that the relationships between body weight and various linear measurements can be used to predict body weight in two populations of Spanish ibex, *Capra pyrenaica* (Schinz, 1838) and one of Chamois *Rupicapra rupicapra* (L., 1757).

Material and methods

Ibex specimens ($n = 169$) came from the Sierra de Cazorla, Southeast of the Iberian peninsula, and also from the Sierra de Gredos, Central Mountain Range, two of the main places within the area of distribution of the species. The chamois specimens ($n = 47$) came from the Cantabrian Mountains, Northwest of the Iberian peninsula. In both species specimens were obtained for a broad study of their biology and ecology from shootings during 1980–1984. Attention was paid to obtain a representative sample of each population in terms of sex, age and seasonality.

The specimens were weighed on a platform scale with a precision of ± 0.1 kg. Horn length (HORN) was measured by using a cord which was applied closely to the convoluted surface of the horn from tip to base. This measurement provided a better estimator of horn length than the simple measurement of the shortest distance between the base of the horn and its tip. Body length (LCC) was measured from the extreme of the snout to the base of the tail using a metric tape. During this procedure the stretched-out neck was bent so that it lay on the same plane as the trunk. The chest girth (PTOR) was taken by placing the tape around the body behind the front legs. All measurements were made in millimeters with a precision of ± 1 mm.

sexual selection which favors the development of secondary sexual characteristics, mainly related to fighting (SCHAFER and REED 1972; CLUTTON-BROCK et al. 1980).

The relationship between horn-length and body weight gives a higher correlation index in males than in females of the two species considered. NIEVERGELT (1966) found similar results for the Alpine ibex where the length of the horn showed the greatest correlation with body weight of males.

Regressions between horn length and body weight have been reported in other bovids (GRAY and SIMPSON 1979; BUNNELL 1980; POPP 1985). In this study, differences in the degree of fitness of horn length/body weight regression were found. These differences can be seen at three levels; specific (Ibex specimens showing higher fitness than the Chamois), sexual (males having higher fitness than females), or population in Ibex females specimens, from Gredos being better fitted than those from Cazorla. On the other hand, other studies (FANDOS 1986; FANDOS and VIGAL in press) suggested that the annual growth rate also shows differences connected with species, sex or population. Thus, we conclude from these two facts, that annual growth rate could be related to the degree of fitness of this regression.

A second reason why ibex shows a high correlation between body weight and horn length, can be explained by the methods used in the present study. The sample of specimens studied was considered to be sufficiently representative of the population concerning sex and age aspects, trying to avoid the error described by BUNNELL (1980) which occurs when all the age groups are not included in the sample. It must also be pointed out that these specimens were collected throughout the whole year, thus avoiding seasonal variations (GRAY and SIMPSON 1979).

Chest girth measurement (PTOR) provides the best estimate of body weight in the chamois in both sexes. Similar results for this species have been obtained by SCHROEDER and REDLICH (1976). However, R^2 values between these two measurements (0.86 and 0.68; table) is lower than the one obtained by BUNNELL (1980) for the mountain goat (*Oreamnus americanus*).

In chamois, when estimating body weight from the three variables, R^2 value for females was lower than for males (Table). This fact may be related to the higher energetic expenses of reproduction (gestation and suckling) in females, as suggested by GRAY and SIMPSON (1979) to explain the rates of development of horns in the Barbary sheep (*Ammotragus lervia*).

Acknowledgements

We would like to thank B. COOK and S. REIG for their help in translating the manuscript into an English version.

Zusammenfassung

Gewichtsabschätzungen für den spanischen Steinbock, Capra pyrenaica und die Gemse, Rupicapra rupicapra (Mammalia, Bovidae)

Es wurde das Verhältnis zwischen Körpergewicht und den Maßen für Hornlänge, Brustumfang und Körperlänge für den spanischen Steinbock, *Capra pyrenaica* (Schinz, 1838) und die Gemse, *Rupicapra rupicapra* (L., 1758), analysiert. Für jede Population und für beide Geschlechter waren getrennte Berechnungen linearer Regressionen notwendig. Alle Beziehungen sind statistisch signifikant ($p < 0.01$). Die besten Voraussagen über das Körpergewicht erlauben für den männlichen Steinbock die Hornlänge, für den weiblichen Steinbock die Körperlänge sowie der Brustumfang für beide Geschlechter der Gemse.

References

- BLOOD, D. A.; LOVAAS, A. L. (1966): Measurements and weight relationships in Manitoba elk. J. Wildl. Manage. 30, 135–140.
BUNNELL, F. L. (1978): Horn growth and population quality in Dall sheep. J. Wildl. Manage. 42, 755–764.

- (1980): Weight estimation of Dall's sheep and mountain goat. *Wildl. Soc. Bull.* **8**, 291–297.
- CLUTTON-BROCK, T. H.; ALBON, S. D.; HARVEY, P. H. (1980): Antlers, body size and breeding group size in the Cervidae, *Nature* **258**, 565–567.
- FANDOS, P. (1986): Aspectos ecológicos de la población de cabra montés en las Sierras de Cazorla y Segura, Jean. Thesis Doc. Universidad Complutense, Madrid, 1–475 p.
- FANDOS, P.; VIGAL, C. R. (in press): Body weight and horn length in relation to age of the Spanish wild goat. *Acta theriol.*
- FERNANDEZ-LOPEZ, J. M.; GARCIA GONZALEZ, R. (1986): Craniométrie comparée entre les chamois pyrénéen et cantabrique. *Mammalia* **50**, 87–97.
- GEIST, V. (1966): The evolutionary significance of mountain sheep horns. *Evolution* **20**, 558–566.
- GRAY, G. C.; SIMPSON, C. D. (1979): Weight estimation of Babary sheep from horn length. *Wildl. Soc. Bull.* **7**, 285–288.
- HALL-MARTIN, A. J. (1977): Giraffe weight estimation using dissected leg weight and body measurements. *J. Wildl. Manage.* **41**, 740–745.
- MC EWAN, E. H.; WOOD, A. J. (1966): Growth and development of the barren ground caribou. I. Heart girth, hind foot length, and body weight relationships. *Can. J. Zool.* **44**, 401–411.
- NIEVERGELT, B. (1966): Der Alpensteinbock (*Capra ibex* L.) in seinem Lebensraum. *Mammalia depicta* 1. Hamburg, Berlin: Paul Parey.
- POPP, J. W. (1985): Horn and body size among antelopes. *Säugetierkde. Mitt.* **32**, 245–248.
- RIDEOUT, C. B.; WORTHEN, C. L. (1975): Use of girth measurement for estimating weight of mountain goats. *J. Wildl. Manage.* **39**, 705–708.
- ROSS, J. G. (1958): A method of estimating live weights in small shorthorn zebu cattle from linear body measurements. *E. Afr. Agric. J.* **23**, 193–194.
- SCHAFER, W. M.; REED, C. A. (1972): The co-evolution of social behavior and cranial morphology in sheep and goats (Bovidae, Caprini). *Fieldiana: Zoology* **61**, 1–88.
- SCHROEDER, W.; REDLICH (1976): Untersuchungen an der bayerischen Gams. I. Körpermaße und Gewichte. Tagungsbericht. 2nd. Conf. Internationales Gamswild. Ljubljana.

Authors' addresses: PAULINO FANDOS, Museo Nacional de Ciencias Naturales, c/Jose Gutierrez Abascal 2, E-28006 Madrid; CARLOS R. VIGAL and JOSE M. FERNANDEZ-LOPEZ, Unidad de Zoología Aplicada, Comunidad Autónoma de Madrid, Aptdo 127, El Encín, Alcalá de Henares, Madrid 1, España

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Mammalian Biology \(früher Zeitschrift für Säugetierkunde\)](#)

Jahr/Year: 1989

Band/Volume: [54](#)

Autor(en)/Author(s): Vigal Carlos R., Fandos Paulino, Fernandez-Lopez Jose M.

Artikel/Article: [Weight estimation of Spanish ibex, *Capra pyrenaica*^ and Chamois, *Rupicapra rupicapra* \(Mammalia, Bovidae\)&SÂámò 239-242](#)