

Z. Säugetierkunde 57 (1992) 112–115
© 1992 Verlag Paul Parey, Hamburg und Berlin
ISSN 0044-3468

WISSENSCHAFTLICHE KURZMITTEILUNGEN

Bats from Mana Pools National Park in Zimbabwe and the first record of *Eptesicus rendallii* from the country

By I. L. RAUTENBACH and M. B. FENTON

Transvaal Museum, Pretoria, South Africa and Department of Biology, York University, North York, Ontario, Canada

Receipt of Ms. 28. 10. 1991
Acceptance of Ms. 3. 12. 1991

The bat fauna of Africa is not well documented compared to other African mammals. For example, while the bats of Zimbabwe have received some attention (FENTON 1975; SMITHERS and WILSON 1979; HUTTON 1986), at most locations there has been no specific survey of bats. Even at Mana Pools National Park on the Zambezi River (15° 44' S; 29° 21' E), the site of some bat research (e.g. ALDRIDGE et al. 1990; FENTON et al. 1990; OBRIST et al. 1989), there is virtually no information about the bat fauna.

Between 18:00 and 21:00 we used 12 m mist nets (126 net hours in November 1987) and two macro-mist nets (36 net hours in June 1990; RAUTENBACH 1985) to sample bats at sites within 10 km of the headquarters of Mana Pools National Park. We sampled sites in the dry woodland some distance away from the river, and on the flood plain of the Zambezi River in the predominately *Acacia albida* woodland along the Zambezi River, a park-like habitat with dispersed tall canopy trees and little understory (DUNHAM 1989). Here we record the first record of *Eptesicus rendallii* from Zimbabwe and the occurrence of 20 species of bats in Mana Pools National Park (Table 1), 15 of which are recorded for the first time.

The bats recorded for Mana Pools National Park are compared with the species compositions of two other sites in southern Africa (Table 1). These bat faunas are typical of open woodland in southern Africa. *Nycticeius schlieffenii* and *Scotophilus borbonicus* together accounted for 80 % of the 136 bats netted at Mana Pools National Park between 18:00 and 21:00 over 12 nights in June 1990. In comparing the faunas (see Table), it is clear that a number of species are rarely encountered, while others are taken only in the immediate vicinity of their roosts. The larger species richness recorded for Pafuri, may be ascribed to greater ecological complexity and more intensive surveying.

Eptesicus rendallii is widespread in open woodland savannahs of subsaharan Africa (SMITHERS 1983) and ranges southwards into northern Botswana (SMITHERS 1971), central Zambia (ANSELL 1978), southern Malawi (ANSELL and DOWSETT 1988), and the Tete and Mocambique districts of Mocambique (SMITHERS and LOBAO TELLO 1976). Our specimens are the first from Zimbabwe and they have been prepared as standard skin and skull vouchers and deposited in the national mammal collection of the Transvaal Museum. In wing characteristics (aspect ratio 6.71 male, 6.18 female; wingloading 7.81 N m⁻² and 6.93 N m⁻²) and size (6.0 and 6.5 g) these bats would be included in the Group 1 of ALDRIDGE and RAUTENBACH (1987), species with high flight manoeuvrability that should forage in relatively cluttered situations.

We prepared standard bone marrow karyotypes from the two *E. rendallii* using the in vivo technique of ROBBINS and BAKER (1978) as modified by BAKER et al. (1982). At least

The bats captured at Mana Pools National Park compared with species recorded in the Sengwa Wildlife Area in Zimbabwe (18° 10' S; 28° 13' E) and the Pafuri area of the Kruger National Park in South Africa (22° 25' S; 31° 13' E)

While the actual numbers of bats from Mana Pools National Park are shown, for the other two sites relative abundance (for comparable sampling effort) is indicated as follows: * <3; + = 3–10; ++ = 11–20; +++ = 21–30; ++++ = 31–40, etc. The “r” identifies bats captured emerging from roost sites, not in nets set in woodland. The “a” is from SMITHERS and WILSON (1979); the “b” from FENTON et al. (1990); the “c” from ALDRIDGE et al. (1990); the “d” from HUTTON (1986); and the “e” is from THOMAS VOLPERS (pers. comm.)

Species	Mana Pools	Sengwa	Pafuri	Species	Mana Pools	Sengwa	Pafuri
<i>Epomophorus wahlbergi</i>	—	+	+	<i>Pipistrellus nanus</i>	1	++	+
<i>Epomophorus crypturus</i>	1	++	*	<i>Chalinolobus variegatus</i>	6	*	*
<i>Rousettus aegyptiacus</i>	+	+	++	<i>Laephotis botswanae</i>	—	+	*
<i>Taphozous mauritanicus</i>	+ ^a	*	*	<i>Eptesicus rendallii</i>	2	—	—
<i>Nycteris grandis</i>	+ ^b	—	—	<i>Eptesicus bottenotus</i>	—	—	*
<i>Nycteris woodii</i>	—	+	*	<i>Eptesicus cf. melckorum</i>	3	?	+
<i>Nycteris thebaica</i>	+ ^c	++	+	<i>Eptesicus zuluensis</i>	1	*	+
<i>Rhinolophus bildebrandtii</i>	1	++	+	<i>Eptesicus capensis</i>	4	+++	++
<i>Rhinolophus fumigatus</i>	—	+	+	<i>Scotophilus dinganii</i>	—	+	++++
<i>Rhinolophus clivosus</i>	—	*	*	<i>Scotophilus borbonicus</i>	52	++++	++++
<i>Rhinolophus darlingi</i>	1	—	*	<i>Nycticeius schlieffenii</i>	55	++++	++++
<i>Rhinolophus landeri</i>	* ^d	+ ^r	*	<i>Kerivoula argentata</i>	—	*	*
<i>Rhinolophus simulador</i>	—	+ ^r	*	<i>Kerivoula lanosa</i>	—	—	*
<i>Rhinolophus swynnii</i>	—	—	*	<i>Otomops mariesseni</i>	—	*	—
<i>Hipposideros commersoni</i>	1 ^e	+	*	<i>Tadarida fulminans</i>	—	*	*
<i>Hipposideros caffer</i>	—	+++ ^r	+++ ^r	<i>Tadarida brevittata</i>	—	+	—
<i>Glossotis percivali</i>	—	* ^r	—	<i>Tadarida ansoorgei</i>	—	—	*
<i>Miniopterus schreibersii</i>	1	+	*	<i>Tadarida nigeriae</i>	—	+	—
<i>Myotis welwitschii</i>	—	*	*	<i>Tadarida chapini</i>	—	++	—
<i>Myotis tricolor</i>	—	—	*	<i>Tadarida pumila</i>	7	—	+ ^r
<i>Myotis bocagii</i>	—	—	*	<i>Tadarida midas</i>	1	—	+ ^r
<i>Pipistrellus rueppellii</i>	1	++	+	<i>Tadarida condylura</i>	—	—	+ ^r
<i>Pipistrellus rusticus</i>	—	—	+++				

five representative chromosome spreads were examined from each bat to determine diploid (2n) and fundamental numbers (FN). Photomicrographic enlargements of suitable spreads were used in the final analyses and to prepare a karyogram.

Our data show that the standard karyotype of *E. rendallii* is $2n = 38$, $FN = 50$, and the autosomal complement is seven pairs of bi-armed chromosomes, and a size-graded series of 11 acrocentric pairs. A prominent secondary constriction is apparent near the centromere of one of the larger pairs of acrocentric chromosomes. The X chromosome is a large, submetacentric, and the Y a small submetacentric. Our findings agree with the earlier description of the karyotype of *E. rendallii* (McBEE et al. 1987), with the exception of the Y chromosome, which is described for the first time.

Some workers (e.g. HELLER and VOLLETH 1984) would group *E. rendallii* together with other karyotypically variable *Eptesicus*-like African bats, as *Pipistrellus* since they deviate from what is considered to be the typically conservative karyotype of 'true' *Eptesicus* ($2n = 50$). Based on bacular morphology others (e.g., HILL and HARRISON 1987) also consider *E. rendallii* as a *Pipistrellus*. However, the complexity of the problem makes it prudent to report our record as *Eptesicus rendallii*, pending the results of more in-depth systematic studies.

Acknowledgements

We thank the Director of the National Parks and Wildlife Management Department for permission, and Warden F. BUYEYE and C. JAKOPO for advice and assistance in the field. R. B. MARTIN, Assistant Director (Research) provided logistical help, while M. B. CUMMING, M. K. MUSGRACE, J. S. TAYLOR and T. VOLPERS helped in the field. This study was supported by grants from the Natural Sciences and Engineering Research Council of Canada to MBF and the South African Foundation for Research Development to ILR.

References

- ALDRIDGE, H. D. J. N.; OBRIST, M.; MERRIAM, H. G.; FENTON, M. B. (1990): Roosting, vocalizations and foraging by the African bat, *Nycteris thebaica*. *J. Mammalogy* 71, 242–246.
- ALDRIDGE, H. D. J. N.; RAUTENBACH, I. L. (1987): Morphology, echolocation and resource partitioning in insectivorous bats. *J. Animal Ecology* 56, 763–778.
- ANSELL, W. F. H. (1978): The mammals of Zambia. Chilanga: The National Parks and Wildlife Service.
- ANSELL, W. F. H.; DOWSETT, R. J. (1988): Mammals of Malawi: an annotated check list and atlas. Zennor: The Trendline Press.
- BAKER, R. J.; HAIDUK, M. W.; ROBBINS, L. W.; CADENA, A.; KOOP, B. F. (1982): Chromosomal studies of South American bats and their systematic implications. Special publ. Museum Texas Tech. Univ. 6, 303–327.
- DUNHAM, K. M. (1989): Long-term changes in Zambezi riparian woodlands, as revealed by photo-panoramas. *African J. Ecology* 27, 263–275.
- FENTON, M. B. (1975): Observations on the biology of some Rhodesian bats, including a key to the Chiroptera of Rhodesia. Life Sciences Contributions, Royal Ontario Museum 104, 1–27.
- FENTON, M. B.; SWANEPOEL, C. M.; BRIGHAM, R. M.; HICKEY, M. B. C. (1990): Foraging behaviour and prey selection by large slit-faced bats (*Nycteris grandis*; Chiroptera: Nycteridae). *Biotropica* 22, 2–8.
- HELLER, K.-G.; VOLLETH, M. (1984): Taxonomic position of "*Pipistrellus societatis*" Hill, 1972 and the karyological characteristics of the genus *Eptesicus* (Chiroptera: Vespertilionidae). *Z. zool. Syst. Evolut.-forsch.* 22, 65–77.
- HILL, J. E.; HARRISON, D. L. (1987): The baculum in the Vespertilioninae (Chiroptera: Vespertilionidae) with a systematic review, a synopsis of *Pipistrellus* and *Eptesicus* and the descriptions of a new genus and subgenus. *Bull. British Museum (Natural History), Zoology Series* 52, 225–305.
- HUTTON, J. M. (1986): The status and distribution of bats in Zimbabwe. *Cimbebasia* (A) 8, 219–236.
- MCBEE, K.; SCHLITTER, D. A.; ROBBINS, R. L. (1987): Systematics of African bats of the genus *Eptesicus* (Mammalia: Vespertilionidae). 2. Karyotypes of African species and their generic relationships. *Annals Carnegie Museum* 56, 213–222.
- OBRIST, M.; ALDRIDGE, H. D. J. N.; FENTON, M. B. (1989): Roosting and echolocation behavior of the African bat, *Chalinolobus variegatus*. *J. Mammalogy* 70, 828–833.
- RAUTENBACH, I. L. (1985): A new technique for the efficient use of macro-mistnets. *Koedoe* 28, 81–86.

- ROBBINS, L. W.; BAKER, R. J. (1978): Karyotypic data for African mammals, with a description of an in vivo bone marrow technique. *Bull. Carnegie Museum of Natural History* 6, 188–210.
- SMITHERS, R. H. N. (1971): The mammals of Botswana. Museum Memoir No 4. Salisbury: Trustees of the National Museums of Rhodesia.
- (1983): The mammals of the Southern African Subregion. Pretoria: University of Pretoria.
- SMITHERS, R. H. N.; LOBAO TELLO, J. L. P. (1976): Check list and atlas of the mammals of Mocambique. Museum Memoir No 8. Salisbury: Trustees of the National Museums and Monuments of Rhodesia.
- SMITHERS, R. H. N.; WILSON, V. J. (1979): Check list and atlas of the mammals of Zimbabwe Rhodesia. Museum Memoir No 9. Salisbury: Trustees of the National Museums and Monuments.

Authors' addresses: I. L. RAUTENBACH, Transvaal Museum, P.O. Box 413, Pretoria, 0001 South Africa, and M. B. FENTON, Department of Biology, York University, North York, Ontario, Canada, M3J 1P3

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Mammalian Biology \(früher Zeitschrift für Säugetierkunde\)](#)

Jahr/Year: 1992

Band/Volume: [57](#)

Autor(en)/Author(s): Rautenbach I. L., Fenton M. B.

Artikel/Article: [Bats from Mana Pools National Park in Zimbabwe and the first record of *Eptesicus rendallii* from the country 112-115](#)