

WISSENSCHAFTLICHE KURZMITTEILUNG**On the karyotype of the Long-eared hedgehog, *Hemiechinus auritus* (Gmelin, 1770) (Mammalia: Insectivora), in Turkey**

By E. ÇOLAK, N. YİĞİT, M. SÖZEN, and Ş. ÖZKURT

Department of Biology, University of Ankara, Ankara and

Department of Biology, University of Gazi, Kirşehir, Turkey

Receipt of Ms. 16. 12. 1996
Acceptance of Ms. 09. 06. 1997

Hemiechinus auritus is distributed throughout Libya, Egypt, Israel, Lebanon, Syria, Turkey, USSR, Iran, Afghanistan, Pakistan, Mongolia, and China (ELLERMAN and MORRISON-SCOTT 1951; BOBRINSKY et al. 1965; NIETHAMMER 1969; HARRISON 1972; CORBET 1978, 1988; OSBORN and HELMY 1980; SCHOENFELD and YOM-TOV 1985; HARRISON and BATES 1991). The karyotype of this species of hedgehog was described from Daghestan (ORLOV 1969), Egypt (DE HOND 1972), Iraq (BHATNAGAR and EL-AZAWI 1978), Afghanistan (GROPP et al. 1969), India (SOBTI and GILL 1980), but not from other regions. The aim of the present study is to describe karyological characteristics of *H. auritus* in Turkey.

We collected 12 specimens from four localities (Fig. 1) (Aralyk 2, Ceylanpinar 3, Harran 5, Nizip 2) in Turkey and karyotyped four specimens from Ceylanpinar, Harran, and Nizip in southeastern Turkey. Karyotype preparations were made from bone marrow of animals treated with colchicine according to FORD and HAMERTON (1956). 25 metaphase cells of each animal were examined.

H. auritus has $2n = 48$, $NFa = 92$ and $NF = 96$. All the autosomal pairs are bi-armed. The karyotype has a large submetacentric, a large subtelocentric pair, six dot-like chromo-




Fig. 1. Recorded localities (▲) of *H. auritus*. 1. Nizip, 2. Harran, 3. Ceylanpinar, 4. Aralyk




Fig. 2. The karyotype of a male *H. auritus* from Harran.

somes, and 18 pairs of submetacentric and metacentric ones. The X chromosome is a large metacentric and the Y chromosome is the smallest metacentric (Fig. 2).

The karyotype of *H. auritus* also contains $2n = 48$ chromosomes in Iraq (BHATNAGAR and EL-AZAWI 1978), in Daghestan (ORLOV 1969), and in Egypt (GROPP et al. 1969). According to BHATNAGAR and EL-AZAWI (1978), the karyotype of *H. auritus* in Iraq consists of 43 macro- and 5 micro-chromosomes. We found the same karyotype in Turkish specimens, except the large metacentric X chromosome and the metacentric Y chromosome. The karyotype of *H. auritus* from Turkey is similar to that given by ZIMA and KRAL (1984).

In *Echinaceus europaeus*, a karyotype of $2n = 48$ chromosomes has one medium-sized pair of acrocentric (ZIMA and KRAL 1984), while there is no acrocentric chromosome in *H. auritus*. In most species of hedgehogs, the diploid chromosome number was described as being 48 (GEISLER and GROPP 1967; HSU and BENIRSCHKE 1968; GROPP et al. 1969; NATARAJAN and GROPP 1971; GROPP and NATARAJAN 1972). These authors noted large subtelocentric chromosomes in the karyotypes of other hedgehogs, as observed in the karyotype of *H. auritus* in Turkey. According to DOĞRAMACI and GÜNDÜZ (1993), *E. concolor* in Turkey has $2n = 48$, NF_a = 90 and NF = 94, the autosomes contain four pairs of large subtelocentric and one pair of small acrocentric, whereas acrocentric chromosomes are absent in the karyotype described in this study for *H. auritus*.

The sex chromosomes are variable in both *H. auritus* and the other hedgehogs. The X chromosome is metacentric (the smallest one of the macro-chromosomes) (BHATNAGAR and EL-AZAWI 1978) and sub-metacentric (HSU and BENIRSCHKE 1968) for *E. europaeus* and large metacentric (DOĞRAMACI and GÜNDÜZ 1993) for *E. concolor*. We found the X chromosome to be large metacentric for *H. auritus*, which is different from that given by BHATNAGAR and EL-AZAWI (1978), the same in *E. concolor* (DOĞRAMACI and GÜNDÜZ 1993).

The Y chromosome is micro-chromosome and subtelocentric in *H. auritus* and a medium-sized submetacentric in *Paraechinus aethiopicus* (BHATNAGAR and EL-AZAWI 1978), micro-submetacentric in *E. e. europaeus* and micro-metacentric in *E. concolor* (DOĞRAMACI and GÜNDÜZ 1993). In *H. auritus* from Turkey the Y chromosome is micro-metacentric which is different from *H. auritus* from Iraq, *P. aethiopicus* from Iraq but similar to *E. concolor* from Turkey. This showed that the Y chromosome is variable in interpopulation as well as among intrapopulation in the family Erinaceidae.

Literature

- BHATNAGAR, A. N.; EL-AZAWI, T. F. (1978): A karyotype study of chromosomes of two species of hedgehogs, *Hemiechinus auritus* and *Paraechinus aethiopicus* (Insectivora: Mammalia). *Cytologia* **43**, 53–59.
- BOBRINSKY, N. A.; KUZNETSOV, B. A.; KUZYAKIN, A. P. (1965): Key to the Mammals of the USSR, 2nd ed. Moscow: Izdatel stvo Prosveshchenie, (in Russian).
- CORBET, G. B. (1978): The mammals of the Palaearctic Region: a taxonomic review. London and Ithaca: British Museum (Nat. Hist.). Cornel Univ. Press.
- CORBET, G. B. (1988): The family Erinaceidae: a synthesis of its taxonomy, phylogeny, ecology, and zoogeography. *Mammal Rev.* **18**, 117–172.
- DOĞRAMACI, S.; GÜNDÜZ, I. (1993): Türkiye *Erinaceus concolor* (Mammalia: Insectivora) Türünün Taksonomisi ve Yayılışı. *Doğa-Tr. J. Zoology* **17**, 267–288.
- ELLERMAN, J. R.; MORRISON-SCOOT, T. C. S. (1951): Checklist of Palearctic and Indian Mammals 1758 to 1946. London: Brit. Mus. Nat. Hist.
- FORD, C. E.; HAMERTON, J. L. (1956): Colchicine-hypotonic citrate squash sequence for mammalian chromosomes. *Stain Tech.* **31**, 247–251.
- GEISLER, M.; GROPP, A. (1967): Chromosome polymorphism in the European hedgehog, *Erinaceus europaeus* (Insectivora). *Nature* **214**, 396.
- GROPP, A.; NATARAJAN, A. T. (1972): Karyotype and heterochromatin pattern of the Algerian hedgehog. *Cytogenetics* **11**, 259–269.
- GROPP, M.; CITOLER, P.; GEISLER, M. (1969): Karyotype variation and heterochromatin pattern in hedgehogs (*Erinaceus* and *Hemiechinus*). *Chromosome* **27**, 228–307.
- HARRISON, D. L. (1972): The Mammals of Arabia: Lagomorpha and Rodentia. Vol. 3. London: Ernest Benn. Pp. 385–670.
- HARRISON, D. L.; BATES, P. J. J. (1991): The Mammals of Arabia. Sec. ed. Kent: Harr. Zool. Museum. Pub.
- HOND'T, H. A. DE (1972): Karyological studies on two insectivores of Egypt. *Proc. Egypt. Acad. Sci.* **25**, 171–174.
- HSU, T. C.; BENIRSCHKE, K. (1967): An atlas of mammalian chromosomes. Vol. 1. New York: Springer Verlag.
- NATARAJAN, A. T.; GROPP, A. (1971): The meiotic behaviour of autosomal heterochromatin segments in hedgehogs. *Chromosoma* **35**, 143–152.
- NIETHAMMER, J. (1969): Zur Taxonomie der Ohrenigel in Afghanistan (Gattung *Hemiechinus*). *Z. Säugetierkunde* **34**, 257–274.
- ORLOV, V. N. (1969): Chromosome complements of the hedgehogs from East Europe. Mat. II. vsesoj. sov. pomlekopit. N. N. Voroncov (ed.): 6–7 (in Russian).
- OSBORN, D. J.; HELMY, I. (1980): The contemporary land mammals of Egypt (including Sinai). *Fieldiana Zool.* **5**, 1–579.
- SCHOENFELD, M.; YOM-TOV, V. (1985): The biology of two species of hedgehogs, *Erinaceus europaeus concolor* and *Hemiechinus auritus aegyptius*, in Israel. *Mammalia* **49**, 339–355.
- SOBTI, R. C.; GILL, S. S. (1980): Chromosome number in fifteen mammals from India. *Mamm. Chrom. Newslett.* **21**, 107.
- ZIMA, J.; KRAL, B. (1984): Karyotypes of European mammals I. *Acta. Sc. Nat. Brno* **8**, 15–17.

Authors addresses: ERCÜMENT ÇOLAK, NURI YİĞİT, and MUSTAFA SÖZEN, Department of Biology, University of Ankara, 06100 Beşevler/Ankara, Turkey; ŞAKIR ÖZKURT, Department of Biology, University of Gazi, Kırşehir, Turkey.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Mammalian Biology \(früher Zeitschrift für Säugetierkunde\)](#)

Jahr/Year: 1997

Band/Volume: [62](#)

Autor(en)/Author(s): Colak Ercümert, Yigit Nuri, Sözen Mustafa, Özkurt Sakir

Artikel/Article: [On the karyotype of the Long-eared hedgehog, *Hemiechinus auritus* \(Gmelin, 1770\) \(Mammalia: Insectivora\), in Turkey 372-374](#)