



Bayerische  
Staatsammlung  
für Paläontologie und Geologie

- München, 01.07.2017
- Manuscript received 30.12.2016; revision accepted 21.01.2017
- ISSN 0373-9627
- ISBN 978-3-946705-00-0

### *Short Communication*

## First report of sea cucumbers (Echinodermata: Holothuroidea) from the latest Cretaceous of Bavaria, Germany

Mike Reich<sup>1,2,3</sup>

<sup>1</sup>SNSB - Bayerische Staatsammlung für Paläontologie und Geologie, Richard-Wagner-Straße 10, 80333 Munich, Germany

<sup>2</sup>Ludwig-Maximilians-Universität München, Department für Geo- und Umweltwissenschaften, Paläontologie und Geobiologie, Richard-Wagner-Straße 10, 80333 Munich, Germany

<sup>3</sup>GeoBio-Center der Ludwig-Maximilians-Universität München, Richard-Wagner-Straße 10, 80333 Munich, Germany

E-mail: m.reich@lrz.uni-muenchen.de

Zitteliana 89, 285–289.

**Key words:** fossil Holothuroidea; Cretaceous; Maastrichtian; Bavaria; Germany

**Schüsselwörter:** fossile Holothuroidea; Kreide; Maastrichtium; Bayern; Deutschland

The Bavarian Gerhardtsreit Formation ("Gerhardtsreiter Mergel" / "Gerhardtsreiter Schichten"; cf. Böhm 1891; Hagn 1960; Wagreich et al. 2004), also known as Gerhartsreit Formation ("Gerhartsreiter Schichten"; Hagn et al. 1981, 1992; Schwarzhans 2010; Pollerspöck & Beaury 2014) or "Gerhardsreuter Schichten" (Egger 1899; Hagn & Hözl 1952; de Klasz 1956; Herm 1979, 2000) is exposed in Upper Bavaria near Siegsdorf, southern Germany (cf. Hagn et al. 1981). The black to grey-black, fine sandy micaceous clayish marls were deposited on the northern Helvetic shelf (northwestern Tethyan realm; Herm 1979, 2000; Hagn et al. 1992), and are Maastrichtian in age.

The formation is well known for its diverse and well preserved assemblages of microfossils (e.g., Egger 1899; Ziegler 1959; Ohmert 1970; Hagn et al. 1981, 1992; Kirsch 1991), invertebrates, especially molluscs (Böhm 1891; de Klasz 1956; Hagn et al. 1992), and vertebrates (chondrichthyan and osteichthyan fishes; e.g., Schwarzhans 1996, 2010; Pollerspöck & Beaury 2014; Pollerspöck 2015). Collections have been made from the strata for over 150 years (Gümbel 1861: "Gosauschichten von Siegsdorf", Gümbel 1870: "Gosaumergel von Götzreuth").

While examining micropalaeontological sieve residues of this formation, besides other microscopic echinoderm remains (Echinoidea, Ophiuroidea) a single holothurian calcareous ring element was found.

Whereas modern holothurians are common and ecologically important members inhabiting all marine environments, from the intertidal to the deepest oceanic trenches worldwide (e.g., Pawson 1982; Kerr & Kim 2001; Kerr 2003), fossil holothurians remain relatively rare (e.g., Frizzell & Exline 1956, 1966; Gilliland

1993; Smith 2004) due to different reasons (Reich 2013). There are nearly 1,700 valid extant sea cucumber species (Smiley 1994; Kerr 2003; Paulay pers. comm.) known worldwide. The fossil record (since the Middle Ordovician; Reich 1999, 2010), by contrast, is discontinuous in time and recorded ranges of species with around 1,000 reported forms (Reich 2013, 2014, 2015b) since the early 19th century.

Although well-preserved articulated material of holothurians are known from around a dozen fossil Lagerstätten localities (Reich 2013) most of the fossil record of the group comprises isolated body-wall ossicles and elements of the calcareous ring (Pawson 1980; Smith et al. 2013; Reich 2015a).

The Cretaceous record of Holothuroidea was quite poor and understudied (Frizzell & Exline 1956; Müller 1969; Reich 1995, 1997) in comparison to Triassic and Jurassic holothurian faunas (Reich 2013). However, first gaps were filled by several papers published in the last 15 years (e.g., Al-Tamimi et al. 2001; Reich 2001a, 2001b, 2002a, 2002b, 2003a, 2003b, 2003c, 2003d; Reich & Wiese 2010; Reich & Ansorge 2014; Sadreddin & Al-Tamimi 2006; Al-Tamimi & Saqqa 2008).

The only known Cretaceous holothurian from Bavaria is represented by an isolated body-wall wheel ossicle ("*Calcarina rotula*", originally misinterpreted as a benthic foraminifer) from the Cenomanian of the "Einbachgraben" (Bavarian Alps), described by Joseph Georg Egger (1824–1913) in 1899 (cf. Reich 1997) which is in fact a member of the Apodida: Chiridotidae ("*Jumaraina rotula*"). Thus, unequivocal, identifiable holothurian fossils are important because this echinoderm group has largely been neglected by palaeontologists.

Images of the specimen figured here were captured with a Keyence VHX 5000 digital microscope with live depth composition (50–100 single stacked images). Figures were processed using Adobe Photoshop®.

Abbreviations: SNSB-BSPG – Bavarian State Collection of Palaeontology and Geology, Munich, Germany; CR – calcareous ring, R – radial element(s) of the calcareous ring, IR – interradial element(s) of the calcareous ring, RI to RV – Ludwig (1889) letters for designation of dorsal/ventral position of R: RI = mid-ventral, RII = left-ventral, RIII = right-ventral, RIV = left-dorsal, RV = right-dorsal.

## Systematic Palaeontology

The taxonomy, descriptive terminology and systematic scheme employed in this paper was adapted and modified from or is conform to those of Heding (1933), Gage & Billett (1986), Smirnov (2012), and Reich (2015a).

Phylum Echinodermata Bruguière, 1791

[ex Klein, 1734]

Subphylum Eleutherozoa Bell, 1891

Class Holothuroidea de Blainville, 1834

Subclass Holothuriacea Smirnov, 2012

Order Molpadida Haeckel, 1896

?(stem group) fam. nov.

gen. et sp. nov. A

Fig. 1A-F

Type material: SNSB-BSPG 2016 VIII 151, radial CR element (?RI); isolated from the sieve residue of micropalaeontological sample #3857 (collected by Herbert Hagn in 1977).

Type locality: Gerhardtsreiter (Gerhartsreiter) Graben, SE of Siegsdorf, Upper Bavaria, Germany.

Type stratum and age: Gerhardtsreit (Gerhartsreit) Formation; Upper Cretaceous: Maastrichtian. – The precise biostratigraphy of this formation was the subject of debate during the last decades: (1) Lower Maastrichtian (e.g., de Klasz 1956; Ziegler 1959; Hagn 1961; Herm 1979), (2) Middle Maastrichtian (e.g., Hagn et al. 1981; Martini 1981; Hagn et al. 1992), or (3) upper Lower Maastrichtian to lower Upper Maastrichtian (Ohmert 1967), as well as (4) Upper Maastrichtian (e.g., Ohmert 1970; Herm 2000). However, the reported nannofossils (e.g., *Lithraphidites quadratus*; cf. Martini 1981; UC20 zone of Burnett 1998) and several foraminifera (e.g., *Bolivinoides draco giganteus*, *Bolivinoides draco draco*; cf. Koch 1977, Frenzel 2000) indicate an upper Lower Maastrichtian age (probably the fastigata belemnite zone).

Palaeoenvironment: Based on teleost fish oto-

liths described from the type stratum, Schwarzhans (2010) interpreted the environment as open or deeper shelf. This is in accordance to observations on the palaeoecology (cf. Frenzel 2000) of the recorded benthic foraminiferal fauna.


Description: The present specimen of an radial CR element is elongate and nearly subsquarish in outline; the whole element tends to be slightly curved with the inner face being concave and both lateral articulation faces inclined slightly inwards or being chamfered.

The muscular process bearing the notch for the passage of the radial nerve (Fig. 1B) is short, and facing somewhat outward (Fig. 1C, E) in part. The additional anterior process (non-muscular process; Fig. 1A-B) is slightly longer. On the external face (Fig. 1A) a medium-sized depression lies between the two anterior processes. On the inner face a posterior indent (encompassed by two short posterior extensions; Fig. 1A-F) extends into a furrow which divides anteriorly into three (?four) branches (Fig. 1B, C, E). Stereom structure is very dense.

Dimensions: Length of R: 1.19 mm; maximum width: 0.66 mm.

Discussion and remarks: Since only one specimen (of the diagnostic CR) of this taxon has been so far recorded I designated it as "gen. et sp. nov. A". The present R resembles the CR of molpadiid/caudinid holothurians but, by contrast, the typical posterior prolongations are short ("rudimentary") like in a few "non-typical" molpadids (e.g., *Eupyrgus*, *Cherbonniera*; Reich & Rössner 2016). However, by comparison with modern molpadid holothurian material (e.g., Hôzawa 1928; Heding 1933; Hatanaka 1939) it is possible to draw conclusions about some of the soft tissue components that were related to the CR (cf. Fig. 1A-F). Due to the size relationship of both anterior processes (shorter muscular process!) the position of the present CR element was probably mid-ventral (= RI). The central groove would have housed the anterior extension of the radial canal, leading to several tentacular canals (cf. also Hatanaka 1939). The insertion area of retractor/longitudinal muscles is located below the passage for the radial nerve (muscular process) (Fig. 1A-B). The new taxon is clearly different from modern molpadiid, caudinid and eupyridid sea cucumbers that have a differing morphology and combination of anterior and posterior processes. It probably represents a stem group member (of our modern molpadids) rather than a representative of extant families.

Molpadid sea cucumbers are medium in size with a body which is often sausage-shaped, tapering posteriorly to form a conspicuous tail. These forms are deposit-feeders, usually burrowing into soft sediments or sand, often orientated with the anterior end downwards in the substrate and with the posterior end near the substrate surface for respiration (e.g.,



**Figure 1:** Radial CR element (probably RI) of an molpadid holothurian (gen. et sp. nov. A) from the Early Maastrichtian Gerhardtsreit Formation of Bavaria (Gerhardtsreiter Graben, near Siegsdorf), Germany (SNSB-BSPG 2016 VIII 151). **(A)** outer view; **(B)** inner view; **(C)** lateral (oblique) view (of left face); **(D)** lateral view (of right face); **(E)** anterior view; **(F)** posterior view. The left figure shows the overall morphology viewed from all directions, and the right figure with same configuration indicates interpretation of soft tissue anatomy related to the CR.

Pawson 1977, 1982). The group contains around 100 modern species in 9 genera and three families. Members of the Molpadida live in shallow or deep water with a bathymetric range of 0–5,250 m (e.g., Pawson 1977; Pawson & Liao 1992; Pawson et al. 2001; Pawson & Vance 2007; Davey & O’Loughlin 2013; Reich & Rössner 2016).

Late Cretaceous molpadid CR material was unknown so far. Hess (1975) and Gilliland (1992) reported on a few examples from the older Swiss and British Jurassic. Molpadid body-wall ossicles from Upper Cretaceous strata were recently described by Reich (2003d) and Reich & Ansorge (2014). The fossil record of Molpadida is poorly known, therefore each additional find can add significantly to our knowledge of extinct Holothuroidea or can help in identifying calibration points for phylogenies.

**Occurrence:** So far known only from the Lower Maastrichtian of the Bavarian Alps, Germany.

The newly discovered fossil constitutes the first reported occurrence of Holothuroidea from the Maastrichtian of Bavaria as well as of molpadid CR material from Late Cretaceous strata worldwide.

## Acknowledgements

This short article is dedicated to Winfried Werner, Munich, on the occasion of his 65<sup>th</sup> birthday. Funding for this research was supported in part by the German Research Foundation (Project: RE2599/6-2). Thanks are also given to T.R. Stegemann (Göttingen/München) for critically reading the first draft of the manuscript. I also received useful comments on this short paper by reviews of Andreas Kroh and Alexander Nützel.

## References

- Al-Tamimi M, Saqqa WA. 2008. Holothurian Sclerites from the Upper Part of Ajlun Group (Late Cretaceous) of Salhoub Area, North Jordan. *Dirasat, Pure Sciences* 35(2), 81–95.
- Al-Tamimi M, Saqqa W, Sadeddin W. 2001. Holothurian sclerites from the Early Late Cretaceous, south of Al-Mastabah Area (North Jordan). *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* [2001](8), 463–482.
- Bell FJ. 1891. On the Arrangement and Inter-relations of the Classes of Echinodermata. *Annals and Magazine of Natural History* (6th series) 8, 206–215.
- Blainville HMD de. 1834. *Manuel d’Actinologie et de Zoophytologie*. 2 vols. Paris & Strasbourg, Levrault, viii + 694 p.
- Böhm J. 1891. Die Kreidebildungen des Fürbergs und Sulzbergs

- bei Siegsdorf in Oberbayern. *Palaeontographica* 38(1-2), 1–106.
- Bruguière JG. 1791. Encyclopédie méthodique, ou par ordre de matières: par une Société de Gens de Lettres, de Savans et d'Artistes. Précédée d'un Vocabulaire universel, servant de Table pour tout l'Ouvrage, ornée des Portraits de MM. Diderot & d'Alembert, premiers éditeurs de l'Encyclopédie. Tableau encyclopédique et méthodique des trois règnes de la nature. [5] L'helminthologie, ou les ver infusoires, les vers intestins, les vers mollusques, etc. - Vers Echinodermes. Paris, Panckoucke, viii + 132 p.
- Burnett JA. 1998. Upper Cretaceous. In: PR Bown (Ed), *Calcareous Nannofossil Biostratigraphy*. London, Chapman & Hall, 132–199.
- Davey N, O'Loughlin M. 2013. The caudinid sea cucumbers of New Zealand (Echinodermata: Holothuroidea: Molpadida: Caudinidae). *Zootaxa* 3613(4), 357–368.
- Egger JG. 1899. Foraminiferen und Ostrakoden aus den Kreidemergeln der Oberbayerischen Alpen. Abhandlungen der Königlich bayerischen Akademie der Wissenschaften (II. Classe) 21(I), 1–230.
- Frenzel P. 2000. Die benthischen Foraminiferen der Rügener Schreibkreide (Unter-Maastricht, NE-Deutschland). Neue Paläontologische Abhandlungen 3, 1–361.
- Frizzell DL, Exline H. 1956. Monograph of Fossil Holothurian Sclerites. *Bulletin of School of Mines and Metallurgy (Technical Series)* 89[1955](1), 1–204.
- Frizzell DL, Exline H. 1966. Holothuroidea - Fossil Record. In: RC Moore (Ed), *Treatise on Invertebrate Paleontology*, U, Echinodermata 3 [Asterozoa-Echinozoa]. Lawrence, Kan., University of Kansas Press & Boulder, Colo., Geological Society of America, U646–U672.
- Gage JD, Billett DSM. 1986. The family Myriotrichidae Théel (Echinodermata: Holothuroidea) in the deep northeast Atlantic Ocean. *Zoological Journal of the Linnean Society* 88(3), 229–276.
- Gilliland PM. 1992. Holothurians in the Blue Lias of southern Britain. *Palaeontology* 35(1), 159–210.
- Gilliland PM. 1993. The skeletal morphology, systematics and evolutionary history of holothurians. *Special Papers in Palaeontology* 47, 1–147.
- Gümbel CW. 1861. Geognostische Beschreibung des bayerischen Alpengebirges und seines Vorlandes. Gotha, J. Perthes, xx + 950 p.
- Gümbel CW. 1870. Vergleichung der Foraminiferenfauna aus den Gosaumergeln und den Belemniten-Schichten der bayrischen Alpen. *Sitzungsberichte der Königlich bayerischen Akademie der Wissenschaften zu München (Mathematisch-physikalische Classe)* [1870](II), 278–288.
- Haeckel E. 1896. Systematische Phylogenie der Echinodermen. In: E Haeckel, Entwurf eines Natürlichen Systems der Organismen auf Grund ihrer Stammesgeschichte. Zweiter Theil: Systematische Phylogenie der Wirbellosen Thiere (Invertebrata). Berlin, G. Reimer, 348–504.
- Hagn H. 1960. Die stratigraphischen, paläogeographischen und tektonischen Beziehungen zwischen Molasse und Helvetikum im östlichen Oberbayern. *Geologica Bavaria* 44, 1–208.
- Hagn H. 1961. Klassische und neue Aufschlüsse mit Faunen der Oberkreide und des Tertiärs in den östlichen Bayerischen Alpen und angrenzenden Gebieten. *Paläontologische Zeitschrift* 35(3/4), 146–170.
- Hagn H, Darga R, Schmid R. 1992. Erdgeschichte und Urwelt im Raum Siegsdorf. Fossilien als Zeugen der geologischen Vergangenheit. Siegsdorf, Eigenverlag Gemeinde Siegsdorf, 241 p.
- Hagn H, Herm D, Martini E. 1981. F 2: Gerhartsreiter Graben. In: H Hagn (Ed), *Die Bayerischen Alpen und ihr Vorland in mikropaläontologischer Sicht*. Exkursionsführer zum 17. Europäischen Mikropaläontologischen Kolloquium in Oberbayern, September 1981. *Geologica Bavaria* 82, 239–242.
- Hagn H, Hözl O. 1952. Geologisch-paläontologische Untersuchungen in der subalpinen Molasse des östlichen Oberbayerns zwischen Prien und Sur mit Berücksichtigung des im Süden anschließenden Helvetikums. *Geologica Bavaria* 10, 1–208.
- Hatanaka M. 1939. A study of the caudate holothurian, *Molpadia roretzii* (v. Marenzeller). *Science Reports of the Tōhoku Imperial University (Fourth Series: Biology)* 14(2/3), 155–190.
- Heding SG. 1933. The *Caudina* of Asamushi, the so-called *Caudina chilensis* (Johs. Müller). *Science Reports of the Tōhoku Imperial University (Fourth Series: Biology)* 8(2), 127–142.
- Herm D. 1979. Die süddeutsche Kreide – Ein Überblick. In: J Wiedmann (Ed), *Aspekte der Kreide Europas*. IUGS Series A 6, 85–106.
- Herm D. 2000. Kreide in den deutschen Alpen. In: *Stratigraphische Kommission Deutschlands (Ed), Stratigraphie von Deutschland III: Die Kreide der Bundesrepublik Deutschland*. Courier Forschungsinstitut Senckenberg 226, 147–162.
- Hess H. 1975. Die fossilen Echinodermen des Schweizer Juras. *Veröffentlichungen aus dem Naturhistorischen Museum Basel* 8, 1–130.
- Hōzawa S. 1928. On the Changes occurring with Advancing Age in the Calcareous Deposits of *Caudina chilensis* (J. Müller). *Science Reports of the Tōhoku Imperial University (Fourth Series: Biology)* 3(3), 362–378.
- Kerr AM. 2003. Holothuroidea (Sea Cucumbers). In: D Thoney (Ed), *Grzimek's Animal Life Encyclopedia*. Second Edition. Volume I. Lower Metazoans and Lesser Deuterostomes. New York, N.Y., Gale Group Publishers, 417–431.
- Kerr AM, Kim J. 2001. Phylogeny of Holothuroidea (Echinodermata) inferred from morphology. *Zoological Journal of the Linnean Society* 133(1), 63–81.
- Kirsch K-H. 1991. Dinoflagellatenzysten aus der Oberkreide des Helvetikums und Nordultrahelvetikums von Oberbayern. *Münchener Geowissenschaftliche Abhandlungen (A: Geologie und Paläontologie)* 22, 1–306.
- Klasz I de. 1956. Stratigraphie der helvetischen Zone (Helvetikum-Zone). In: O Ganss, I de Klasz, K Götzinger, F Vogel, *Geologie des Blattes Bergen*. *Geologica Bavaria* 26, 42–72.
- Klein JT. 1734. *Naturalis Dispositio Echinodermatum: Accessit Lucubratiuncula De Aculeis Echinorum Marinorum, Cum Spicilegio De Belemnitis*. Gedani [= Gdansk], Schreiber, 78 + 36 p.
- Koch W. 1977. Biostratigraphie in der Oberkreide und Taxonomie von Foraminiferen. *Geologisches Jahrbuch (A: Allgemeine und regionale Geologie BR Deutschland und Nachbargebiete, Tektonik, Stratigraphie, Paläontologie)* 38, 11–123.
- Ludwig H. 1889–1892. Die Seewalzen. In: HG Bronn (Ed), *Klassen und Ordnungen des Thier-Reichs, wissenschaftlich dargestellt in Wort und Bild*. Zweiter Band. Abtheilung: Echinodermen (Stachelhäuter). I Buch. Leipzig, C.F. Winter'sche Verlagshandlung, vi + 460 p. [1889: 1–176, 1890: 177–240, 1891: 241–376, 1892: 377–460].
- Martini E. 1981. Nannoplankton in der Ober-Kreide, im Alttertiär und im tieferen Jungtertiär von Süddeutschland und dem angrenzenden Österreich. In: H Hagn (Ed), *Die Bayerischen Alpen und ihr Vorland in mikropaläontologischer Sicht*. Exkursionsführer zum 17. Europäischen Mikropaläontologischen Kolloquium in Oberbayern, September 1981. *Geologica Bavaria* 82, 345–356.
- Müller AH. 1969. Reste seltener Holothurien (Echinodermata) aus dem Mesozoikum Europas. *Monatsberichte der Deutschen Akademie der Wissenschaften zu Berlin* 11(8/9), 662–671.
- Ohmert W. 1967. Die Ostracoden-Gattung *Cythereis* aus der Oberkreide Südbayerns. *Inaugural-Dissertation zur Erlangung der Doktorwürde der Hohen Naturwissenschaftlichen Fakultät der Ludwig-Maximilians-Universität zu München*. München/Bamberg, Eigenverlag (Bamberger Fotodruck), viii + 1–192 + vii p. Anh.
- Ohmert W. 1970. Die Ostracodengattung *Golocythere* aus der Oberkreide Südbayerns. *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und historische Geologie* 10, 205–228.
- Pawson DL. 1977. Molpadiid Sea Cucumbers (Echinodermata: Holothuroidea) of the Southern Atlantic, Pacific, and Indian Ocean. In: DL Pawson (Ed), *Biology of the Antarctic Seas VI*. Antarctic Research Series 26, 97–123.
- Pawson DL. 1980. Holothuroidea. In: TW Broadhead, JA Waters (Eds), *Echinoderms. Notes for a Short Course*. University of Tennessee, Department of Geological Sciences, Studies in

- Geology 3, 175–189, [references: 215–235].
- Pawson DL. 1982. Holothuroidea. In: SP Parker (Ed), Synopsis and Classification of Living Organisms, Volume 2. New York, N.Y., McGraw-Hill, 813–818.
- Pawson DL, Liao Y. 1992. Molpadiid sea cucumbers of China, with descriptions of five new species (Echinodermata: Holothuroidea). Proceedings of the Biological Society of Washington 105(2), 373–388.
- Pawson DL, Vance DJ. 2007. *Heteromolpadija joyceae*, a new species of sea cucumber from New Zealand, with a key to New Zealand Molpadiidae (Echinodermata: Holothuroidea: Molpadiida). Zootaxa 1608, 35–39.
- Pawson DL, Vance DJ, Ahearn C. 2001. Western Atlantic sea cucumbers of the Order Molpadiida (Echinodermata: Holothuroidea). Bulletin of the Biological Society of Washington 10, 311–327.
- Pollerspöck J. 2015. Ammenhaie – Opfer des Kreide–Tertiär Massensterbens. Fossilien 32(2), 35–39.
- Pollerspöck J, Beaury B. 2014. *Parasquatina zitteli* nov. sp. (Elastomorpha: Orectolobiformes) aus dem Maastricht von Oberbayern (Gerhartsreiter Schichten, Siegsdorf) und Bemerkungen zur Verbreitung der Ordnung Orectolobiformes. Zitteliana A 54, 147–164.
- Reich M. 1995. Erster sicherer Nachweis der Elasipoda (Holothuroidea, Echinodermata) aus der Kreide, sowie Bemerkungen zu den Holothurienresten der Oberkreide. Archiv für Geschiebekunde 1(11), 681–688.
- Reich M. 1997. Holothurienreste (Echinodermata) aus der Oberkreide – ein Überblick. Zeitschrift der Deutschen Geologischen Gesellschaft 148(1), 71–79.
- Reich M. 1999. Ordovizische und silurische Holothurien (Echinodermata). Greifswalder Geowissenschaftliche Beiträge 6, 479–488.
- Reich M. 2001a. *Pravuscucumis deeckeii* paragen. et parasp. nov. (Echinodermata: Holothuroidea) aus dem Turonium des Ostseegebietes. Geschiebekunde aktuell 17(2/3), 97–106.
- Reich M. 2001b. Holothurians from the Late Cretaceous of the Isle of Rügen (Baltic Sea). In: M Barker (Ed), Echinoderms 2000. Lisse etc., A. A. Balkema Publishers, 89–92.
- Reich M. 2002a. Die Holothurien (Echinodermata) der Oberkreide. Unpublished Inauguraldissertation, Leopold-Franzens University Innsbruck. Innsbruck, 584 p.
- Reich M. 2002b. Holothurien (Echinodermata) aus der Oberkreide des Ostseeraumes: Teil 1. Myriotrichidae Théel, 1877. Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen 224(3), 373–409.
- Reich M. 2003a. Holothurien (Echinodermata) aus der Oberkreide des Ostseeraumes: Teil 2. *Ypsilothuriidae* Heding, 1942. Neues Jahrbuch für Geologie und Paläontologie, Monatshefte [2003] (8), 498–512.
- Reich M. 2003b. Holothurien (Echinodermata) aus der Oberkreide des Ostseeraumes: Teil 3. Chiridotidae Östergren, 1898. Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen 228(3), 363–397.
- Reich M. 2003c. Holothurien (Echinodermata) aus der Oberkreide des Ostseeraumes: Teil 4. Synaptidae Burmeister, 1837. Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen 229(1), 75–95.
- Reich M. 2003d. Holothurien (Echinodermata) aus der Oberkreide des Ostseeraumes: Teil 5. Molpadiidae J. Müller, 1850. Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen 229(2), 231–253.
- Reich M. 2010. The early evolution and diversification of holothurians (Echinozoa). In: LG Harris, SA Böttger, CW Walker, MP Lesser (Eds), Echinoderms: Durham. Boca Raton/London etc., CRC Press, Taylor & Francis Group, 55–59.
- Reich M. 2013. How many species of fossil holothurians are there?. In: C Johnson (Ed), Echinoderms in a Changing World. Boca Raton/London etc., CRC Press, Taylor & Francis Group, 23–51.
- Reich M. 2014. Supplement to: „How many species of fossil holothurians are there?“. Göttingen Contributions to Geosciences 77, 161–162.
- Reich M. 2015a. Different pathways in early evolution of the holothurian calcareous ring? In: S Zamora, I Rábano (Eds), Progress in Echinoderm Palaeobiology. Cuadernos del Museo Geominero 19, 137–145.
- Reich M. 2015b. Supplement 2 to: “How many species of fossil holothurians are there?” Zitteliana A 55, 121–122.
- Reich M, Ansorge J. 2014. Santonian sea cucumbers (Echinodermata: Holothuroidea) from Sierra del Montsec, Spain. Göttingen Contributions to Geosciences 77, 147–160.
- Reich M, Rössner GE. 2016. Unusual calcareous ring morphology in molpadiid sea cucumbers (Echinodermata)? Zitteliana 88, p. 41.
- Reich M, Wiese F. 2010. Apodid sea cucumbers (Echinodermata: Holothuroidea) from the Upper Turonian of the Isle of Wolin, NW Poland. Cretaceous Research 31(4), 350–363.
- Sadeddin W, Al-Tamimi M. 2006. Two New Holothurian Sclerite Genera from the Late Cretaceous of Jordan. Journal of Applied Sciences 6(11), 2426–2431.
- Schwarzans W. 1996. Otoliths from the Maastrichtian of Bavaria and their evolutionary significance. In: G Arratia, G Viohl (Eds), Mesozoic Fishes – Systematics and Paleoecology. München, Verlag Dr. Friedrich Pfeil, 417–431.
- Schwarzans W. 2010. Otolithen aus den Gerhartsreiter Schichten (Oberkreide: Maastricht) des Gerhartsreiter Grabens (Oberbayern). Palaeo Ichthyologica 4, 1–100.
- Smiley S. 1994. Holothuroidea. In: FW Harrison, F-S Chia (Eds), Microscopic Anatomy of Invertebrates 14[Echinodermata], 401–471.
- Smirnov AV. 2012. System of the Class Holothuroidea. Paleontological Journal 46(8), 793–832.
- Smith AB. 2004. Echinoderms (other than Echinoids). In: R Seeley, R Cocks (Eds), Encyclopaedia of Geology. Oxford, Elsevier, 334–341.
- Smith AB, Reich M, Zamora S. 2013. Comment on supposed holothurian body fossils from the middle Ordovician of Wales (Botting and Muir, *Palaeontologia Electronica*: 15.1.9A). Palaeontologia Electronica 16.1.2A, 1–6.
- Wagreich M, Egger H, Schnabel W, Summesberger H, Lukeiner A. 2004. Kreide. In: WE Piller, CW Erhart (Coord), Die stratigraphische Tabelle von Österreich 2004 (sedimentäre Schichtfolgen). Wien, Kommission für die paläontologische und stratigraphische Erforschung Österreichs der Österreichischen Akademie der Wissenschaften und Österreichische Stratigraphische Kommission.
- Ziegler JH. 1959. *Lituola grandis* (Reuss) aus dem Untermaastricht der Bayerischen Alpen. Ein Beitrag zur Systematik und Stammbaugeschichte der Lituolidae (Foram.). Palaeontographica (A: Paläozoologie - Stratigraphie) 112(1/4), 59–132.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Zitteliana Serie A+B gemeinsam](#)

Jahr/Year: 2017

Band/Volume: [89](#)

Autor(en)/Author(s): Reich Mike

Artikel/Article: [First report of sea cucumbers \(Echinodermata: Holothuroidea\) from the latest Cretaceous of Bavaria, Germany 285-289](#)