

#### für Paläontologie und Geologie

München, 2019

- Manuscript received 10.10.2017; revision accepted 12.02.2018; available online: 13.12.2018
- ISSN 0373-9627
- ISBN 978-3-946705-05-5

## On the occurrence of the Indonesian ammonite *Macrocephalites keeuwensis* Boehm [M & m] from Kachchh (Western India)

Sreepat Jain\*

Department of Applied Geology, School of Applied Natural Sciences, Adama Science and Technology University, P.O. Box 1888, Adama, Ethiopia

\*E-mail: sreepatjain@gmail.com

Zitteliana 93, 3-24.

#### Abstract

The dimorphic occurrence of the Indonesian ammonite *Macrocephalites keeuwensis* Boehm is reported co-occurring with the Late Bathonian *Macrocephalites* cf. *mantataranus* Boehm [M] from the basal sediments (ash gray marl-limestone intercalations) of the Jara Dome (Kachchh, Western India). Coeval sediments from the adjoining Jumara Dome also record *M. cf. mantataranus* Boehm [M] and *M. keeuwensis* var. aff. forma *flexuosa* Boehm [m]. Based on these finds, the age of the Indonesian *M. keeuwensis* Association is re-evaluated as also the age of the nominal species that now occurs in Kachchh, Madagascar and Europe. A new Late Bathonian age is proposed for the Association that hitherto was assigned on balance to straddle between "late Early Callovian–latest Bathonian". It is speculated that *M. keeuwensis* Boehm [M & m] gave rise to two distinct lineages and the paper documents the separation of these two different lines. One that gave rise to the Early Callovian *M. lamellosus* (Sowerby), and which in turn to the late Early Callovian *Subkossmatia opis* Spath. On the other hand, the dimorphic pair also gave rise to the early–middle Early Callovian *Macrocephalites formosus* (Waagen), which in turn gave rise to the late Early Callovian *M. semilaevis* Spath. *M. keeuwensis* [M & m], in turn, arose from the late Middle Bathonian *M. bifurcatus* transient *intermedius* Boehm (from one of its varieties, A or B). All forms occur in Kachchh. *M. lamellosus* (Sowerby) [M & m] and *M. formosus* [M & m] are now reported from Kachchh.

Key words: Macrocephalites, Bathonian, Indonesia, Kachchh, Western India

#### 1. Introduction

The presence of an Middle Bathonian assemblage of Indonesian macrocephalitids in Kachchh (*Macrocephalites bifurcatus* transient *intermedius* [M], *M. bifurcatus* transient aff. *bifurcatus* [M] and *M. cf. etheridgei* Spath [m] have been noted from the Jumara Dome, Kachchh (Western India; Fig. 1) (Jain 2002, 2014; see also Roy et al. 2007). However, similar records of Indonesian ammonites from higher up in the Late Bathonian, are rare (Jain & Desai 2014).

The present contribution reports the occurrence of the dimorphic Indonesian ammonite *Macrocephalites keeuwensis* Boehm [M & m] from the ash gray marl-limestone intercalations from the core of Jara Dome (Figs 2, 3) from where recently, the Late Bathonian *M.* cf. *mantataranus* Boehm [M] was also recorded (Jain & Desai 2014). From coeval sediments from the adjoining Jumara Dome (Fig. 1), the presence of *M.* cf. *mantataranus* Boehm [M] (Jain & Desai 2014) and now *M. keeuwensis* var. aff. forma



**Fig. 1:** The Kachchh basin is characterized by residual basement ridges, now exposed as domal outcrops or uplifts viz. the Kachchh Mainland, Pachchham, Khadir, Bela, Chorad and Wagad. These constitute islands in an otherwise flat lying surface with the latter five being collectively referred as "Islands". Two domal outcrops within the Kachchh Mainland are studied for this contribution – Jara and Jumara.

*flexuosa* Boehm [m] are also recorded (Fig. 3). Based on these finds, the age of the Indonesian *M. keeuwensis* Association is re-evaluated as Late Bathonian nominal species that occurs in Kachchh, Madagascar and Europe (Fig. 4). Its biostratigraphic and biogeographic implications are also discussed.

#### 2. Geological setting

Kachchh is a pericratonic epicontinental basin borne out from the rifting of the Indian plate from the Gondwana in the Late Triassic with the deposition of nearly 1500 m of carbonates and siliclastics (Jain et al. 1996). Residual basement ridges, now exposed as domal outcrops or uplifts that provide an opportunity to study its faunal and sedimentary history during the Mesozoic, characterize the basin. The six major uplifts (Kachchh Mainland, Pachchham, Khadir, Bela, Chorad and Wagad (Fig. 1), constitute islands in an otherwise flat lying surface with the latter five being collectively referred as "Islands" (Fig. 1). The domal outcrops within the Kachchh Mainland (like Jumara and Jara, the localities studied for this contribution; see Fig. 1) follow an East–West trending lineament with the general regional slope of the basin being WSW.

#### 2.1. Jara Dome

The Jara Dome (Fig. 2), like all other domal outcrops in Kachchh is heavily faulted with a 248 m thick succession of Middle Jurassic rocks exposed in the northwestern fringe of the Kachchh Mainland, western India (23°43'00"N: 68°57'52"E to 23°45'00"N: 65°00'00"E). Here, as well as in other domal outcrops, the exact delineation of the Bathonian–Callovian boundary has remained elusive in spite of the fact that the basal sediments (ash gray marl–limestone alternations within the core of the dome) bear striking lithological resemblance with the adjacent well-dated Late Bathonian Sponge beds of the Patcham Formation exposed at the Jumara Dome (Callomon 1993; Jain & Pandey 2000). Alt-



**Fig. 2: (a)** Jara is a heavily faulted quaquaversal domal outcrop yielding Late Bathonian-?Middle Oxfordian ammonites. **(b, c)** The core of the dome is best exposed as a linear outcrop within the center of the dome from where dimorphic occurrence of the Indonesian ammonites *Macrocephalites keeuwensis* Boehm is reported here associated with the Late Bathonian *Macrocephalites* cf. *mantataranus* Boehm [M] from the basal sediments (marl–limestone intercalations). The late Early–early Middle Callovian Zoophycos Beds I and II (Ridge Sandstone Member) form prominent ridges, followed by shales and then another cliff–forming Early–?Middle Oxfordian Dhosa Oolite bed. The shales between the core and the Zoophycos Bed I are *marcocephalites*–bearing, forming biozones (a). A detailed biozonation will be published elsewhere. Fig. 2b, courtesy Bhawanisigh G. Desai.

hough, Jumara is located barely ~12 km east of Jara (Fig. 1), unlike at Jumara, stratigraphically controlled ammonite records from the basal sediments at Jara have only been noted in passing (Bhaumik et al. 1993; Krishna & Ojha 2000), resulting in conflicting ages that have ranged from Late Bathonian to Early Callovian (Prasad 1998; Bhaumik et al. 1993). Even the nannofossil record has been equally ambiguous (Rai, 2003). Recently, the find of M. cf. mantataranus Boehm [M] from the basal sediments at Jara (Jain & Desai 2014) have enabled to fix a definite Late Bathonian age to the core sediments of the Jara Dome (Fig. 2: site 1). From the same sediments, the dimorphic pair of the Indonesian ammonite Macrocephalites keeuwensis Boehm [M & m] is now reported and described below.

#### 2.2. Jumara Dome

The Jumara Dome (Fig. 1; 23°41'13.6"N; 69°02' 51.00"E), the depocentre of the basin is a classical Jurassic locality that exhibits not only the most expanded Callovian sequence for the Kachchh basin but also for the entire northern peri-Gondwana margin (see Jain 2014 for a detailed description and faunal content of the dome). The dome possesses 422 m thick sequence of Middle Jurassic strata constituting 8 beds in the Patcham Formation (Bathonian; A1–A8; 48 m), 46 beds in the Lower Chari Formation (Lower Callovian; B1–B45 and C1; 229 m), 37 beds in the Middle Chari Formation (Middle Callovian (C2–C37 and D1; 68 m) and 14 beds (D2–D15) in the Upper Chari Formation (Late Callovian; 74 m). Bed E1, the Dhosa Oolite bed (Early–?Middle Oxfordian) tops the Chari Formation (see also Jain 1996). The present study is restricted to the topmost unit of the Patcham Formation, bed A8, the ash gray marl–limestone alternations (Fig. 3).

### 3. Age of *Macrocephalites keeuwensis* Boehm

*Macrocephalites keeuwensis* Boehm was first described from Indonesia, where the fossil record is characterized by "...faunas that can be securely dated are very intermittent" and "the stratigraphic record, at least as revealed by ammonites, is regionally highly discontinuous" (Westermann & Callomon 1988: p. 6). The Bathonian–Callovian fauna of Indonesia collected from 15 stratigraphical separate assemblages from Taliabu in the Sula Islands (the classical area of "Keeuw") and Strickland River, Papua New Guinea (see also Sato et al. 1978; Westermann et al. 1979) were recently re-evaluated and placed chronologically by Westermann & Callomon (1988) (see Fig. 4).

The youngest of all is the *Macrocephalites keeuwensis* Association dated on balance "as late Early Callovian, although latest Bathonian cannot be



Fig. 3: Late Bathonian exposures at Jara (a) and Jumara Domes (b) with their ammonite content.



Fig. 4: Indonesian ammonite associations, ammonite content and correlation with Kachchh and NW European faunal content (see text for explanation).

ruled out" (Westermann & Callomon 1988: p. 6). The Association is characterized by the presence of *M. keeuwensis* Boehm with several of its varieties ( $\alpha$ – $\delta$ ; see also Boehm 1912; Spath 1927–33; Westermann & Callomon 1988). Besides the nominal species, this youngest assemblage also contains endemic *Macrocephalites* cf. *bujabesari*, *M.* cf. *wichmanni*, *M.* cf. *folliformis*, *Oxyerites sulaensis* Westermann and Callomon, and the cosmopolitan *Choffatia* aff. or cf. *furcula* (see Fig. 4). One of the specimens from this assemblage (figured by Boehm 1912: pl. 38, fig. 2) closely matches with the recently collected *Macrocephalites* from the Late Bathonian Orbis Zone of Franconia, South Germany (Dietl & Callomon 1988).

Now the dimorphic pair of Macrocephalites keeuwensis Boehm [M & m] (Figs 5, 6) in association with typical Late Bathonian M. cf. manatatranus Boehm [M] (Jain & Desai 2014) from the Jara Dome and M. keeuwensis var. aff. forma flexuosa [m] (Fig. 7) associated with M. cf. manatatranus Boehm [M] (Fig. 8) along with *M. triangularis* Spath [M] and Sivajiceras congener [M] from the top of the Patcham Sponge Beds (bed A8 of present work) from the Jumara Dome are reported here. At Jumara, coeval levels have also yielded characteristic Orbis-Discus Zone fossils of Oxycerites cf. orbis (Roy et al. 2007), O. cf. oppeli (Pandey et al. 2012) and Bullatimorphites cf. hannoveranus (Bardhan et al. 2002) (Fig. 3). Hence, based on the common presence of Indonesian Macrocephalites keeuwensis Boehm [m] in Kachchh and Madagascar (Thierry 1978) and in Europe (Fig. 6f), the M. keeuwensis Association in Indonesia can now be unequivocally dated as Latest Bathonian (Fig. 4).

In central Nepal, only the subjacent Bifurcatus and Apertus Association of Sula Island (Indonesia) (Fig. 4) have been recorded with an unconformity at the top of the Apertus Association (Cariou & Enay 1999). The Apertus Association ammonites come from two

levels (see Cariou & Enay 1999: p. 704). From the lower level, bed 10 yielded Macrocephalites apertus Spath, M. mantataranus Boehm, M. bifurcatus transient bifurcatus Boehm [m], M. bifurcatus transient intermedius (Spath) [M], Homeoplanulites (Parachoffatia) aff. evolutum Sandoval and Oxycerites sulaensis Westermann and Callomon. Bed 11 yielded Macrocephalites triangularis Spath, M. chrysoolithicus (Waagen), Choffatia cf. madani Spath and Oxycerites sulaensis Westermann and Callomon. Cariou & Enay (1999) considered their bed 10 as Late Bathonian (=Apertus Association of Sula Island) and their bed 11 as Early Callovian. Clearly, the latter should have also been dated as Latest Bathonian, in commonality with the Kachchh Late Bathonian Triangularis Zone fauna (Fig. 2; see also Jain 2014).

#### 4. Discussion

Based on present finds, Macrocephalites keeuwensis Boehm [m] (Fig. 6a-d) now occurs in definitely Late Bathonian sediments in Kachchh, Madagascar and Europe (Orbis Zone; see Dietl & Callomon 1988; Thierry 1978) (Fig. 4). M. cf. manatatranus Boehm [M] has already been recorded from the Triangularis Zone (= Orbis–Discus Zones) of Jumara (Fig. 8) in association with M. keeuwensis var. aff. forma flexuosa [m] (Fig. 7), *M. triangularis* Spath [M] and Sivajiceras congener (Waagen) [M] and the characteristic Orbis-Discus Zone fossils of Oxycerites cf. orbis (Roy et al. 2007), O. cf. oppeli (Pandey et al. 2012), Bullatimorphites cf. hannoveranus (Bardhan et al. 2002) from the top of the Patcham Sponge Beds (bed A8 of present work) (Fig. 3) (see also Jain & Desai 2014). M. manatatranus Boehm [M] also occurs in Central Nepal in Late Bathonian sediments (Cariou and Enay, 1999). M. triangularis and S. congener are long ranging forms (late Middle to Latest Bathonian; Jain



**Fig. 5:** *Macrocephalites keeuwensis* Boehm [M]. **(a–c)** Jara, specimen no. Ja/SJ/1999/Core/1, measured diameter = 172 mm, phragmocone = 132 mm. **(d–f)** Specimen lent to the author by the late Dr. G.E.G. Westermann (Canada), specimen no. J2023W, Locality 4A, Assemblage V1 of Tikong, Indonesia, phragmocone = 103 mm, measured diameter = 115.3 mm (see text for explanation).



Fig. 6: Macrocephalites keeuwensis Boehm [m]. (a–d) Jara specimen no. Ja/SJ/1999/Core/2, measured diameter = 107 mm, phragmocone = 80 mm. (e) *M. keeuwensis* var. B [m] refigured from Boehm (1912: pl. 38, fig. 1; see also Westermann & Callomon 1988: pl. 13, fig. 1). (f) *Macrocephalites* cf. / aff. *keeuwensis* Boehm, Late Bathonian (Orbis Zone), refigured from Dietl & Callomon (1988: pl. 3, fig. 1).

2014) and *Oxycerites sulaensis* Westermann and Callomon which occurs in the Indonesian *M. keeuwensis* Association is a close homeomorph of the well-known Early Bathonian European *O. yeovilensis* Rollier (Westermann & Callomon 1988). The other

persisting form is that of *Choffatia* (*Homoeoplanu-lites*), which had already appeared in the late Middle Bathonian Bremeri Zone, and is also long ranging.

Contextually, both associations at Sula Island (Westermann & Callomon 1988) and Madagascar

(Thierry 1978) are discontinuous, and hence hard to date precisely, whereas in Central Nepal, the fauna above the Apertus Zone is not present (Cariou & Enay 1999). Macrocephalites apertus Boehm has not been recorded from Kachchh. Hence, the best available choice to date the M. keeuwensis Association lies either in the assemblages of Kachchh or in Europe (Dietl 1981; Dietl & Callomon 1988; Callomon 1993; Jain 2014). Common occurrence of index forms are noted in both the latter two localities (Fig. 4), although in the Franconian Alb, it is much more well-resolved (Dietl 1981; Dietl & Callomon 1988; Mönnig 2015). Nevertheless, based on common species (Fig. 4), the M. keeuwensis Association can be safely dated to lie around the Orbis-Discus Zones of Latest Bathonian age, as also earlier suggested by Westermann & Callomon (1988: p. 6) but due to lack of characteristic species, now noted, hampered precise age determination, then.

Based on present finds and on stratigraphic, morphological and morphometric considerations, it is speculated, and also probably plausible, that M. keeuwensis Boehm [M & m] gave rise to two distinct lineages and the paper documents the separation of these two different lines. One, that gave rise to the Early Callovian M. lamellosus (Sowerby) and that in turn to the late Early Callovian Subkossmatia opis Spath (= the Eucycloceras opis var. opis of Jana et al. 2005) (Fig. 9). In this progression, the ribbing gets straighter, finer, denser and the shell increasingly gets compressed and more eccentrically coiled, but at the same time, retaining the somewhat subquadrate whorl section (Fig. 9). On the other hand, to the early-middle Early Callovian Macrocephalites formosus (Waagen), which in turn gave rise to the late Early Callovian M. semilaevis Spath.

Dimorphic pairs of *M. lamellosus* have now been recorded from the late Early Callovian Sameilaevis Zone (Fig. 10). Earlier, Krishna & Westermann (1987: p. 1576) have suggested that "there is a close agreement between the phragmocones of M. lamellosus [m] and the inner whorls of M. madagascariensis [M] in our samples, indicating that they are matching dimorphs". However, it is always best to look for dimorphism within the morphospecies rather than outside it, as noted in the present record (Fig. 10). The Macroconch, which is complete with an attached operculum, measures 150 mm with a phragmocone of 115 mm (T/H = 0.80; U/D = 0.22) (Fig. 10a, b). The holotype, a microconch and almost complete with only part of the body chamber damaged (Fig. 10c, d) measures 108 mm with a phragmocone of 71 mm (T/H = 0.89; U/D = 0.19). The macroconch shows increased extraumbilication (Fig. 10a). In both dimorphs, mature modification only includes strongly projected ribs and opening of the whorl (Fig. 10). Krishna & Westermann (1987) paired M. lamellosus [m] - M. madagascariensis [M], but, in contrast, Thierry (1978) considered the pairing of M. subcompressus (Waagen) [m] and M. madagascariensis

[M]. The holotype of *M. madagascariensis* [a Macroconch] measures 130 mm with a phragmocone of 109 mm and is much more inflated (T/H = 0.94-0.99) and involute (U/D = 0.14-0.15) at 130 and 109 mm diameters, respectively. Based on present finds, the *M. subcompressus* [m] – *M. madagascariensis* [M] pair seems more plausible.

On the other hand, the Kachchh Macrocephalites keeuwensis [M & m] (Fig. 5a-c), probably also gave rise to the early-middle Early Callovian M. formosus (Sowerby) (Fig. 11a-e) and that in turn gave rise to the late Early Callovian M. sameilaevis (Waagen). Krishna & Westermann (1987: p. 1576) suggested that M. chariensis and M. formosus intergrade and hence considered them to represent as a single species. However, here, following Spath (1927–33), both are considered as separate species with their dimorphic pairs (Fig. 11; M. formosus - Figs. 11a-e; M. chariensis - Figs. 11f-i). Both dimorphs are now recorded from the same bed (bed 23) from the middle-Early Callovian Diadematus Zone from the Jumara Dome. The Macroconch is large, discoidal and smooth measuring 230 mm with its microconch measuring 64 mm (Fig. 11a-e). The dimorphic pairs of M. chaeiensis (Waagen) are almost complete specimens with only part of the body chamber damaged (Fig. 11f-i). The Microconch comes from a stratigraphically slightly lower horizon, the middle-Early Callovian Diadematus Zone (Jumara Dome; diameter = 86 mm; T/H = 0.88, U/D = 0.14) and the Macroconch comes from the late Early Callovian Formosus Zone (Jumara Dome; diameter = 101 mm; T/H = 0.79, U/D = 0.14) (Fig. 12). In both the microconchs (Fig. 11a-f) and the macroconchs of M. chaeiensis, the crowding of ribs is noted; the lack of umbilical seam in all specimens, also suggests maturity (Fig. 11). Basinal biozonation for the localities mentioned (Keera, Jumara and Jara; Fig. 1) is given in Figure 12 and overall correlation with the Western European biozonation is given in Figure 13. A more detailed Early Callovian biozonation is being worked out and will be published elsewhere.

Earlier, Krishna & Cariou (1993) had suggested that *M. formosus* gave rise to *M. sameilaevis* (Fig. 14) and at the same time to *Subkossmatia opis*, also. The other lineage they proposed was their "Chrysoolithicus lineage" – *M. triangularis* – *M. madagascariensis* – *M. transitorius* – *M. chrysoolithicus* – *M. diadematus*. Contrary to their claim, the Kachchh ammonite fauna is not stratigraphically differentiated; the compressed species do occur together with the inflated species, although their relative percentage may differ and they themselves differ laterally in abundance from one dome to the other (Fig. 15).

Additionally, it is also speculated that both microconch [m] and macroconch [M] of *M. keeuwensis* Boehm, in turn, probably arose from *M. bifurcatus* transient *intermedius* (from one of its varieties, A or B). All the above forms have been recorded from Kachchh (Jain 1996, 2014).



Fig. 7: Macrocephalites keeuwensis Boehm aff. var. forma flexuosa [m]. (a–c) Jumara, specimen no. Ju/22/122. (d–e) Jumara, specimen no. Ju/22/148. Both specimens (a–e) come from the top of the Sponge Beds (bed A8; see Fig. 3), Patcham Formation, Jumara, (b) is the close–up of (a) to show the umbilical region. (f) Macrocephalites keeuwensis var. forma flexuosa [m] refigured from Westermann & Callomon 1988: pl. 12, fig. 4; IMC 444, Assemblage VI; see Fig. 4). (g–i) Macrocephalites keeuwensis var. forma flexuosa [M] refigured from Westermann & Callomon 1988: pl. 13, fig. 3a, b; IMC 447, Assemblage VI; see Fig. 4). Fig. 7i is reconstructed from Fig. 7g.



**Fig. 8:** *Macrocephalites* cf. *mantataranus* Boehm [M], fully septate specimen from bed A8 (Sponge Beds), Jumara, specimen no. Ju/22/8c, measured diameter = 110 mm (H: 52 mm; T = 46; U = 15 mm (see Jain, 1996; Jain and Desai, 2014).

#### 5. Paleobiogeography

Seyed-Emami et al. (2015) noted in their description of Macrocephalitinae from Iran that Latest Bathonian-Earliest Callovian duration is marked by the dominance of Macrocephalites and Homoeoplanulites in both Subboreal and Submediterranean Provinces, whereas in Kachchh (and in general, within the Indo-Madagash Province), Macrocephalites and Choffatia dominate (and to a minor extent subgenus Subgrossouvria and Genus Indosphinctes). Genus Macrocephalites, in Europe, occurs in the Late Bathonian of Southern Germany where now both Macrocephalites triangularis and M. cf. or aff. keeuwensis (Dietl & Callomon 1988; = M. cf. keeuwensis of Mönnig 2015) have been recorded; the latter comes from the middle-Late Bathonian Hannoveranus subzone, Retrocostatum Zone (Dietl & Callomon 1988; Mönnig

1996, 2014) (Fig. 13). Sporadic occurrences of Late Bathonian–Early Callovian *M. madagascariensis* and the Early Callovian *M. chrysoolithicus*, *M. lamellosus* and *M. subcompressus* have also been mentioned occasionally (see also Cariou et al. 1990).

It seems that the small, compressed and sharply ribbed Macrocephalites keeuwensis Boehm [m] (Fig. 4) is cosmopolitan, following the Late Bathonian transgression and migrated from Indonesia via Kachchh to Madagascar and on to Europe, along the southern peri-gondwana margin. But, it did not reach further to Iran, Tibet and the Russian platform, however, the genus, as such, did (see also Mönnig 2015). Compressed forms such as M. lamellosus Sowerby [m] and M. cf. subcompressus (Waagen) [m] have been recorded from Iran and M. madagascariensis Lemoine and M. semilaevis (Waagen) [M] from Tibet (Yin 2016) suggesting dilution of ammonite faunal provincialism and increased exchanges between the Indo-Madagascan and other provinces throughout the Early Callovian, and more so at the base of Early Callovian.

#### 6. Systematic paleontology

All specimens presented here are part of the set of specimens that formed the basis for the author's dissertation and are accordingly kept in the collcetion of the Department of Geology, University of Rajasthan (Jaipur, India).

Genus Macrocephalites Zittel, 1884

Type species: Ammonites macrocephalus Schlotheim, 1813

Macrocephalites keeuwensis Boehm 1912 [M and m]

Figs. 5-7, 16, 17

#### Microconch [m]

- 1890 *Ammonites calloviensis* Etheridge, p. 175, pl. 29, fig. 5.
- 1908 Macrocephalites cf. macrocephalus compressus Boehm, p. 331, pl. 12, fig. 2.
- Macrocephalites keeuwensis a Boehm, p. 160, pl.
   36, figs 2–4 only [fig. 3a, b, lectotype].
- 1912 Macrocephalites keeuwensis  $\beta$  Boehm, p. 161, pl. 38, figs 1 and 2a, b only.
- 1912 *Macrocephalites keeuwensis*  $a \beta$  Boehm, p. 163, pl. 41, fig. 3.
- 1913 Macrocephalites keeuwensis Boehm, p. 15, pl. 5, fig. 1.
- 1928 Dolikephalites keeuwensis Spath, p. 205 [lectotype designated: Boehm 1912, pl. 36, fig. 3a, b].
- 1928 Dolikephalites flexuosus Spath, p. 210, pl. 32, fig.
  4. [for Etheridge 1890, pl. 29, fig. 5; holotype refigured].
- 1958 *Macrocephalites flexuosus* Collignon, pl. 18, fig. 77.
- 1978 Macrocephalites (Dolikephalites) keeuwensis Sato et al., p. 12.



**Fig. 9:** Proposed evolution of *Macrocephalites keeuwensis – M. lamellosus – Subkossmatia opis*. **(a–d)** *Subkossmatia opis* Spath. a: Microconch refigured from Jana et al. (2005), their *Eucycloceras opis* var. *opis* [m], specimen no. JUM 378, bed 6 from Keera, diameter = 98 mm. b–d: holotype [m] refigured from Spath (1928: pl. 39, figs 2, 7). **(e–g)** *Macrocephalites lamellosus* (Sowerby). Microconch holotype refigured from Spath (1928: pl. 39, figs 2, 7). **(e–g)** *Macrocephalites lamellosus* (Sowerby). Microconch holotype refigured from Spath (1928: pl. 39, figs 2, 7). **(e–g)** *Macrocephalites lamellosus* (Sowerby). Microconch holotype refigured from Spath (1928: pl. 19, fig. 8a, b), measured diameter = 108 mm; phragmocone = 71 mm, Keera Dome, Kachchh (see text and Fig. 10c, d for details). **(h–j)** *Macrocephalites keeuwensis* Boehm [m] from Jara, present study, see text and Fig. 6a–c for details. In this progression (*M. keeuwensis–M. lamellosus–S. opis*), the ribs get straighter, finer, denser and the shell increasingly gets compressed and more eccentrically coiled, but at the same time, retaining the somewhat subquadrate whorl section.



**Fig. 10:** *Macrocephalites lamellosus* (Sowerby) [M and m]. The dimorphic pair have been recorded from the late Early Callovian Sameilaevis Zone (see Fig. 13). **(a, b)** [M]: specimen no. Ju/12/25, Jumara, bed B36, Sameilaevis Zone, measured diameter = 150 mm, phragmocone = 115 mm (at D: 150, H: 66.5; T: 53.3; U = 33.6; T/H = 0.80; U/D = 0.22). **(c, d)** [m]: holotype refigured from Spath (1928, pl. 19, figs. 8a–b), measured diameter = 108 mm; phragmocone = 71 mm (at D: 108, H: 49.7; T: 44.3; U = 20.4; T/H = 0.89; U/D = 0.19). The microconch come from the adjoining Keera Dome, Kachchh (see Fig. 1), middle–Early Callovian Diadematus Zone (see Fig. 13). Note that the Macroconch is complete with attached operculum (b) and the microconch has only part of the body chamber damaged and is almost complete. In both dimorphs, ribbing projects strongly at the end of the whorl. Krishna & Westermann (1987) had suggested dimorphic pairing between *M. lamellosus* [m] and *M. madagascariensis* [M]. However, Thierry (1978) considered the pairing of *M. subcompressus* [m] and *M. madagascariensis* [a Macroconch] measures 130 mm with a phragmocone of 109 mm; it is much more inflated (T/H = 0.94–0.99) and involute (U/D = 0.14–0.15) at 130 and 109 mm diameters, respectively. Based on present find, the latter seems more plausible.



**Fig. 11:** *Macrocephalites formosus* (Sowerby) [M and m] and *M. chaeiensis* (Waagen) [M and m]. **(a–e)** *M. formosus* (Sowerby) [M and m], both from bed B23, middle–Early Callovian Diadematus Zone, Jumara (see Fig. 13). a–b: Macroconch, specimen no. Ju/12/20, measured diameter = 230 mm, c–e: Microconch, specimen no. Ju/12/50, measured diameter = 64 mm. f–i: *Macrocephalites chaeiensis* (Waagen) [M and m], almost complete specimens with part of the body chamber damaged. **(f)** [m], specimen no. Ju/12/40, bed B29, middle–Early Callovian Diadematus Zone (see Fig. 13), Jumara measured diameter = 86 mm (H = 45, T = 39.6, U = 8.1, T/H = 0.88, U/D = 0.14). **(g–i)** [M], specimen no. Ju/9/185, bed B29, late Early Callovian Formosus Zone, measured diameter = 101 mm (H = 57.4, T = 55.7, U = 13.82, T/H = 0.79, U/D = 0.14) (see Fig. 13). In both the microconchs (a–f), crowding of ribs is noted. Lack of umbilical seam in all specimens, also suggests maturity.

- 1978 Macrocephalites (Dolikephalites) cf. subcompressus – Sato et al., p. 12.
- 1978 Macrocephalites macrocephalus keeuwensis dim. micr. [= Dolikephalites flexuosus] – Thierry, p. 285, pl. 21, fig. 9 a, b [Boehm 1912, pl. 36, fig. 3 a, b, refigured = lectotype Spath 1928, textfig. 103].
- 1988
   Macrocephalites keeuwensis Boehm 1912 [m] –

   Westermann & Callomon, p. 63, pl. 11, fig. 5; pl.
   12; pl.

   12; pl.
   13, figs 1, 2.
- 1988 Macrocephalites cf. / aff. keeuwensis Boehm 1912 [m] – Dietl & Callomon, pl. 3, fig. 1
- cf. 1996 Macrocephalites subcompressus (Waagen) Jain, p. 81, pl. 10, fig. 4
- cf. 1996 Macrocephalites flexuosa (Spath) Jain, p. 85, pl. 11, fig. 3

#### Macroconch [M]

- cf. 1912 Macrocephalites keeuwensis a γ Boehm, p. 164, pl. 42, fig. 2.
  1912 Macrocephalites keeuwensis a γ Boehm, p. 164.
  - 1912 Macrocephalites keeuwensis α γ Boehm, p. 164, pl. 43, fig. 1.
- cf. 1912 *Macrocephalites* sp. Boehm, p. 166, pl. 44, fig. 1a, b.

1978 Macrocephalites (Macrocephalites) aff. fomosus mantataranus – Sato et al., p. 12 only.

cf. 1978 Macrocephalites macrocephalus keeuwensis [=M. keeuwensis  $a - \gamma$ ] – Thierry, p. 282, pl. 18, fig. 2a, b [?Boehm 1912, pl. 42, fig. 2 refigured], textfig. 102 (part).

Type: Boehm 1912: pl. 36, fig. 3a, b; lectotype designated by Spath 1928, p. 205 [m].

Material: 5 fragmentary specimens; One unillustrated macroconch [M] lent by Late G. E. G. Westermann, (1994, during the author's dissertation) from locality 4A, Assemblage V1 of Tikong, Indonesia (specimen no. J2023W mentioned in Westermann and Callomon, 1988, p. 63) (Fig. 3d–f); 2 specimens from the core of Jara Dome – one [m], var. B (specimen no. Ja/SJ/1999/Core/1); +2.5 m from the base; one [M] (specimen no. Ja/SJ/1999/Core/2) (Fig. 3a– c); 2 specimens from top of Sponge Bed (A8) of the Jumara Dome – var. aff. forma *flexuosa* [m] (specimen nos. Ju/22/122 and 148) (Fig. 7a–e).

Remarks [M]: In 1994, during my dissertation, the late Dr. G.E.G. Westermann (Canada) lent me a specimen (J2023W from Locality 4A, Assemblage V1 of Tikong, Indonesia) (Fig. 4, figured in Fig. 5d-f) of a macroconch of Macrocephalites keeuwensis Boehm which he had mentioned in his publication (Westermann & Callomon 1988) as material on p. 63 and on page 66 it was specifically mentioned - "The second, crushed specimen is somewhat larger with a (reconstructed) estimated diameter of 180 mm, and its body chamber has entirely smooth flanks." I am here illustrating this specimen for the first time as now comparable forms have been found in Kachchh from Jara; both are described below (Fig. 5). The phragmocone of J2023W measures 103 mm and adding a standard 3/4 additional whorl for body chamber, the estimated diameter measures to 180

mm. The Macroconch of *Macrocephalites keeuwensis* Boehm described by Westermann & Callomon (1988: pl. 13, fig. 3a–c) is the Macroconch of one of their formas (forma *flexuosa*) (re-figured in Fig. 7g–i; Fig. 7i is reconstructed from 7h).

Thierry (1978: p. 282) noted that 20 macroconch specimens were available to him from Boehm's collection (Geological Institute of Utrecht, Holland), only 3 were (in part) with preserved body chamber measuring 163, 148 and 130 mm diameters. He choose the specimen illustrated in plate 42, figure 2 (Boehm 1912) from Keeuw (Taliabu Island, Sula Islands), Indonesia as type (refigured here in Fig. 16). This is a smooth, compressed form with a somewhat triangular whorl section; the body chamber occupies half to three quarters of a whorl. Thierry (1978) further noted that M. keeuwensis Boehm is a small-sized species (adult diameter 150-170 mm), discoidal, ornamented with moderately fine to medium ribbing. The phragmocone measures 90-100 mm with an oval whorl section. At this stage, the primary ribs are numerous and straight, slightly flexing towards the end of the phragmocone and dividing (mono-, di-, or polyschizotomes) into secondary ribs, to which are added numerous intercalary ribs, all passing the ventral border straight. Umbilical edge remains sharp and the walls vertical (Fig. 16). The early stages are unknown.

Description: Macroconch [M]-Shell is large (diameter = 172 mm; phragmocone = 132 mm; maximum estimated diameter = 231 mm), nearly complete (judging by the absence of umbilical seam) and discoidal. Early whorls, though not clearly discernable, are moderately depressed (see Figs. 5b-c; T/H = 1.20). At the body chamber, the umbilicus is narrow (U/D = 0.14) and funnel-shaped (66% broader at the top than at the bottom; conical; Fig. 5a). Umbilical margin is sharp. Flanks are moderately converging with a broadly rounded, somewhat tabulate venter. Whorl section changes from subtriangular to somewhat subquadrate, from the phragmocone to the body chamber, respectively. Ribbing is moderately fine with long primaries that arise from just below the umbilical shoulder with a slight backward bent at the umbilical region and then project forward, dividing indistinctly on the outer third of the flank into two or three secondaries that cross the venter straight. There are ~15 primaries per half whorl at the body chamber. They persist well into the body chamber but fade by 160 mm with an eventual smoothening of the lateral flanks. The only ribbing modifications noted at the body chamber is that the primaries become somewhat more forwardly projected than they are on the visible end part of the phragmocone (Fig. 5a); in the latter part, they are somewhat straighter, radial and closely spaced. However, with regards to lateral smoothening (the other adult modification), in the crushed Indonesian specimen (Fig. 5d), it is somewhat early (at the end of the phragmocone at



Fig. 12: Biozonation at the three domal localities of Kachchh discussed in this contribution. (a) After Krishna and Ojha (1996). (b, c) present study from Jumara and Jara Domes, respectively. Both Jara and Jumara records are from the Late Bathonian Triangularis Zone.

102 mm) and very pronounced with secondaries restricted to the outer third of the flank, thereafter.

Microconch [m]-Macrocephalites keeuwensis Boehm var. B [m]: Shell of medium diameter (107 mm), and nearly complete (as suggested by the absence of umbilical seam and the strong attenuation of ribbing; Figs. 6a, b), moderately evolute (U/D = 0.16-0.20, compressed (T/H = 0.79) and densely ribbed with ~23 primaries and ~51 secondaries per half whorl. Phragmocone (80 mm) is typically involute (U/D = 0.16), with a compressed subrectangular to trapezoidal whorl section (Fig. 6d). The primaries arise obliquely on the umbilical wall, curve with strong adoral concavity across the umbilical margin and reach with gentle projection to almost middleflank. At the end of the body chamber, they tend to project more strongly and are thus are more concave at the umbilical region. The secondaries arise mostly by virgatotome polyfurcation and by intercalation at about middle-flank but at somewhat uneven whorlheight. They are more or less markedly rectiradiate and cross the venter straight at the phragmocone and with a slight and broad adoral convexity at the body chamber. The umbilical wall remains vertical, and the edge sharp. The flanks become subparallel to gently converging with a broad, evenly rounded

and slightly tabulate venter. The ribbing, even at the body chamber, remains dense but increases somewhat in spacing; it is still very sharp and prominent. At the body chamber, the shell egresses, resulting in a somewhat elliptical coil.

Macrocephalites keeuwensis Boehm aff. var. forma flexuosa [m]

Shell evolute, compressed and fragmentary but shows strong projection and increased closeness of ribs (crowding) on the body chamber (Fig. 7a, b, d) suggesting shell maturity reminiscent of that noted in *Macrocephalites keeuwensis* var. forma *flexuosa* Boehm [m] (Westermann & Callomon 1988: pl. 11, fig. 5a) (refigured in Fig. 7f [m]). The umbilical edge, though little available for observation, is characteristically sharp and the wall slanting inward. Primaries and secondaries, per quarter whorl, number ~12 and ~28, respectively. No other comparable Kachchh macrocephalitids are known at the coeval Bathonian–Callovian interval.

After	Noi	thwest Europe	1997.	Kachchh							
Thierry et al., 1997, Seyed-Emami et al., 2015, Mönnig. 2014				After Jain, 2014, This study							
Subzone				mn	Assemblage						
Middle Callovia	Anceps	Stuebeli		S. opis E. eucyc	Subkossmatia opis, Reineckeia (R.) anceps, Macrocephalites subcompressus, Rehmanni (L.) reis: Eucycleceras eucyclum						
Early Callovian	Gracilis	Patina	M. tumidus	M. semilaevis	Macrocephalites semilaevis, M. magnumbilicatus, M. subcompressus, Macrocephalites sp., Rehmanni (Loczyceras) reissi						
		Michalskii			Macrocephalites formosus, M. dimerus, M. lamellosus, M. chariensis, M. subcompressus B. (Kheraiceras) bullatus, Parapatoceras tuberculatum, Hectococeras proximum, Indosphinotes peregrinus, Coliotia oxyptycha, Reineckeia tyranniformis, Indosphinctes urbanus	Formosus					
		Laugieri	dicosmur gracilis		Macrocephalites diadematus, M. chrysoolithicus, M. dimerus, M. cf. dimerus, M. formosus, M. lamellosus, M. chariensis, M. subcompressus Hecticoceras michalski, Oxycerites cf. pleurocamya, Paralcidia sp., Parapatoceras tuberculatum, P. distans, Bullatimorphites (Bomburites) densicostatum, Paroecotraustes (Oecotraustes) sp., Calliphylloceras disputable.						
		Pictava	W.	dematus		Diadematus					
		Grossouvrei		M. dia oolithicus formosus losus	Subgróssouvria gudjinsirensis						
		Prahecquense	M. gracilis ssp. A	M. chryss	Macrocephalites dimerus, M. diadematus, M. chrysoolithicus, M. transitorius, M. formosus, M. lamellosus, M. chariensis, M. cf. sphaerodalis, M. subcompressus, M. elephantinus						
	Bullatus	Bullatus / Jadnoti W. verus / Jadnoti	M. verus	A. transitorius	Choffatia madani, Choffatia sp., Indosphinctes sp., Subgrossouvria sp., Indosphinctes urbanus, Indosphinctes errans, Subgrossouvria gudjinsirensis var. tenius, Macrocephalites madagascariensis, M. dimerus,	Di					
			M. verus / jaquoti	M. madagascariensis	M. transitorius, M. elephantinus, M. formosus, M. lamellosus, M. chariensis, M. diadematus, M. subcompressus, Parapatoceras sp., Choffatia sp., Choffatia baluchistanensis, B. (K.) cf. cosmopolitum, M. madagascariensis, M. dimerus, M. transitorius, M. elephantinus, M. lamellosus, M. habyensis, M. intermedius, M. diadematus, M. subcompressus, Bullatimorphites (K.) bullatus, Bullatimorphites sp., B. (Kheraiceras) cosmopolitum, I. peregrinus, Calliphylloceras sp., Choffatia subbakeriae, Choffatia baluchistanensis, Subgrossouvria sp., Indosphinctes sp.	Madagascariensis					
					GAP						
Middle Bathonian Late Bathonian	Discus	Discus Angulicostatum (Hollandi)	M. jaquoti A. triangularis vensis	M. keeuwensis M. mantataranus	<ul> <li>Oxycerites orbis, Sivajiceras congener, Choffatia sp., Homoeoplanulites sp., Macrocephalites triangularis, M. cf. dimerus, M. cf. madagascariensis, M. cf. mantataranus, M. keeuwensis, Parapatoceras sp., Oxycerites sp., Paracenoceras sp., P. hexagonoides, P. wandaensis, P. jumarensis, Oxycerites cf. oppeli, Subgrossouvria sp., Choffatia baluchistanensis, Choffatia subbakeriae</li> </ul>	riangularis					
	Bremeri Retrocostatum	Hannoveranus (Histricoides)	A M. cf. keeuv	trianoularis	Epistrenoceras sp., B. (Kheraiceras) cf. hannoveranus, Choffatia spp., Homoeoplanulites spp., Sivajiceras congener, Macrocephalites triangularis, Paracenoceras sp., Choffatia balinensis	F					
		Blanazense / Julii		W	Procerites cf. hodsoni, Procerites spp., Choffatia spp., Homoeoplanulites spp., Sivajiceras congener, Macrocephalites triangularis, Subgrossouvria sp., Procymatoceras cf. intumenscens, Parapatoceras tuberculatum, Micromphalites (Clydomphalites) clydocromphalus.	Procerites					
		Fortecostatum		M. cf. etheridgei M. bifurcatus	Wagnericeras sp., Sivajiceras congener, Procerites hians, P. (Gracilisphinctes) arkelli, P. (G.) intermedius, Macrocephalites triangularis, M. cf. etheridgei, M. bifurcatus aff. bifurcatus, M. bifurcatus intermedius, Parapatoceras distans, Reineckeia sp. A & B, Epimorphoceras decorum,						
		Bullatimorphus		w. triangularis	Clydoniceras traingulare, Bullatimorphites s.s.	Arkell					
L											

**Fig. 13:** Bathonian–Callovian ammonite assemblages and biozonation at Kachchh and correlation with the Western European biozonation. Zitteliana 93 18



**Fig. 14: (a, b)** *Macrocephalites semilaevis* (Waagen) [M], fully septate specimen, specimen no. Ju/7/76, bed B36, late Early Callovian Semilaevis Zone (see Fig. 13), Jumara, diameter = 101 mm.

Remarks: Macroconch [M]: Thierry (1978: p. 282, pl. 18, fig. 2a, b; pl. 20, fig. 5a, b) illustrated two macroconch specimens; the one on plate 18, figure 2a, b, the holotype, is fragmentary and was refigured from Boehm's plate 42, figure 2 (refigured here for reference in Fig. 16); it measures 165 mm with a phragmocone of 135 mm.

Both Indonesian and the Jara macroconchs (Figs. 5, 16) resemble somewhat the early-middle Early Callovian *Macrocephalites formosus* (Waagen) [M] (Fig. 11a, b) recorded from almost all domal outcrops in Kachchh (Jain & Pandey 2000; Krishna & Ojha 1996), Madagascar (Thierry 1978) and Europe (Krishna & Cariou 1990; Cariou et al. 1993). In Kachchh, its acme marks a Zone, the Formosus Zone (Figs 12, 13), which is delimited at the base by the occurrence of *Collotia oxytypha* Neumayr and at the



**Fig. 15:** Domal (from distal to proximal transect of the basin; Jara to Habo, respectively) distribution of macrocephalitid species within the Kachchh Basin. The ammonite fauna is not stratigraphically differentiated; the compressed species occur together with the inflated species, although their percentages may differ and they themselves differ laterally in abundance from one dome to the other.



Fig. 16: Macrocephalites keeuwensis Boehm [M]. holotype refigured from Boehm (1912: pl. 42, fig. 2). It measures 165 mm with a phragmocone of 135 mm (see text for explanation and Appendix for dimensions).

top by *Hecticoceras proximum* Elmi (Krishna & Ojha 1996; Jain & Pandey 2000) and hence, is correlated with the standard European Proximum subzone, Gracilis Zone (Fig. 13). The Formosus Zone is best developed at Jumara (Krishna & Westermann 1987) where the nominal macroconch species, has a long range, from the Earliest Callovian Madagascariensis Zone to the middle–Early Callovian Formosus Zone, itself (Figs 12, 13).

The macroconch of *Macrocephalites formosus* is a large form (maximum estimated shell diameter in excess of 410 mm; phragmocone measures about 194 mm; Fig. 11a, b) (Bhaumik et al. 1993), is involute (U/D = 0.07), moderately compressed (T/H = 0.90) with smooth flanks, at or just after the phragmocone (although in some specimens, smoothening starts much before the end of the phragmocone). Umbilicus is deep and cylindrical with a venter that is narrow and acutely rounded (Fig. 11b). The large size, smooth flanks, deep cylindrical umbilicus and the narrow acutely rounded venter distinguish M. formosus from the macroconch of M. keeuwensis Boehm (compare Figs 5, 16 with Fig. 11) or any other Indonesian forms (Westermann & Callomon 1988; Thierry 1978).

The other closely comparable form is Collignon's *M. formosus* (1958: pl. 13, fig. 66) from his Macrocephalites triangularis Zone, Andranomantsy III, Madagascar. However, it is much smoother with pronounced extraumbilication and body chamber modifications and a vertical umbilical wall even at 182 mm shell diameter.

The macroconch of European *Macrocephalites gracilis* (Spath) (Cariou & Hantzpergue 1997: pl. 19, fig. 7a, b) in its outer whorl smoothening is comparable, but *M. gracilis* is more evolute, with a subtriangular whorl section, an acutely rounded venter with high vertical umbilical walls. It is the index of the European late Early Callovian Gracilis Zone (Fig. 13).

Microconch [m]: Westermann & Callomon (1988) identified four *M. keeuwensis* morphs (two formae and two varieties):

- 1 Forma *typica*: It is moderately involute (U/D = 0.17-0.20), compressed (W/H  $\leq 0.8$ ); densely ribbed (P  $\geq 18-22$ ) and includes the Lectotype.
- Forma *flexuosa*: It is the most abundant form. It is involute (U/D ≤ 0.17), compressed (W/H ≤ 0.8) and densely ribbed (P ≥19). It includes *Dolikephalites flexuosus* Spath. Forma *flexuosa* is now





Fig. 17: Whorl section (T/H) versus Umbilical width (U/D) of *Macrocephalites keeuwensis* Boehm [M and m] discussed in this contribution (see Appendix for corresponding dimensions).

recorded here and described below.

 Var. A: It possesses weakly compressed whorls (W/H > 0.8) with subquadrate body chamber, medium umbilicus (U/D = 0.15–0.19) becoming moderately evolute at the body chamber. The ornamentation is coarse (P ≤ 19) and resembles *Macrocephalites subcompessus* (Waagen) from which it is distinguished by its sharp umbilical edge (as most Indonesian forms are).

Var. B: It is relatively evolute (U/D ≥ 0.20), particularly the body chamber (U/D = 0.25), and compressed (W/H ≤ 0.8). The ribs are strongly flexed and moderately dense (P = 17–20). It resembles *Subkossmatia* and *Eucycloceras*, from which its



Fig. 18: Proposed lineages. It is speculated, that *M. keeuwensis* Boehm [M and m] gave rise to two distinct lineages; one, that gave rise to the Early Callovian *M. lamellosus* (Sowerby) and that in turn to the late Early Callovian *Subkossmatia opis* Spath. On the other hand, to the early–middle Early Callovian *Macrocephalites formosus* (Waagen), which in turn gave rise to the late Early Callovian *M. semilaevis* Spath.

sharp umbilical edge and dense, sharp ribbing on the body chamber distinguish it. Var. B is recorded here and described below.

#### Macrocephalites keeuwensis Boehm var. B [m]

The specimen figured is in its morphological and dimensional characteristics (Figs 6a–c, 17) an exact match with Boehm's (1912: pl. 38, fig. 1) *Macrocephalites keeuwensis* Boehm var. B [m] from "Keeuw" (refigured here in Fig. 6e), as also the European Late Bathonian (Orbis Zone) *Macrocephalites* cf. / aff. *keeuwensis* Boehm (Dietl & Callomon 1988; refigured here in Fig. 6f).

Thierry's (1978: pl. 21, fig. 9a) *Macrocephalites macrocephalus keeuwensis* (= *Dolikephalites flexuo-sus* sensu Spath) is somewhat comparable but is much smaller (86 mm) and more densely ribbed (26 primaries and 62 secondaries per half whorl).

Jeannet's (1955: pl. 22, fig. 1) *Macrocephalites* (*Pleurocephalites*) *uhligi* Lemoine and his *Macrocephalites* (*Kamptokephalites*) *lamellosus* Sowerby (1955: pl. 26, fig. 3) from Herznach, Switzerland (Enodatum subzone; top of Calloviense Zone; late Early Callovian) are comparable forms but possess a straighter ribbing pattern with a distinctly rounded umbilical shoulder.

The Kachchh *Macrocephalites lamellosus* (Sowerby) [m] (Fig. 10) which occurs stratigraphically just above (Earliest Callovian) the Jara microconch (Latest Bathonian) (Fig. 13), is also a closely comparable form but lacks the distinct primaries that show a strong adoral concavity across the umbilical margin as also the characteristic sharp umbilical edge, besides the former's microconch is much smaller and inflated with a broadly rounded venter (Fig. 10c–d). The dimorphic pair for *M. lamellosus* (Sowerby) has now been recorded from Kachchh (Fig. 10). The

Macroconch comes from the late Early Callovian Semilaevis Zone (bed B36 from Jumara); it is complete with an attached operculum (Fig. 10a, b) and measures 150 mm with a phragmocone of 115 mm. The holotype, a microconch, is almost complete at 108 mm from Early Callovian sediments of Keera (refigured here in Fig. 10c, d).

The Jara specimen (Fig. 6a–d) in its evoluteness and to some extent, in its ribbing pattern, closely resembles the latest Early-earliest Middle Callovian Subkossmatia opis (see refigured holotype in Fig. 9b-d and another example refigured in Fig. 9a from Jana et al. 2005, his Eucycloceras opis var. opis [m] from Keera) of the Eucycloceratinae (see also Jain & Pandey 2000 for stratigraphic distribution) as recognized by Spath (1928: p. 202, 210, pl. 39, figs 2, 7) (Fig. 13). As also noted by Spath, in M. keeuwensis Boehm only the body chamber egresses, while in Subkossmatia egression begins at the intermediate whorls, e.g. the entire ultimate septate whorl egresses, as noted in the fully septate holotype of S. opis (Fig. 9a). Additionally, Subkossmatia differs in possessing increasingly and more strongly projected secondary ribs. However, Westermann & Callomon (1988) noted that M. keeuwensis Boehm var. B is often indistinguishable from the nucleus of Subkoss*matia*, unless the septal suture is preserved showing the convex or "raised" umbilical saddle envelope of the former. Besides these, there are no other comparable forms of *M. keeuwensis* Boehm var. B [m].

Macrocephalites keeuwensis Boehm aff. var. forma flexuosa [m]

Two fragmentary specimens were found from the top of the Sponge Bed (bed A8; see Fig. 7a–e) associated with *M*. cf. *mantataranus* Boehm (Jain & Desai 2014) (Fig. 8). Based on this association, their close resemblance to the ribbing pattern of *M. keeuwensis* Boehm var. forma *flexuosa* [m] (Fig. 7f) and the fact that there are no other comparable compressed and densely ribbed macrocephalitid forms at this level, their affinity to *M. keeuwensis* Boehm var. forma *flexuosa* [m] is suggested (Fig. 7f).

Another closely comparable form is *Dolikephalites flexuosus* Spath (1928: p. 210, pl. 32, fig. 4), which has already been considered to be a synonymy of *M. keeuwensis* Boehm [m] (Westermann & Callomon 1988). Additionally, no Kachchh macrocephalitid form possesses such a sharp umbilical edge as the specimens presented here; a characteristic feature of Indonesian forms (Westermann & Callomon 1988). The crowding of the ribs and their strong forward projection suggests maturity of these fragmentary specimens (Fig. 7a, b, d). The macroconch (Fig. 7g–i; refigured from Westermann & Callomon 1988: pl. 13, fig. 3a, b) shows the same ribbing pattern, sharp umbilical edge and inwardly projected umbilical walls.

#### 7. Conclusions

Co-occurrence of the dimorphic pairs of the Indonesian ammonites Macrocephalites keeuwensis Boehm [M & m] and M. keeuwensis var. aff. forma flexuosa Boehm [m] with M. cf. mantataranus Boehm [M] from Jara and Jumara Domes (Mainland Kachchh) enables to date not only the Indonesian *M. keeuwensis* Association but also the strata where the nominal species M. keeuwensis Boehm occurs in Kachchh, Madagascar and Europe as definitely Late Bathonian. The Late Bathonian age is additionally strengthened by reports from the Jumara Dome of characteristic Orbis-Discus Zone fossils of Oxycerites cf. orbis (Roy et al. 2007), O. cf. oppeli (Pandey et al. 2012) and Bullatimorphites cf. hannoveranus (Bardhan et al. 2002). Additionally, it is speculated, that *M. keeuwensis* Boehm [M & m] gave rise to two distinct lineages and the paper documents the separation of these two different lines (Fig. 18). One that gave rise to the Early Callovian M. lamellosus (Sowerby), and that in turn led to the late Early Callovian Subkossmatia opis Spath, as well as to the early-middle Early Callovian Macrocephalites formosus (Waagen), which in turn gave rise to the late Early Callovian M. semilaevis Spath (Fig. 18). The dimorphic pair of both *M. lamellosus* Sowerby and *M. for*mosus (Waagen) are recorded. All the above forms now occur in Kachchh, Western India.

#### Acknowledgements

The author is grateful to the late Drs. J.H. Callomon (United Kingdom) and G.E.G. Westermann (Canada), who provided literature and sample help from time to time. Special thanks to Dr. Eckhard Mönnig (Coburg, Germany) for identifying the Jara Macroconch of *Macrocephalites keeuwensis* Boehm and Dr. Volker Dietze (Stuttgart, Germany), who considerably improved the manuscript through constructive comments.

#### 8. References

- Bardhan S, Sardar SK, Jana SK. 2002. The Middle Jurassic *Kheraiceras* Spath 1924 from the Indian Subcontinent. In: H Summesberger, K Histon, A Daurer (Eds.), Cepahalopods Present and Past. Abhandlungen der Geologischen Bundesanstalt 57, 265–277.
- Bhaumik D, Datta K, Jana SK, Bardhan S. 1993. Taxonomy and intraspecific variation of *Macrocephalites formosus* from the Jurassic Chari Formation, Kutch, Western India. Journal of Geological Society of India 42, 163–179.
- Boehm G. 1908. Jura von Rotti, Timor, Babar und Buru. Neues Jahrbuch für Mineralogie, Geologie und Paläontologie 25, 324–343.
- Boehm G. 1912. Beiträge zur Geologie von Niederländisch-Indien. I. Abteilung: Die Südküsten der Sula-Inseln Taliabu und Mangoli. 4. Abschnitt; Unteres Callovium. Palaeontographia Suppl. IV, 124–179.
- Callomon JH. 1993. On *Perisphinctes congener* Waagen, 1875, and the age of the Patcham Limestone in the Middle Jurassic of Jumara, Kutch, India. Geologische Blätter von Nordost-Bayern 43, 227–246.

- Cariou E, Enay R. 1999. The Bathonian–Callovian ammonites from Thkkhola (Central Nepal): Biochronology and paleogeographical aspect. Geobios 32, 701–726.
- Cariou E, Hantzpergue P. 1997. Biostratigraphie du Jurassique ouest-européen et méditerranéen: zonations parallèles et distribution des invertébrés et microfossiles. Bulletin des Centres de Recherche Elf Exploration–Production, Mémoir 17, 440 p.
- Cariou E, Krishna J, Enay R. 1990. Dynamique biogéographique des ammonites téthysiennes au Callovien inférieur et corrélations stratigraphiques entre les marges indienne et européenne. Comptes Rendus de l'Académie de Sciences de Paris 311, 373–379.
- Collignon M. 1958. Atlas des fossiles caracteristiques de Madagascar. Tome 2 (Bathonien–Callovien). Service geologique, Tananarive p. 3.
- Dietl G, Callomon JH, 1988. Der Orbis–Oolith (Ober–Bathonium, Mittl. Jura) von Sengenthal/Opf., Frank. Alb, und seine Bedeutung fur die Korrelation und Gliederung der Orbis–Zone. Stuttgarter Beiträge zur Naturkunde, Serie B 142, 1–31.
- Dietl G. 1981. Über *Macrocephalites* (Ammonoidea) aus dem Aspidoides–Oolith und die Bathonium/Callovium–Grenzschichten der Zollernalb (SW–Deutschland). Stuttgarter Beiträge zur Naturkunde, Serie B 68, 1–15.
- Etheridge E. 1890. Our present knowledge of the palaeontology of New–Guinea. Records of the Geological Survey of New South Wales 1, 172–179.
- Jain S, Callomon JH, Pandey DK. 1996. On the earliest known occurrence of the Middle Jurassic ammonite genus *Reineckeia* in the Late Bathonian of Jumara, Kachchh, Western India. Paläontologische Zeitschrift 70, 129–143.
- Jain S, Desai BG. 2014. Biostratigraphic Implications of the Discovery of Late Bathonian Indonesian Ammonite *Macrocephalites* cf. *mantataranus* Boehm [M] from the core of Jara Dome, Kachchh, Western India. Journal of the Palaeontological Society of India 59, 1–11.
- Jain S, Pandey DK. 2000. Middle Jurassic Ammonite Biozonation in Kachchh, western India. Bulletin of the Indian Geologists Association 33, 1–12.
- Jain S. 1995. Biostratigraphical and Paleoecological Studies of Middle Jurassic (Upper Bathonian–Lower Callovian) Beds Exposed in Jumara Dome, Kachchh, Western India. Unpublished Thesis, 202 p.
- Jain S. 2002. Middle Jurassic Ammonite Biozonation in Western India: Global Implications, GSA Annual Meeting, Colorado. Paper No. 141–10, p. 34.
- Jain S. 2014. The Arkelli Chronozone: A new early Middle Bathonian standard ammonite zone for Kachchh, western India (southern Jurassic Tethys). Zitteliana A 54, 91–136.
- Jana SK, Bardhan S, Halder K. 2005. Eucycloceratin ammonites from the Callovian Chari Formation, Kutch, India. Palaeontology 48, 883–924.
- Jeannet A. 1955. Die Macrocephaliten des Callovien von Herznach (Aargau). Eclogae Geolgicae Helvetiae 47, 223–267.
- Krishna J, Cariou E. 1993. The Tethyan Macrocephalitinae: Evolutionary Environmental and Dispersal Strategies. Geobios 15, 217–226.
- Krishna J, Ojha JR. 1996. The Callovian Ammonoid Chronology in Kachchh, W. India. Geo Research Forum 1–2, 151–165.
- Krishna J, Ojha JR. 2000. The intrabasinal correlation in the Middle Jurassic Callovian Stage of Kachchh (Gujrat) and Ammoniods-Foraminifer integration. Geophytology 28, 101–120.
- Krishna J, Westermann, GEG. 1987. Faunal association of the Middle Jurassic ammonite genus *Macrocephalites* in Kach-

chh, Western India. Canadian Journal of Earth Sciences 24, 1570–1582.

- Mangold C, Rioult, M. 1997. Ammonites Bathonien. In: Cariou E, Hantzpergue P. (Coord.), Biostratigraphie du Jurassique Ouest–Européen et Méditerranéen. Bulletin des Centres de Recherche Elf Exploration-Production 17, 55–62.
- Mönnig E. 2014. The stratigraphy of the Bathonian–Callovian boundary (Middle Jurassic) in Northern Germany. Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen 274, 271–290.
- Mönnig K. 1995. Der Macrocephalen–Oolith von Hildesheim. Mitteilungen des Riemer-Museums Hildesheim, Neue Folge 5, 1–77.
- Mönnig K. 2015. The stratigraphy of the Bathonian–Callovian boundary (Middle Jurassic) in Northern Germany. Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen 274, 271–290.
- Pandey B, Krishna J, Pathak DB, Kumar A. 2012. Ammonoid Biostratigraphy of Bathonian Succession at Jumara, Kachchh, Western India. Journal Indian Geological Congress 4, 7–18.
- Prasad S. 1998. Ammonite Biozonation of the Middle–Late Jurassic Sediments with Special Reference to Keera and Jara, Kachchh District, Gujrat. Journal of Geological Society of India 52, 25–40.
- Rai J. 2003. Early Callovian nannofossils from Jara Dome, Kachchh, western India. Journal of the Geological Society of India 61, 283–294.
- Roy P, Bardhan S, Mitra A, Jana SK. 2007. New Bathonian (Middle Jurassic) ammonite assemblages from Kutch, India. Journal of Asian Earth Sciences 30, 629–651
- Sato T, Westermann GEG, Skwarko SK, Hasibaun F. 1978. Jurassic biostratigraphy of the Sula Islands, Indonesia. Bulletin of the Geological Survey of Indonesia 4, 1–28.
- Schlotheim EF von. 1813. Beiträge zur Naturgeschichte der Versteinerungen in geognostischer Hinsicht. Leonard's Taschenbuch für die gesammte Mineralogie 7, 3–134.
- Seyed–Emami K, Roufian A, Mönnig K. 2015. Macrocephalitinae (Ammonoidea, Middle Jurassic) from North and Central Iran. Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen 278, 257–279.
- Spath LF. 1927–33. Revision of the Jurassic Cephalopod fauna of Cutch. Memoirs of the Geological Survey of India. Palaeontologica Indica, New Series 9, 1–945.
- Thierry J, Cariou E, Elmi S, Mangold C, Marchand D, Rioult, M. 1997. Ammonites – Callovien. In: Cariou E, Hantzpergue P. (Coord.), Biostratigraphie du Jurassique Ouest–Européen et Méditerranéen. Bulletin des Centres de Recherche Elf Exploration-Production 17, 56–78.
- Thierry J. 1978. Le genre *Macrocephalites* au Callovien inferiur (Ammonites, Jurassique moyen). Systematique et evolution, biostratigraphie, biogeographie. Europe et domaine indo-malgeche. Mémoires Géologiques de l'Université de Dijon 4, 451.
- Westermann GEG, Callomon JH. 1988. The Macrocephalitinae and associated Bathonian and Early Callovian (Jurassic) ammonoids of the Sula Islands and Papua New Guinea. Palaeontographica Abt. A 203, 1–90.
- Westermann GEG, Sato T, Skwarko SK. 1979. Brief report on the Jurassic biostratigraphy of the Sula Islands, Indonesia. Newsletter on Stratigraphy 7, 96–101.
- Yin J. 2016. Bathonian–Callovian (Middle Jurassic) ammonites from northwestern Qiangtang Block, Tibet, and the revised age of the Suowa Formation. Proceedings of the Geologists' Association 127, 247–265.
- Zittel KA von. 1884. Handbuch der Paläontologie, 2 vols. München/ Leipzig, Oldenbourg, 893 p.

**Appendix:** Measurements of *Macrocephalites keeuwensis* Boehm [M & m] in mm. T = Whorl thickness; H = Whorl height; T/H = Whorl section; U/D = Umbilical width.

Species	Sp. no.	Measured at	D	Н	Т	U	T/H	U/D
	J2023W	Body chamber	115.3	52.1	38	14.9	0.73	0.13
	Ja/SJ/1999/	Body chamber	172	75.6	59.6	23.7	0.79	0.14
	Core/2	Phragmocone	132	46.1	43	15	0.93	0.11
	Boehm 1912:	Body chamber	163	78	61	21	0.78	0.13
Boehm [M]	pl. 42, fig. 2 (holotype)	Phragmocone	135	70	56	20	0.8	0.15
	Boehm 1912: pl. 44, fig. 1a, b	Body chamber	105	53	50	14	0.94	0.13
	Boehm 1912: pl.	Body chamber	102	54.5	47.1	13.5	0.86	0.13
	36, fig. 1	Phragmocone	82	45	42	13	0.93	0.16
	Ja/SJ/1999/	Phragmocone	80	38.5	30.5	13	0.79	0.16
var. B [m]	Core/1	Body chamber	107	44.2	34.7	21.4	0.79	0.20

# **ZOBODAT - www.zobodat.at**

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Zitteliana Serie A+B gemeinsam

Jahr/Year: 2019

Band/Volume: 93

Autor(en)/Author(s): Jain Sreepat

Artikel/Article: <u>On the occurrence of the Indonesian ammonite Macrocephalites</u> keeuwensis Boehm [M & m] from Kachchh (Western India) 3-24