


Zitteliana

An International Journal
of Palaeontology and Geobiology

Series A/Reihe A
Mitteilungen der Bayerischen Staatssammlung
für Paläontologie und Geologie

45



München 2005

Zitteliana

An International Journal of Palaeontology and Geobiology

Series A/Reihe A

Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Geologie

45

CONTENTS/INHALT

| | |
|---|-----|
| DHIRENDRA K. PANDEY & FRANZ T. FÜRSICH Jurassic corals from southern Tunisia | 3 |
| THORSTEN KOWALKE Mollusca in marginal marine and inland saline aquatic ecosystems – examples of Cretaceous to extant evolutionary dynamics | 35 |
| JOACHIM GRÜNDEL Gastropoden aus dem oberen Callovium (Lamberti- Zone) der Tongrube Dubki bei Saratov, Russische Plattform | 65 |
| SIMON SCHNEIDER, WOLFGANG WITT & ERDİNÇ YIGİTBAŞ Ostracods and bivalves from an Upper Pleistocene (Tyrrhenian) marine terrace near Altınova (İzmit Province, Turkey) | 87 |
| RENATE MATZKE-KARASZ & WOLFGANG WITT Ostracods of the Paratethyan Neogene Kılıç and Yalakdere Formations near Yalova (İzmit Province, Turkey) | 115 |
| JÜRGEN KRIWET A comprehensive study of the skull and dentition of pycnodont fishes (Neopterygii, Pycnodontiformes) | 135 |
| JEAN GAUDANT & BETTINA REICHENBACHER <i>Hemitrichas stapfi</i> n. sp. (Teleostei, Atherinidae) with otoliths <i>in situ</i> from the late Oligocene of the Mainz Basin | 189 |
| ALFRED SELMEIER <i>Capparidoxylon holleisii</i> nov. spec., a silicified <i>Capparis</i> (Capparaceae) wood with insect coprolites from the Neogene of southern Germany | 199 |
| INKEN JULIANE MUELLER-TÖWE <i>Short Communication</i> : Phylogenetic relationships of the Thalattosuchia | 211 |
| Instructions for authors Hinweise für Autoren | 215 |

Editors-in-Chief/Herausgeber: Reinhold Leinfelder, Michael Krings
Production and Layout/Bildbearbeitung und Layout: Martine Focke, Lydia Geißler

Editorial Board

A. Altenbach, München
B.J. Axsmith, Mobile, AL
F.T. Fürsich, Würzburg
K. Heißig, München
H. Kerp, Münster
J. Kriwet, Berlin
J.H. Lipps, Berkeley, CA
T. Litt, Bonn
O.W.M. Rauhut, München
B. Reichenbacher, München
J.W. Schopf, Los Angeles, CA
G. Schweigert, Stuttgart
F. Steininger, Frankfurt a.M.
W. Werner, München

Bayerische Staatssammlung für Paläontologie und Geologie
Richard-Wagner-Str. 10, D-80333 München, Deutschland
<http://www.palaeo.de/zitteliana>
email: zitteliana@lrz.uni-muenchen.de

Für den Inhalt der Arbeiten sind die Autoren allein verantwortlich.
Authors are solely responsible for the contents of their articles.

Copyright © 2005 Bayerische Staatssammlung für Paläontologie und Geologie, München

Die in der Zitteliana veröffentlichten Arbeiten sind urheberrechtlich geschützt.
Nachdruck, Vervielfältigungen auf photomechanischem, elektronischem oder anderem Wege
sowie die Anfertigung von Übersetzungen oder die Nutzung in Vorträgen, für Funk und Fernsehen
oder im Internet bleiben – auch auszugsweise – vorbehalten und bedürfen der schriftlichen Genehmigung
durch die Bayerische Staatssammlung für Paläontologie und Geologie, München.

ISSN 1612-412X

Druck: Gebr. Geiselberger GmbH, Altötting

Cover illustration: Ostracod *Callistocythere intricatoides* (RUGGIERI, 1953) from the Thyrrenian of Altinova (Turkey). Left: Right valve, external view, BSPG 1980 X 1313 (length 0.640 mm). Right: Left valve, external view, BSPG 1980 X 1314 (length 0.646 mm). SEM Photograph: R. MATZKE-KARASZ (LMU München, Department für Geo- und Umweltwissenschaften, Sektion Paläontologie)

Umschlagbild: Ostrakode *Callistocythere intricatoides* (RUGGIERI, 1953) aus dem Thyrrenium von Altinova (Türkei). Links: Rechte Klappe, Außenansicht, BSPG 1980 X 1313 (Länge 0,640 mm). Rechts: Linke Klappe, Außenansicht, BSPG 1980 X 1314 (Länge 0,646 mm). REM-Foto: R. MATZKE-KARASZ (LMU München, Department für Geo- und Umweltwissenschaften, Sektion Paläontologie)

Capparidoxylon holleisii nov. spec., a silicified *Capparis* (Capparaceae) wood with insect coprolites from the Neogene of southern Germany

By
Alfred Selmeier*

Department für Geo- und Umweltwissenschaften – Sektion Paläontologie, Ludwig-Maximilians-Universität
München, Richard-Wagner-Straße 10, 80333 Munich, Germany

Manuscript received July 28, 2005; revision accepted September 10, 2005.

Abstract

A silicified dicotyledonous wood from Attenfeld, southern Franconian Alb, Germany, is assignable to the form genus *Capparidoxylon* SCHENK, 1883 based on anatomical features that are identical to those seen in the wood of extant members in the genus *Capparis*. The new species *Capparidoxylon holleisii* nov. spec. is introduced for the fossil. The genus *Capparis* is today comprised of some 250 species, and widely distributed in dry environments in the Mediterranean, the Palaeotropis, and in South America. A special feature of the fossil is insect feeding channels filled with coprolites. *Capparidoxylon holleisii* nov. spec. represents the first record of a silicified Capparaceae wood from the Neogene of Europe.

Key words: Deciduous tree, Capparaceae, South Germany, Neogene, Wood anatomy, Insect coprolites

Kurzfassung

Ein verkieseltes Laubholz aus Attenfeld, südliche Franconialb, Süddeutschland, läßt sich der Formgattung *Capparidoxylon* SCHENK, 1883 zuordnen. Für dieses Holz wird die neue Art *Capparidoxylon holleisii* nov. spec. eingeführt. Die Anatomie des Holzes wird beschrieben und durch Abbildungen dokumentiert. Die Feinstruktur ist weitgehend ähnlich und vergleichbar mit dem Xylem der artenreichen Gattung *Capparis* (250), derzeit weit verbreitet im Mittelmeerraum, der Paläotropis und in Südamerika. Eine Besonderheit des Fossils sind die erhalten gebliebenen Bohr- und Fraßgänge mit Coprolithen von holzbewohnenden Insekten. Die vorliegende Arbeit beschreibt erstmals einen verkieselten Holzrest der Familie Capparaceae aus neogenen Schichten Europas.

Schlüsselwörter: Holz-anatomie, Laubbaum, Capparaceae, Insektenbefall, Süddeutschland, Neogen


1. Introduction

With the Alps emerging, their northern foreland developed into an elongate depression. This depression, the Molasse basin, was subsequently filled with mountain debris. With the onset of the Upper Freshwater Molasse period, continental conditions prevailed in the northern alpine foreland. Giant alluvial fans transported alpine debris into a westward oriented river system (GLASER et al. 2004). Gravel, sands, and fine sediments were deposited and formed the “Tertiärhügelland”, which is located south of the river Danube, between the cities Augsburg and Passau. The Early and Middle Miocene sediments of the North Alpine Foreland Molasse Basin continue to yield one of the world’s richest regional records of silicified wood (SELMEIER 1998b). The number of specimens gathered to date ranges well over 10.000, of which some 1.000 specimens have been scientifically analyzed. These specimens come from 61 stratigraphically well dated localities by using the Coexistence Approach, and demonstrate the potential of fossil silicified wood in palaeobotanical and palaeoenvironmental studies (BÖHME et al., in preparation).

Nevertheless, wood of Capparaceae (formerly Cappari-daceae) has not been recorded to date. The specimen presented here was collected by P. HOLLEIS from a site west of Attenfeld, southern Franconian Alb. The town of Attenfeld is situated between the rivers Altmühl (north) and Danube (Textfig. 1). The fossils from the Franconian Alb predominantly occur in sandy or loamy sediments, usually close to the surface. They can easily be picked from farmland or at waysites.

There exist only 2 silicified woods today that have been assigned to the family Capparaceae, i.e. *Capparidoxylon Geinitzi* SCHENK (1883) from Egypt (Kairo) and *Forchhammerioxylon scleroticum* KRUSE (1954) from Wyoming (U.S.A.). Even the extensive list of silicified woods recorded for the Middle Eocene Clarno Nut Beds flora (145 genera) does not contain the family Capparaceae (cf. WHEELER & MANCHESTER 2002). The specimen of Attenfeld is thus the third fossil record worldwide

* E-mail: a.selmeier@lrz.uni-muenchen.de



Textfigure 1: Geographic position of Attenfeld and other Miocene localities of the southern Franconian Alb that have yielded diverse silicified wood assemblages.

for the Capparaceae, and the first record for this family from the Neogen of Europe. The specimen from Attenfeld displays insect boreholes and channels filled with coprolites.

In this paper, the microscopic anatomy of *Capparidoxylon holleisii* nov. sp. is described and compared with that seen in the wood of extant representatives of the Capparaceae. The diagnostic value of various anatomical features is discussed.

2. Material and Methods

The specimen is preserved as silicified cellular permineralization. All slides were prepared using standard thin-section techniques. A diamond saw was used to cut thick sections (wafers) of transverse, longitudinal tangential, and radial surfaces. These sections were ground down with a Logitech Lapp & Polish LP 50 (Edinburgh, Scotland) grinding machine until they were thin enough to permit analysis of the anatomical details by transmitted light microscopy (cf. HASS & ROWE 1999).

The description of the wood anatomical details generally follows the list of microscopic features for hardwood identification compiled by the IAWA COMMITTEE (1989), counting each vessel separately (WHEELER 1986), and the recommendations of WHEELER & BAAS (1998). Photo documentation was accomplished with a Zeiß Ultraphot analog camera and a

digital Leica Q 550 IW. Images were processed, and plates and textfigures constructed using Adobe Photoshop 7.0.

For authors of plant names and taxonomy of plant families, genera, and species, four publications were available, i.e. BRUMMIT & POWELL (1992), KUBITZKI & BAYER (2003), MABBERLEY (1997), and SCHUBERT & WAGNER (1984).

3. Systematic Palaeontology

Order Capparales (formerly Rhoeadales)

Family Capparaceae (formerly Capparidaceae)


Organ genus *Capparidoxylon* SCHENK, 1883 (without diagnosis)

Type species: *Capparidoxylon Geinitzi* SCHENK, 1883 (without diagnosis)

Capparidoxylon holleisii nov. sp.

Pl. 1, Figs 1-4; Pl. 2, Figs 1-4; Textfigs 2-6.

Material: Silicified wood, specimen measures 13.3 x 6,8 x 4,6 cm, weight 456 g, surface somewhat smooth, light brown, inside darker.



Textfigure 2: *Capparidoxylon holleisii* nov. sp. Cross section [holotype]. Selection of typical vessel forms, radial multiples of 2-10, occasionally forming small clusters, and vessels intergrading with tracheids; BSPG 2004 XV 1 and HOLLEIS IX 309; scale bar = 200 µm.




Plate 1: *Capparidoxylon bolleisii* nov. sp. from the Neogene of the southern Franconian Alb, Bavaria (Germany); BSPG 2004 XV 1 (Fig. 1) und HOLLEIS IX 309 (Figs 2-4).

Fig. 1: Cross section. Indistinct growth ring boundary, vessels diffuse-porous; scale bar = 1000 μ m, x 15.

Fig. 2: Cross section. Vessels in radial multiples of up to five. Ground tissue and dark rays visible; scale bar = 300 μ m, x 70.

Fig. 3: Cross section. Ground tissue, vessels, and ten boreholes filled with coprolites recognizable; scale bar = 500 μ m, x 55.

Fig. 4: Cross section. Ground tissue, vessels, and a long radially-orientated borehole filled with coprolites; scale bar = 500 μ m, x 55.

Holotype: BSPG 2004 XV 1, hand specimens and seven thin sections, reposit in the Bayerische Staatssammlung für Paläontologie und Geologie, Munich, Germany. An additional cross section (IX 309) and a few cutting fragments of the holotype specimen are kept in the Fossil Wood Collection of the "Historischer Verein und Museum Neuburg" (cf. SELMEIER 2001).

Diagnosis: Growth rings indistinct. Vessels diffuse-porous, medium-sized (50-100 µm), solitary (not more than 10%), radial multiples up to 10, some in clusters, vessels sometimes intergrading with vascular tracheids, perforation simple, vessel element members relatively short, many vessel lumina filled with yellow or dark deposits. Fibres non-septate, medium thick-walled. Axial parenchyma scanty, paratracheal, and/or sparsely vasicentric. Rays 1-2-(3)-seriate, faintly heterocellular (KRIBS 1935; type III), sheat cells occasionally occur, 4-26 cells high, 6-12 rays per mm tangential.

Etymology: The specific epithet refers to P. HOLLEIS who collected the fossil presented here and hundreds of other valuable silicified woods from the southern Franconian Alb.


Locality: Attenfeld, Bavaria, map TK 1:25.000, No. 7233, Neuburg a. d. Donau, southern Franconian Alb, Germany (Textfig. 1).

Age and horizon: Neogene sediments of the Alpine foredeep, southern Franconian Alb. The Lower part of the Upper Freshwater Molasse (OSM, Obere Süßwasser Molasse) is correlated with the lower Karpatien-Badenien based on mammal fossils (uppermost MN4-MN5), Late Karpatian, 17.0-16.3 Ma. (DOPPLER et al. 1996, 2002).

| Length and width, µm | Length and width, µm |
|----------------------|----------------------|
| 221 : 117 | 559 : 104 |
| 273 : 208 | 676 : 78 |
| 274 : 234 | 819 : 468 |
| 312 : 234 | 1326 : 143 |
| 442 : 182 | 1664 : (156-364) |

Table 1: Size of 10 insect borings filled with coprolites, measured from cross sections of *Capparidoxylon holleisii*.

Description: Secondary xylem of dicotyledonous wood without bark or pith. Microscopic structure of radial section poorly preserved. Growth rings faint and indistinct (Pl. 1, Fig. 1). Vessels randomly arranged, a specific pattern is not recognizable (Pl. 1, Fig. 1-2), diffuse-porous, and either solitary (not more than 10%), in radial multiples of 2-10 or forming



Textfigure 3: *Capparidoxylon holleisii* nov. sp. Longitudinal tangential section. Rays 1-3-seriate, faintly heterocellular, tendency to sheat cells; BSPG 2004 XV 1; scale bar = 200 µm.

small clusters (rare); vessels partly intergrading with vascular tracheids (Textfig. 2), solitary vessels tangential diameter up to 95 (radial 52-122) µm, tangential diameter of radial multiples of two 55-129 (mean 95) µm, radial 80-259 (mean 185) µm; individual vessels of multiples flattened; multiples radially up to 683 µm; perforations simple with transverse to oblique end walls (Textfig. 4), vessel member length short, 165-427 (mean 283) µm, intervessel pits crowded and alternate; very small to minute, difficult to observe due to meagre preservation; in cross- and longitudinal sections, many vessel lumina filled with yellowish deposits (Pl. 2, Figs 1-2; Textfig. 4), they often appear to accumulate at the end of vessel elements (longitudinal sections). Fibres non-septate (Pl. 2, Fig. 4), medium thick-walled, polygonal in cross section, more or less quadratic or round, tangential diameter 15-22 µm, radial 11-23 µm. 4-(6) rows of fibres between two rays, locally only 1-2 rows. The axial parenchyma is paratracheal and usually sparsely vasicentric. Rays 1-2(3) seriate, somewhat heterocellular (KRIBS 1935; type III), 3(96 µm)-26(738 µm) cells high (Pl. 2, Figs 3-4; Textfig. 3); uniseriate rays, e.g., with 12 cells (365 µm), uniseriate rays partly 2-seriate, 143-857 µm high; rays composed of procumbent, sometimes square and upright cells, sheat cells occasionally present, vertically up to 62 µm; 6-12 rays per mm tangential.





Plate 2: *Capparidoxylon holleisii* nov. sp. from the Neogene of the southern Franconian Alb, Bavaria (Germany); BSPG 2004 XV 1.


Fig. 1: Cross section. Vessels diffuse-porous, vessel lumina partly filled with yellow deposits, solitary vessels rare; scale bar = 500 µm, x 40.

Fig. 2: Longitudinal tangential section. Vessel members filled with brown-yellow deposits; scale bar = 300 µm, x 60.


Fig. 3: Longitudinal tangential section. Rays 1-3-seriate, faintly heterocellular; scale bar = 200 µm, x 125.

Fig. 4: Longitudinal tangential section. Fibres non-septate, rays faintly heterocellular, 1-3-seriate; scale bar = 200 µm, x 125.





Textfigure 4: *Capparidoxylon holleisii* nov. sp. Longitudinal tangential section. Short vessel members with simple perforation plates; BSPG 2004 XV 1; scale bar = 300 μm , x 60.



Textfigure 5: *Capparidoxylon holleisii* nov. sp. Longitudinal tangential section. Vertical insect boring filled with oval to round coprolites, presumably from *Anobium* sp.; BSPG 2004 XV 1; scale bar = 200 μm , x 120.

Crystals and storied structure not observed.

Coprolites: Numerous feeding channels occur that are particularly conspicuous in cross sections (e.g., Pl. 1, Figs 3-4; Textfig. 5). These are filled with coprolites; small coprolites are round and 27-55 μm in diameter, the larger ones oval and measure 75-112 (mean 89) \times 62-70 (mean 65) μm . The radial length of the borings varies from 0.2 to 1.6 mm (Tab. 1). *Capparidoxylon holleisii* represents only the second fossil from the southern Franconian Alb that displays insect damage. The perpetrators of these feeding channels probably were *Anobium* sp. (MOMBACHER 2003). A first silicified wood with feeding channels filled with coprolites assignable to *Anobium* sp. has been reported from Egweil (SELMEIER 1984b), a site located a few kilometers east of Attenfeld (Textfig. 1).

Synopsis of microscopic features:

- a) IAWA Committee (1989): 2, 5, 10, (11), 13, 22, (24/25), (41/42), 46, 51, 58, 60, 75, 97, 100, 105.
- b) Fossil Wood Datasheet for Species (WHEELER 1991a, b): 42, 45, 50, 51, 54, 58, 62, 65, 71, 83, 86, 87, 95, 106, 110, 113.

4. Discussion

4.1 Comments on Anatomical Features


Vessels – Distinct vessel diameter classes are a typical feature for 12 species of the Saharan and Sahelian Capparaceae (NEU-

MANN et al. 2001). This feature is highly diagnostic, but may be difficult to observe since the smaller vessels cannot be separated with confidence from vasicentric tracheids in cross sections (Textfig. 2). Quantitative features such as vessel diameter and number of vessels per mm^2 are highly variable.

Deposits – In woody plants, various deposits commonly occur and are irregularly distributed. Their formation is often related to processes of heartwood formation. In the fossil presented here, conspicuous yellow deposits, probably consisting of mustard-colored oil glycosides (cf. ROTH et al. 1994), are clearly visible in cross- and longitudinal sections (Pl. 2, Figs 1-2; Textfig. 4). Gums and other deposits include a wide variety of chemical compounds (HILLIS 1987).

Tracheids – 82% of the Saharan extant woody plants possess vascular/vasicentric tracheids (NEUMANN et al. 2001). These tracheids are effective as a subsidiary system for water conduction. Within the Sahelian woody plants that display vascular/vasicentric tracheids, the family Capparaceae is well represented with 11 species (NEUMANN et al. 2001). Unfortunately, this important feature is often difficult to observe without maceration.

Rays – Wood samples of a single plant species often display considerable size differences with regard to the width of the multiseriate rays. The fossil presented here, displaying mixed procumbent, square, and upright cells, is presumably part of a twig or young branch, but not a mature wood. The young tissues usually display a higher amount of square and upright cells. The ray height is also a highly variable feature.



Textfigure 6: *Capparidoxylon holleisii* nov. sp. (left, BSPG 2004 XV 1), and *Capparis speciosa* (right, RAKF No.3223). Note the similarities in cellular composition of the rays; scale bars = 200 µm, x 115.

Crystal-containing ray cells – In heterocellular rays with mixed procumbent and square or upright cells, crystals are usually present in both cell types. Crystals have not been observed from the fossil. However, MILLER (1978) found that sectioning and mounting techniques may remove some of the crystals.

4.2 Affinities

Capparidoxylon holleisii is characterized by indistinct growth rings, solitary vessels with approximately 50 to 100 µm tangential diameter, radial multiples of up to 10, the occasional occurrence of vessel clusters, vessels that sometimes intergrade with vascular tracheids, short vessel members, simple perforations, conspicuous yellow deposits, non-septate fibres, scanty paratracheal axial parenchyma, rays 1-2-(3)seriate, faintly heterocellular rays, and the occasional occurrence of sheath cells. This complement of wood anatomical in nearly all microscopic details clearly resembles that seen in woods members in the extant genus *Capparis*. A detailed summary of the similarities between the fossil *C. holleisii* and selected extant species of *Capparis* is provided in Tabs 2-4.

4.2.1 Comparison to Extant Wood

A detailed microscopic examination of the wood structure of selected extant Capparaceae was carried out to specify

the nearest living equivalent of the fossil. The Xylothech of the Wood Research Institute, Technical University Munich, Germany (STERN 1988) holds material of the following genera: *Boscia* sp. (LAM.), *Buchholzia coriacea* (ENGL.) A.CHEV., *Capparis cynophallophora* L., *C. speciosa* GRISEB., *C. spinosa* L., *C. tomentosa* LAM., *Crataeva adansonii* DC., *C. tapia* L. and *Maerua* sp. FORSSK. This material was included in the comparative analysis. *Capparis speciosa* GRISEB. (RAKF No.3223) is very similar to the present fossil, especially with regard to the cellular composition of the rays (Textfig. 6). Moreover, descriptions and microphotographs (GREGORY 1994) of the wood structure of extant Capparaceae in the literature were considered; these include data on *Capparis maroniensis* R. BEN. (DÉTIENNE & JACQUET 1983); *C. cartilaginea* DECNE., *C. decidua* (FORSSK.) EDGEW., *C. ovata* DESF., *C. spinosa* L., *Cleome droserifolia* (FORSSK.) DEL., *Maerua crassifolia* FORSSK. (FAHN et al. 1996); *C. decidua* (FORSSK.) EDGEW. (JAGIELLA & KÜRSCHNER 1987); *C. acuminata* WILLD., *C. micracantha* D.C., *C. subacuta* MIQ. (JANSSONIUS 1952); *C. decidua* (FORSSK.) EDGEW., *C. spinosa* L., *C. tomentosa* LAM., *Crataeva religiosa* FORST. (NEUMANN et al. 2001); *C. decidua* (FORSSK.) EDGEW. (NILOFARI 1961); *C. spinosa* L. (SCHWEINGRUBER 1991), and other species in the genus *Capparis* (cf. BOUREAU 1957; CARLQUIST 2001; ILIC 1991; METCALFE & CHALK 1950).

Only very few anatomical features distinguish *Capparidoxylon holleisii* from the general characteristics of the wood seen in the extant genus *Capparis*; no extant taxon displays exactly the same complement of wood anatomical features

| Microscopic features | Extant <i>Capparis</i> spp. | <i>Capparidoxylon holleisii</i> |
|--|---|--|
| Growth rings | indistinct | indistinct, faint to fairly distinct |
| Vessels | | |
| Arrangement | diffuse | diffuse |
| Tangential diameter | 53-127 (95) μm | 40-135 μm , up to 150 μm |
| Perforation plates | simple | simple |
| Radial multiples | 2-10 | up to 8(10) |
| Clusters, sometimes intergrading with vascular tracheids | present | present |
| Intervessel pits | alternate, weakly vested, 4-6 μm | alternate, apertures slit-like |
| Vessel element length | 160-200 μm | mean 283 μm |
| Vessels per square mm | 4-10 | 4-8 |
| Solitary vessels per mm | ca. 10% | 10% or less, up to 30% |
| Yellow and/or solid deposits | present in <i>C. decidua</i> | present |
| Fibres | | |
| Septate or non-septate | non-septate | non-septate |
| Pits | simple pits | not observed |
| Axial parenchyma | | |
| Vasicentric | scanty, paratracheal to vasicentric | scanty, paratracheal |
| Fusiform or in 2-3-celled strands | present | not observed |
| Rays | | |
| Width | 1-2-4 (6) seriate | 1-2 (3) seriate |
| Cellular composition | homo- and weakly heterocellular | homo- and weakly heterocellular |
| Sheath cells | occasionally present | present, rare |
| Height | up to 29 (50) cells | up to 26 cells |
| Number per mm tangential | (3)-5-9 | 6-12 |
| Crystals | not observed, probably water-soluble | not observed |

Table 2: Comparison of wood anatomical features of selected extant species of *Capparis* (i.e. *C. cartilaginea*, *C. decidua*, *C. ovata*, *C. spinosa*) and *Capparidoxylon holleisii*.

as that recorded for the fossil. For example, one difference in minute anatomy concerns the absence of two clearly distinct vessel diameter classes in cross section (Tab. 4) in extant *Capparis*. As stated above, the fossil presumably represents part of a twig or branch. It is established knowledge that the wood of branches and twigs may differ in anatomy from the wood of mature trunks. Trunk wood, however, usually forms the base for wood anatomical descriptions of extant plants. Thus, taking the wealth of striking similarities (Tabs 2-4) into account that exist between *Capparidoxylon holleisii* and extant

Capparaceae, the fossil can be assigned to the family *Capparaceae* with confidence.

4.2.2 Fossil *Capparaceae* Wood

The fossil record of *Capparaceae* wood is meagre (ANDREWS 1955). I am aware of only two descriptions of silicified wood specimens that have been assigned to *Capparaceae*, i.e. *Capparidoxylon Geinitzi* SCHENK, 1883 from the Petrified Forest near Cairo in Egypt and *Forchhammerioxylon scleroticum* KRUSE,

| | Number of vessels per multiples | % of solitary vessels | Vessel member length, mean μm | Ray width, No. of cells | Maximum ray height, No. of cells |
|---------------------------------|---------------------------------|-----------------------|--|-------------------------|----------------------------------|
| <i>C. cartilaginea</i> | 2-8(12) | 10 or less | 160 | 1-4 | 30 |
| <i>C. decidua</i> | 2-3(5) | 35 | 180 | 1-3(4) | 40 |
| <i>C. ovata</i> | 2-5(6) | 25 | 200 | (1)3-6 | 50 |
| <i>C. spinosa</i> | 2-6(8) | 20 | 170 | 1-4(5) | 30 |
| <i>Maerua crassifolia</i> | 2-8 | 40 | 140 | 1-3 | 45 |
| <i>Capparidoxylon holleisii</i> | 2-7(10) | 10 or less | 283 | 1-2(3) | 26 |

Table 3: Quantitative comparison of selected wood anatomical features of *Capparidoxylon holleisii* with those seen in selected extant species of *Capparis* and *Maerua* (according to FAHN et al. 1986).

| Species | Vessels | Deposits in vessel lumina | Vascular tracheids | Cellular composition of rays | Perforation plates | Reference |
|--|--|----------------------------------|--|--|--------------------|---------------------------|
| <i>Capparis decidua</i> | two distinct vessel diameter classes | yellow gum-like deposits | commonly present | procumbent, square and upright | simple | NEUMANN et al. 2001 |
| <i>Crateva religiosa</i> | medium sized | – | commonly present | procumbent, square and upright (in branchwood) | simple | Neumann et al 2001 |
| <i>Capparis decidua</i> | tending to be of two classes | gummy contents in many vessels | vessels typically intergrading with vascular tracheids | somewhat heterocellular | simple | FAHN et al 1996 |
| <i>Capparis decidua</i> | diffuse-porous, small, pore groups irregular distributed | mustard-oil glucosides | paratracheal-vasicentric parenchyma | homo- and heterocellular (KRIBS Typ III) | simple | JAGIELLA & KÜRSCHNER 1987 |
| <i>Capparis spp.</i> | – | often filled with solid deposits | – | – | simple | METCALFE & CHALK 1951 |
| <i>Capparidoxylon holleisii</i> nov. sp. | medium sized, two distinct classes not observed | yellow deposits in many vessels | vessels partly intergrading with vascular tracheids | procumbent, square, upright, sheat cells | simple | this paper |

Table 4: Wood anatomical similarities between *Capparidoxylon holleisii* and selected extant species of *Capparis*

| Fossil name | Affinities | Locality and age | Author |
|---------------------------------------|---|----------------------|--------------|
| <i>Capparidoxylon Geinitzi</i> | <i>Capparis</i> (Tourn.) L. | Egypt, Oligocene | SCHENK, 1883 |
| <i>Forchhammerioxylon scleroticum</i> | <i>Forchhammeria</i> , <i>Cadaba</i> , <i>Maerua</i> | Wyoming, USA, Eocene | KRUSE, 1954 |
| <i>Capparidoxylon holleisii</i> | <i>Capparis</i> spp. | Germany, Miocene | this paper |

Table 5: Summary of fossil Capparaceae wood described since 1883.

| Family | Genus | References |
|---------------------|--------------------------------------|---|
| Anacardiaceae | <i>Pistacia</i> | SELMEIER 2000a, 2000b; GOTTWALD 2004 |
| Capparaceae | <i>Capparis</i> | this paper |
| Dipterocarpaceae | <i>Shorea</i> | SELMEIER 1998a |
| Fagaceae | <i>Castanopsis</i> | SELMEIER 1970; GOTTWALD 2002, 2004 |
| Icacinaceae | <i>Cantleya</i> | GOTTWALD 2002, 2004 |
| Juglandaceae | <i>Carya</i> , <i>Juglans</i> | SELMEIER 1986; GOTTWALD 2004 |
| Lauraceae | <i>Laurus</i> , <i>Cinnamomum</i> , | SELMEIER 2003; GOTTWALD 2004 |
| Leg.-Mimosoideae | <i>Acacia</i> , <i>Dichrostachys</i> | GOTTWALD 2002, 2004 |
| Leg.-Papilionoideae | <i>Robinia</i> | GOTTWALD 2004 |
| Meliaceae | <i>Cedrela</i> | GOTTWALD 2002, 2004; SELMEIER 2003a, 2004 |
| Rosaceae | <i>Crataegus</i> | SELMEIER 1984a; GOTTWALD 2004 |
| Rosaceae | <i>Prunus</i> | SELMEIER 1984a |
| Sapotaceae | <i>Bumelia</i> | SELMEIER 1991; GOTTWALD 2004 |
| Ulmaceae | <i>Celtis</i> | SELMEIER 1989; GOTTWALD 2004 |
| Cupressaceae | <i>Cupressus</i> | GOTTWALD 2002 |
| Podocarpaceae | <i>Podocarpus</i> | GOTTWALD 2004 |
| Taxodiaceae | <i>Taxodium</i> | SELMEIER 2003b; GOTTWALD 2004 |
| Taxodiaceae | <i>Glyptostrobus</i> | GOTTWALD 2004 |
| Tempskyaceae | <i>Tempskya</i> | HOLLEIS & GREGOR 1986 |

Table 6: Silicified wood from the locality Attenfeld in the southern Franconian Alb: summary of genera described to date.

1954 from the Eden Valley in Wyoming (U.S.A.). However, the specimen from Egypt is not included in the Fossil Wood Database/Database References (WHEELER 1991a, 1991b). A second *Capparidoxylon* specimen from the Petrified Forest near Cairo is 8 cm long and 5 cm in perimeter, and was collected by SCHLIEFFEN. According to SCHENK (1883), this second specimen is very similar in anatomy to *C. Geinitzi*.

4.3 Silicified Wood from Attenfeld

Some 1.200 specimens of permineralized wood have been collected from farmland and waysides by P. HOLLEIS during the mid 1970s (HOLLEIS 1992). Twenty-two genera from this locality have to date been identified by xylem-microscopic methods (Tab. 6).

5. Concluding Remarks

Several localities in the southern Franconian Alb continue to yield well preserved Miocene silicified wood. Of these localities, Attenfeld produced the most diverse assemblage of wood ever recorded from a single locality in the North Alpine Molasse Foreland, and twenty two different plant genera have been identified based on silicified wood. The extant equivalents of these woody plants from the southern Franconian Alb generally thrive in temperate, subtropical, and tropical climates; they occur today in the Mediterranean to Minor Asia and to parts of South Asia. The fossil wood from Attenfeld must have been reworked and transported to southern places of the Franconian Alb by rivers based on the fact that fossilized roots have never been found to date in Miocene sediments in southern Franconian Alb.

Capparidoxylon holleisii represents the first record to date of silicified Capparaceae wood from the Neogene of Europe. The fossil displays indistinct growth rings, which suggests that relatively stable climate conditions existed during this time. Variability of the ecological conditions, including seasonal drought, soil composition, light regimes, and temperature variability, may profoundly affect xylem structure. However, genetic determination of quantitative anatomical features and differences related to the topographic position of the wood fragment must also be taken into consideration. *Capparidoxylon holleisii*, together with the fossil representatives of *Dichrostachys* and *Pistacia* (Tab. 5), has presumably been subjected to prolonged drought periods. *Capparis* is today widely distributed in arid-/semi-arid regions and open habitat in the Mediterranean and adjacent areas (ZOHARY 1960, 1983).

Acknowledgements

I would like to thank L. LANG (Ehekirchen, Germany) for providing access to the here presented fossil. The author is indebted to D. GROSSER (Munich, Germany) for granting permission to consult the XylotheC, Institut of Wood Research, at the Technical University Munich (Germany). Technical assistance has been provided by C. HEILBIG and R. ROSIN (both Munich, Germany). For valuable comments on the coprolites, I would like to thank M. BAEHR (Munich, Germany).

6. References

- ANDREWS, H. N. (1955): Index of Generic Names of Fossil Plants, 1820-1950; Washington DC (United States Government Printing Office), 261 pp.
- BOUREAU, E. (1957): Anatomie végétale, Tome III; Paris (Presses Universitaires), viii + pp. 525-725.
- BRUMMIT, R. K. & POWELL, C. E. (1992): Authors of Plant Names; Kew (Royal Botanic Gardens), 732 pp.
- CARLQUIST, S. (2001): Comparative Wood Anatomy; Berlin (Springer), 436 pp.
- DÉTIENNE, P. & JACQUET, P. (1983): Atlas d'identification des bois de l'Amazone et des régions voisines; Nogent-sur-Marne (Centre Technique Forestrier Tropical), 640 pp.
- DOPPLER, G. & SCHWERDT, K. (1996): Molasse. – In: FREUDENBERGER, W. (Ed.), Erläuterungen zur Geologischen Karte von Bayern 1:500.000; München (Bayerisches Geologisches Landesamt), 150-168.
- DOPPLER, G., FIEBIG, M. & MEYER, R. K. F. (2002): Erläuterung zur Geologischen Karte 1:100.000, Planungsregion Ingolstadt 10, München (Bayerisches Geologisches Landesamt), 172 pp.
- FAHN, A., WERKER, E. & BAAS, P. (1986): Wood Anatomy and Identification of Trees and Shrubs from Israel and Adjacent Regions; Jerusalem (The Israel Academy of Science and Humanities), 221 pp.
- GLASER, S., DOPPLER, G. & SCHWERDT, K. (2004): GeoBavaria, 600 Millionen Jahre Bayern; München (Bayerisches Geologisches Landesamt), 92 pp.
- GOTTWALD, H. P. J. (2002): Tertiäre Kieselhölzer der Südlichen Frankenalb. – Documenta naturae, **143**: 1-53.
- GOTTWALD, H. P. J. (2004): Neue taxonomische Untersuchungen an 205 tertiären Hölzern und 2 verkieselten Rindenresten aus der Südlichen Frankenalb und anderen Randgebieten mit Aussagen über Herkunft und Flora, Klima und Alter. – Documenta naturae, **153**: 1-93.
- GREGORY, M. (1994): Bibliography of Systematic Wood Anatomy of Dicotyledons (IAWA Journal Supplement 1); Leiden (Rijksherbarium, The Netherlands), 265 pp.
- HASS, H. & ROWE, N. P. (1999): Thin section and wafering. – In: JONES, T. P. & ROWE, N. P. (Eds), Fossil Plants and Spores: Modern Techniques; London (Geological Society), 76-81.
- HOLLEIS, P. (1992): Kieselhölzer aus randlichen Ablagerungen der Oberen Süßwassermolasse (südliche Frankenalb). – Documenta naturae, **7**: 30-35.
- HILLIS, H. E. (1987): Heartwood and Tree Exudates; Berlin etc. (Springer), 268 pp.
- IAWA COMMITTEE (1989): IAWA List of microscopic features for hardwood identification. – IAWA Bulletin, n.s., **10**: 219-332.
- ILIC, J. (1991): CSIRO, Atlas of Hardwoods; Berlin (Springer), 525 pp.
- JAGIELLA, C. & KÜRSCHNER, H. (1987): Atlas der Hölzer Saudi Arabiens (Beihefte zum Tübinger Atlas des Vorderen Orients, Reihe A, Nr. 20); Wiesbaden (L. Reichert), 176 pp.
- JANSONIUS, H. H. (1952): Key to the Javanese Woods on the Basis of Anatomical Features; Leiden (E. J. Brill), 244 pp.
- KRIBS, D. A. (1935): Salient lines of structural specialization in the wood rays of dicotyledons. – Botanical Gazette, **96**: 547-557
- KRUSE, H. O. (1954): Some Eocene dicotyledonous woods from Eden Valley, Wyoming. – The Ohio Journal of Science, **54**: 243 - 268.
- KUBITZKI, K. & BAYER, C. (2003): The Families and Genera of Vascular Plants, V; Berlin etc. (Springer), 418 pp.
- MABBERLEY, D. J. (1997): The Plant Book. 2nd revised edition; Cambridge (Cambridge University Press), 858 pp.
- METCALFE, C. R. & CHALK, L. (1950): Anatomy of the Dicotyledons, 2 vols; Oxford (Clarendon Press), 1500 pp.
- MILLER, R. B. (1978): Potassium calcium sulphate in the secondary xylem of *Capparis*. – IAWA Bulletin, **1978**: 50.
- MOMBACHER, R. (2003): Holz-Lexikon. Nachschlagewerk für die Holz- und Forstwirtschaft, 2 vols, 4th ed.; Leinfelden-Echterdingen (DRW-Verlag Weinbrenner GmbH & Co.), 1450 pp.
- NEUMANN, K., SCHOCH, W., DÉTIENNE, P. & SCHWEINGRUBER, F. H. (2001): Woods of the Sahara and the Sahel; Bern, Stuttgart, Wien

- (P. Haupt), 465 pp.
- NILOOFARI, P. (1961): Textbook of Wood Technology. Vol. I. Iranian Timbers; Teheran (University of Teheran), 155 pp.
- ROTH, L., DAUNDERER, M. & KORMANN, K. (1994): Giftpflanzen, Pflanzengifte, 4th ed.; Hamburg (Nikol Verlagsgesellschaft mbH & Co. KK), 1090 pp.
- SCHENK, A. (1883): Fossile Hölzer. – *Palaeontographica*, **30**: 1-19.
- SCHUBERT, R. & WAGNER, G. (1984): Pflanzennamen und botanische Fachwörter, 8th ed.; Melsungen (J. Neumann-Neudamm), 882 pp.
- SCHWEINGRUBER, F. H. (1991): Atlas mitteleuropäischer Hölzer; Bern, Stuttgart (P. Haupt), 802 pp.
- SELMEIER, A. (1970): *Castanopsis*-Hölzer aus obermiozänen Glimmersanden der südlichen Frankenalb. – Mitteilung der Bayerischen Staatssammlung für Paläontologie und historische Geologie, **10**: 309-320.
- SELMEIER, A. (1984a): Kleinporige Laubhölzer (Rosaceae, Salicaceae) aus jungtertiären Schichten Bayerns. – Mitteilung der Bayerischen Staatssammlung für Paläontologie und historische Geologie, **24**: 121-150.
- SELMEIER, A. (1984b): Fossile Bohrgänge von *Anobium* sp. in einem jungtertiären Lorbeerholz aus Egweil (südliche Frankenalb). – *Archaeopteryx*, **1984**: 1-17.
- SELMEIER, A. (1986): Jungtertiäre Juglandaceen-Hölzer aus der Südlichen Frankenalb Bayerns. – *Archaeopteryx*, **1986**: 61-78.
- SELMEIER, A. (1989): Ein jungtertiäres *Celtis*-Holz (Ulmaceae) aus der Südlichen Frankenalb (Bayern). – *Archaeopteryx*, **7**: 35-50.
- SELMEIER, A. (1991): Ein verkieseltes Sapotaceae-Holz, *Bumelioxylon holleisii* n. gen., n. sp. aus jungtertiären Schichten der Südlichen Frankenalb. – *Archaeopteryx*, **9**: 55-72.
- SELMEIER, A. (1998a): *Shoreoxylon holleisii* n. sp. from Tertiary sediments of Attenfeld, southern Franconian Alb, Germany. – *Archaeopteryx*, **16**: 77-99.
- SELMEIER, A. (1998b): Aufsammlung von Kieselhölzern aus tertiären Schichten Süddeutschlands, der Schweiz und aus Österreich. – Mitteilungen der Bayerischen Staatssammlung für Paläontologie und historische Geologie, **38**: 275-300.
- SELMEIER, A. (2000a): A silicified *Pistacia* wood, Anacardiaceae, from the Upper Miocene of southern Franconian Alb, Germany. – Mitteilung der Bayerischen Staatssammlung für Paläontologie und historische Geologie, **40**: 2-212.
- SELMEIER, A. (2000b): Liste der Tertiärhölzer aus dem nordalpinen Molassebecken und der Südlichen Frankenalb (Deutschland). – *Archaeopteryx*, **18**: 101-109.
- SELMEIER, A. (2001): Die versteinerte Gehölzsammlung des Historischen Vereins Neuburg a. d. Donau. – Neuburger Kollektaneenblatt, **149**: 123-182.
- SELMEIER, A. (2003a): Tertiary *Cedrela* woods (Meliaceae) from the North Alpine Foreland in Southern Germany and Austria. – *Zitteliana*, **A43**: 159-170.
- SELMEIER, A. (2003b): Anatomische Identifizierung tertiärer Holzreste aus der südlichen Frankenalb Bayerns, Deutschland. – *Archaeopteryx*, **21**: 37-52.
- SELMEIER, A. (2004): Versteinertes Zigarrenkistenholz Cedro, Familie Meliaceae, in der Sammlung des Historischen Vereins Neuburg a. d. Donau e. V. – Neuburger Kollektaneenblatt, **152**: 219-241.
- STERN, W. L. (1988): Index Xylariorum. Institutional wood collections of the world, 3. – *IAWA Bulletin*, n.s., **9**: 203-252.
- WHEELER, E. A. (1986): Vessels per square millimetre or vessel groups per square millimetre? – *IAWA Bulletin*, n. s., **7**: 73-74.
- WHEELER, E. A. (1991a): Fossil Wood Database 11 March 1991; Raleigh NC (North Carolina State University), 32 pp.
- WHEELER, E. A. (1991b): Database references: March 1991; Raleigh NC (North Carolina State University), 21 pp.
- WHEELER, E. A. & BAAS, P. (1998): Wood identification – a review. – *IAWA Journal*, **19**: 241-284.
- WHEELER, E. A. & MANCHESTER, S. R. (2002): Woods of the Middle Eocene Nut Beds Flora (*IAWA Journal Supplement 3*); Leiden (Nationaal Herbarium Nederland), 188 pp.
- ZOHARY, M. (1960): The species of *Capparis* in the Mediterranean and the Near Eastern countries. – *Bulletin Research Council Israel*, **80**: 49-64.
- ZOHARY, M. (1983): Pflanzen der Bibel; Stuttgart (Calwer Verlag), 222 pp.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Zitteliana Serie A](#)

Jahr/Year: 2005

Band/Volume: [45](#)

Autor(en)/Author(s): Selmeier Alfred

Artikel/Article: [Capparidoxylon holleisii nov. spec., a silicified Capparis \(Capparaceae\) wood with insect coprolites from the Neogene of southern Germany 199-209](#)