


Zitteliana

The background of the cover is a detailed photograph of a fossilized trilobite. The trilobite is shown in a split view, with the left side showing the segmented body and the right side showing the rounded cephalon. The fossil is embedded in a light-colored, textured rock matrix.

An International Journal
of Palaeontology and Geobiology

Series B/Reihe B
Abhandlungen der Bayerischen Staatssammlung
für Paläontologie und Geologie

26

4th International Symposium
on Lithographic Limestone and Plattenkalk

Eichstätt/Solnhofen, Germany

September 12th-18th, 2005

- Abstracts and Field Trip Guides -

München 2005

Zitteliana

An International Journal of Palaeontology and Geobiology

Series B/Reihe B

Abhandlungen der Bayerischen Staatssammlung für Paläontologie und Geologie

26

4th International Symposium on Lithographic Limestone and Plattenkalk

Eichstätt/Solnhofen, Germany

September 12th-18th, 2005

Organised by

Martina Kölbl-Ebert

Jura-Museum, Eichstätt

Martin Röper

Bürgermeister-Müller-Museum, Solnhofen

Reinhold R. Leinfelder

Staatliche Naturwissenschaftliche Sammlungen Bayerns,
^{LMU}München, Department für Geo- und Umweltwissenschaften, Sektion Paläontologie,
and GeoBio-Center^{LMU}

– Abstracts and Field Trip Guides –

Zitteliana	B 26	96 Seiten	München, 05.09.2005	ISSN 1612-4138
------------	------	-----------	---------------------	----------------

Editors-in-Chief/Herausgeber: Reinhold R. Leinfelder, Michael Krings

Volume Editor: Winfried Werner

Production and Layout/Bildbearbeitung und Layout:

Stefanie Klug

Bayerische Staatssammlung für Paläontologie und Geologie

Richard-Wagner-Str. 10, D-80333 München, Deutschland

<http://www.palaeo.de/zitteliana>

email: zitteliana@lrz.uni-muenchen.de

Für den Inhalt der Arbeiten sind die Autoren allein verantwortlich.

Authors are solely responsible for the contents of their articles.

Copyright © 2005 by Bayerische Staatssammlung für Paläontologie und Geologie, München

Die in der Zitteliana veröffentlichten Arbeiten sind urheberrechtlich geschützt.

Nachdruck, Vervielfältigungen auf photomechanischem, elektronischem oder anderem Wege sowie die Anfertigung von Übersetzungen oder die Nutzung in Vorträgen, für Funk und Fernsehen oder im Internet bleiben – auch auszugsweise – vorbehalten und bedürfen der schriftlichen Genehmigung durch die Bayerische Staatssammlung für Paläontologie und Geologie, München.

ISSN 1612-4138

Druck: Gebr. Geiselberger GmbH, Altötting

Published with the support of the
Deutsche Forschungsgemeinschaft (DFG)

Cover illustration: *Mesolimulus walchi* DESMAREST; horse-shoe crab with its trail; Lower Tithonian, Solnhofen (BSPG AS I 944).

Umschlagbild: *Mesolimulus walchi* DESMAREST; Pfeilschwanzkrebs mit Fährte; Lower Tithonian, Solnhofen (BSPG AS I 944).

Zitteliana

An International Journal of Palaeontology and Geobiology

Series B/Reihe B

Abhandlungen der Bayerischen Staatssammlung für Paläontologie und Geologie

26

CONTENTS/INHALT

MARTINA KÖLBL-EBERT & WINFRIED WERNER Preface	3
Abstracts	6
Geology and Field Trip Guides	
DIETER SCHMID, REINHOLD R. LEINFELDER & GÜNTER SCHWEIGERT Stratigraphy and Palaeoenvironments of the Upper Jurassic of Southern Germany	31
ROMAN KOCH & CHRISTIAN WEISS Field Trip A: Basin-Platform Transition in Upper Jurassic Lithographic Limestones and Dolomites of the Northern Franconian Alb (Germany)	43
MARTIN RÖPER Field Trip B: East Bavarian Plattenkalk – Different Types of Upper Kimmeridgian to Lower Tithonian Plattenkalk Deposits and Facies	57
MARTIN RÖPER Field Trip C: Lithographic Limestones and Plattenkalk Deposits of the Solnhofen and Mörnsheim Formations near Eichstätt and Solnhofen	71
GÜNTER SCHWEIGERT, GERD DIETL & ROMAN KOCH Field Trip D: The Nusplingen Plattenkalk and Other Fossil Sites in the Western Swabian Alb (SW Germany)	87

Zitteliana	B 26	96 Seiten	München, 05.09.2005	ISSN 1612-4138
------------	------	-----------	---------------------	----------------

ABSTRACTS

ARRATIA, G.: The Contribution of Late Jurassic Fishes from Germany to Understand the Evolutionary History of Teleosts	7
BAUSCH, W. M.: Geochemical Position of the Franconian Plattenkalk	7
BAUSCH, W. M.: Revision of the Age Relationships of Maxberg/Solnhofen	8
BENNETT, S. C.: A Review of <i>Ctenochasma</i> : Taxonomy and Ontogeny	8
BILLON-BRUYAT, J.-P.: Comparison of the Tetrapod Assemblages from the Late Jurassic Plattenkalk Deposits of Western Europe	9
BRITO, P. M.: The Lower Cretaceous Crato Formation's Ichthyofauna (Araripe Basin, Northeastern Brazil)	9
BURNHAM, D. A.: The Paleocology of <i>Archaeopteryx</i> - A Re-Evaluation	10
CARANNANTE, G., SIGNORE, M., SIMONE, L. & VIGORITO, M.: Vertebrate-fossil Rich Plattenkalk of Pietrarola (Southern Appennines, Italy). A Sedimentological and Taphonomical Approach	10
CHIAPPE, L. M. & GÖHLICH, U.: A new Theropod Dinosaur from the Upper Jurassic Limestones of Schamhaupten (Bavaria, Germany)	11
DIETL, G. & SCHWEIGERT, G.: The Upper Kimmeridgian Nusplingen Plattenkalk (W Swabian Alb, SW Germany)	11
FUCHS, D.: Coleoid Cephalopods from the Lithographic Limestones of Hâqel (Lebanon) - A Comparison with Solnhofen (Germany)	12
FÜRSICH, F. T., MÄUSER, M., SCHNEIDER, S. & WERNER, W.: Sedimentology and Palaeoecology of Laminated Limestones from the Kimmeridgian of the Northern Franconian Alb (Southern Germany)	13
GAILLARD, C.: Recent and Fossil Chelonian Kingdoms	13
GARASSINO, A., DE ANGELI, A. & SCHWEIGERT, G.: Brachyurans from the Upper Jurassic (Tithonian) of Pfalzpaint and Breitenhill (Bavaria, South Germany)	14
GARASSINO, A. & SCHWEIGERT, G.: The Upper Jurassic Solnhofen Decapod Crustacean Fauna: Review of the Types from Old Descriptions	15
IFRIM, C., STINNESBECK, W. & FREY, E.: Upper Cretaceous (Cenomanian-Turonian and Turonian-Coniacian) Open Marine <i>Plattenkalk</i> - Deposits in NE Mexico	15
JURKOVSEK, B. & KOLAR-JUROKOVSEK, T.: Fossil Assemblages from the Upper Cretaceous Carbon Rich Sediments of Kras, Slovenia	16
KEUPP, H.: Complete Ammonoid Jaw Apparatuses from the Solnhofen Plattenkalks: Implications for Aptychi Function and Microphage Feeding of Ammonoids	17
KLUG, C., SCHWEIGERT, G., DIETL, G. & FUCHS, D.: Coleoid Beaks from the Nusplingen Lithographic Limestone (Late Kimmeridgian, SW Germany)	17
KLUG, S. & KRIWET, J.: Occurrence and Diversity of Palaeospinacid Sharks (Neoselachii, Synechodontiformes) from the Upper Jurassic Lithographic Limestones of South Germany	18
KOCH, R.: New Data on Sedimentology of Upper Jurassic Carbonate Rocks from the Southern Franconian Alb	18
KOCH, R.: Characteristics of Solnhofen Monument Stones	19
KRIWET, J. & KLUG, S.: Late Jurassic Carpetsharks (Neoselachii, Orectolobiformes) from South Germany	19
MARTY, D.: Sedimentology and Taphonomy of Dinosaur Track-bearing Plattenkalke (Kimmeridgian, Canton Jura, Switzerland)	20
MICKLICH, N.: The Fish Fauna of the Messel Pit: Rule or Exception?	20
PAPAZZONI, C. A. & TREVISANI, E.: New Data about the Age and Palaeoenvironment of the „Pesciara di Bolca“ Fossil-Lagerstätte	21
RUSCONI, M., LOMBARDO, C. & TINTORI, A.: New Colobodontid (Actinopterygians) from the Carnian (Late Triassic) of Northern Italy	22
SCHWEIGERT, G.: Ammonite Biostratigraphy as a Tool for Dating Upper Jurassic Lithographic Limestones from South Germany - First Results and Open Questions	22
SCHWEIGERT, G. & GARASSINO, A.: The Lobster Genus <i>Squamosoglyphea</i> BEURLIN, 1930 (Crustacea, Decapoda, Glypheidae) in the Upper Jurassic Lithographic Limestones of Southern Germany	23
SKAWINSKA, J.: Development of a Jurassic Basin in Terms of Palaeohydrodynamic Reconstruction (Miechów Trough, Poland)	24
TINTORI, A., BONA, F. & ZORZIN, R.: Bolca (Eocene, Verona) Fossils under a New Light	24
TINTORI, A. & LOMBARDO, C.: The Middle Triassic Vertebrate Levels in the Western Tethys: Are Fossil Fishes Useful for Biostratigraphy?	25
TISCHLINGER, H.: Ultraviolet Light Investigations of Fossils from the Upper Jurassic Plattenkalks of Southern Franconia	26
VIOHL, G. & ZAPP, M.: Schamhaupten, an Outstanding Fossil-Lagerstätte in a Silicified Plattenkalk (Kimmeridgian-Tithonian Boundary, Southern Franconian Alb, Bavaria)	27
WEISS, C., KOCH, R. & HUANG, W.: Diagenesis of Thin Bedded Carbonate Rocks in the Cambro-Ordovician Sequences of the North China Carbonate Platform	28
WELLNHOFER, P. & RÖPER, M.: The Ninth Specimen of <i>Archaeopteryx</i> from Solnhofen	28
ZEISS, A., SCHULBERT, C. & VIOHL, G.: The Ammonites from Schamhaupten (Southern Franconian Alb, Bavaria) - An Interesting Faunal Association at the Boundary Upper Kimmeridgian/Lower Tithonian	29

Field Trip D

The Nusplingen Plattenkalk and Other Fossil Sites in the Western Swabian Alb (SW Germany)

September 17th/18th, 2005

By

Günter Schweigert¹, Gerd Dietl¹ & Roman Koch²

¹ Staatliches Museum für Naturkunde, Rosenstein 1, D- 70191 Stuttgart, Germany

² Institut für Paläontologie der Universität Erlangen-Nürnberg, Loewenichstr. 28, D-91054 Erlangen, Germany

First day (Saturday 17th, 2005)

The field trip starts in Eichstätt and goes to the western part of the Swabian Alb, to the area of the Upper Danube valley. From the town of Sigmaringen westwards the Danube valley is developed as a canyon in which the Upper Jurassic rocks are well exposed. First we give an overview on the geological setting, common lithologies, and the palaeoenvironment of the Upper Jurassic. Few kilometres north of the Upper Danube valley, two important sites of “Plattenkalk” exist: the Kolbingen Plattenkalk and the more famous Nusplingen Plattenkalk. The first locality is visited on Saturday, the latter on Sunday. The accommodation from Saturday to Sunday will be a guest house in the small village of Beuron, aside a large monastery.

Stop 1: Schaufels and other rocks of the Upper Danube Valley between Sigmaringen and Beuron

Geological Map 1:25 000 sheet 7920 Leibertingen,
R 3505400, H 5328100

The Schaufels rock is one of the most famous rocks for climbing activities in the Upper Danube Valley area. In a facies mapping study the previously undifferentiated Upper Jurassic massive limestone facies (Massenkalk Formation) of an area of ca. 8 km² W of the Schaufels (including the Schaufels itself) ranging from the time equivalents of the “Untere Felsenkalk” (ki2) to the “Liegende Bankkalk” (ki4) formations has been differentiated into 11 facies types (VOLK et al. 2001) (Fig. 2).

The investigated massive limestones can be subdivided into **five facies groups (FG I-V)** and by a more detailed classification into **eleven facies types (FT-1-11)**.




Figure 1: Map showing the location of stop 1-3 (first day) and the fossil site Nusplingen, 12 km N of the River Danube, SW Germany. Modified from DIETL & SCHWEIGERT (2004)

Flaser-bedded limestones and sponge-marls facies (FG-I) can be differentiated into mud- to packstones containing sponges and tuberoids (FT-1), sponge-rich wacke- to floatstones (FT-2), and sponge-marls (FT-3).

Thick-bedded facies types of the Kimmeridgian (FG-II) are built up by wackestones with silicification and sponges (FT-4), and wacke- to packstones with sponges (FT-5).

The peloid-lithoclast-oid sand facies (FG-3) consists of peloid-lithoclast-wackestones with sponges and occasional silicification (FT-6), peloid-lithoclast-wacke- to packstones with sparse sponges (FT-7), ooid-lithoclast pack- to grainstones (FT-8) and stromatolitic boundstones in peloid-lithoclast-oid sands (FT-9).

Micritic breccias and tuberolites of the higher Kimmeridgian (FG-4) are built up of FT-10.

Stratigraphie		Ki 2.1	Ki 2.2	ki 2.3	ki 2.4	ki 3	ki 4
Flaserkalk- und Schwamm-Mergelfazies							
Schwamm- und tuberoïd-führende Mud- Packstones	FT 1	■		■			
Schwammreiche Wacke- Floatstones	FT 2	■		■			
Schwamm-Mergel	FT 3	■		■			
Dickbankige Faziestypen							
Schwammführende Wackestones	FT 4			■			
Schwammführende Wacke- Packstones	FT 5			■			
Peloid-Lithoklast-Ooid-Sandfazies							
Schwammführende Peloid-Lithoklast-Wackestones	FT 6			■		■	
Schwammf. Peloid-Lithokl.-Ooid-Wacke- Packstones	FT 7			■		■	
Ooid-Lithoklast-Pack- Grainstones	FT 8				■		
Stromatolith. Boundstones + Peloid-Lithokl.-Ooid-Sande	FT 9					■	
Brekzien und Tuberolithe							
Mikritische Brekzien und Tuberolithe	FT 10					■	
Mikritreiche Massenkalkfazies							
Mud- Packstones mit mikritischen Lithoklasten	FT 11						■

Figure 2: Distribution of facies groups (FG-I-V) and the subdivision in facies types FT-1 to FT-11 in the Kimmeridgian. Their relative abundance is indicated as rare, common, abundant, and very abundant (VOLK et al. 2001).

The massive limestone facies of the Ki4 consists of mudstones to packstones with micritic lithoclasts (FT-11).

Diagenetically strongly altered sediments (FG-5) are represented by saccharoidal, vuggy dolomite (so-called Lochfels), and recrystallized, orange-brown sediments of FT-7.

The distribution of facies types is controlled by the evolution of relative sea level changes. In the stratigraphic succession there is a trend towards a shallower depositional environment from the ki2 to the ki3, reaching a maximum at the transition ki3/4. Whereas the beginning of ki2 micritic sediments of the flaser-bedded limestones and sponge-marl facies prevail, towards the ki2.4 there is a progressive shift towards more carbonate-rich sediments of the peloid-lithoclast-ooid sand facies. The depositional environment of the sand facies is interpreted as a carbonate platform with minor morphological features where carbonate sand shoals have been deposited in a water depth of less than 10 m (VOLK et al. 2001). From the ki2.4 lenses of ooid-lithoclast-packstones to grainstones (FT-8) and stromatolitic boundstones (FT-9) can be found within the wide-spread areas of peloid-lithoclast-ooid wackestones to packstones (FT-7). The shallow marine environment of moderate to episodic higher water energy is indicated by the presence of primary open interparticle porosity in which marine isopachous cements occur locally followed by granular cements occluding the relic pore space.

A cyclic development in the deposition of these facies types was revealed in the latest study on these sediments from the same area by PAWELLECK & AIGNER (2003). First analyses of

these type of carbonate sands in the area of the Schaufels were carried out by SCHALLER & KOCH (1996) (Fig. 3) who already gave hints to the cyclic development of these shallow water sediments in a profile of 84 m thickness of the Kimmeridgian 2.4 to the Kimmeridgian 3 which gives a detailed analyses of the sediments of this interval as indicated by VOLK et al. (2001). The section can be subdivided from base to top in three units as follows: (A) sponge bearing peloid wackestones to packstones (21 m), (B) ooid-peloid packstones to grainstones (30 m), and (C) carbonate sand facies with abundant microbial crusts and stromatolitic boundstones rich in sponges.

All these interpretations are completely contrasting the former interpretation as massive sponge-algal reefs or sponge-magnafacies to be deposits in a water depth of some tens to even 150 m as reported by many authors. Basic ideas on the sedimentation of carbonate sand facies and the distribution of real spongiolites were published by KOCH et al. (1994) (Fig. 4) from the Swabian Alb opening the window for a new insight in the formation of so-called sponge build-ups (sponge-reefs, sponge-biostroms etc.) in the Upper Jurassic of Southern Germany. This facies development is assumed for most outcrops in the direction to Beuron as indicated by moderate horizontal thick-bedded limestones which are typical of carbonate sand deposits and proven by local samples indicating the same facies. This carbonate sand facies occurs in direction to Sigmaringen up to the quarry of Thiergarten, where dolomitic sponge-buildups are developed (HEISS 1990). This facies marks the slope facies of carbonate sands settled by




Figure 3: The Schaufels section. Besides geological age, thickness and position of samples the following parameter are indicated: the classification according to DUNHAM (1962), the five different facies types according to SCHALLER & KOCH (1996), the carbonate content, and the semiquantitative estimated amount of components (1 = microbial crusts, 2 = sponges, 3 = peloids, 4 = ooids). Furthermore the three large units A, B, and C are indicated (from SCHALLER & KOCH 1996).

sponges and interfingering with a deeper basin facies according to the model of KOCH et al. (1994).

The deposition of sponge-marls is restricted to depositional environments with a lower water energy level. The deposition of more micritic limestones in the ki4 is interpreted as relative sea level rise.

Stop 2: Viewpoint “Knopfmacherfelsen” W of Beuron

Geological Map 1:25 000 sheet 7919 Mühlheim a. d. Donau,
R 3496500, H 5321250

During the Pleistocene the River Danube eroded a canyon into the Upper Jurassic rocks of the southern border of the Swabian Alb. At a tourist viewpoint called “Knopfmacherfelsen” the enormous palaeorelief formed by late Kimmeridgian sponge-microbial mounds and corresponding lithologies is

excellently visible. Although the rock is rather massive in appearance (‘Massenkalk Formation’) and very poor in age-diagnostic guide fossils, some major marker surfaces allow a good resolution even in the logs of drill holes. In the east, at the monastery of Beuron, the Massenkalk rocks have thin marly intercalations pointing to the ‘Glaukonitbank’ marker bed of the Untere Felsenkalk Formation in bedded facies.

South of “Knopfmacherfelsen” the massive rocks are forming an atoll-like structure, which is, however, open to the direction of the Danube valley. In his overview on the development of the Upper Jurassic in Swabia GWINNER (1976) presented a nice sketch of this area (Fig. 5). We are looking at the northern border of this structure; with the castle of Bronnen on top of the rock. The incorporated basin in the South is filled by the bedded facies of the upper Kimmeridgian, the Liegende Bankkalk and Zementmergel formations of late Kimmeridgian age (Beckeri Zone, Ulmense Subzone). These strongly bioturbated basinal deposits are coeval with the laminated Nusplingen Plattenkalk which is located about 10 km to the north. In a Pliocene karstic fill located between




Figure 4: The distribution of spongiolites in massive carbonate rocks (reef-complexes, Massenkalk, etc.). Comparison of the classical interpretation with the new interpretation by KOCH et al. (1994). Note that the peloid-litholasts-sand facies makes up the largest amount of the “Massenkalk”. The spatial distribution of real algal-sponge build-ups in and around the sand facies is interpreted to be caused by varying hydrodynamic conditions triggered by the palaeogeographic distribution of the sand facies.




Figure 5: Sketch of the area in the vicinity of viewpoint “Knopfmacherfelsen” showing the former palaeorelief of the Upper Jurassic prepared by erosion (from GWINNER 1976).

Beuron and “Knopfmacherfelsen” several reworked Jurassic hermatypic corals like *Isastrea* were found (FRANZ et al. 1997). They indicate the local occurrence of coral reefs and shallow conditions in the Upper Jurassic of this area.

Stop 3: The Kolbingen Plattenkalk

Geological Map 1:25 000 sheet 7919 Mühlheim a. d. Donau,
R 3490450, H 5326500

About 5 kilometres N of the village of Kolbingen there is an area where a platy limestone was quarried over centuries, even since the times of the Romans. Today the last quarry is almost abundant, and only temporarily this “Kolbingen Plattenkalk” is still exploited. The world-famous naturalist ALEXANDER VON HUMBOLDT (1823) was the first to mention this locality, and he compared it with the Solnhofen Lithographic Limestone of Franconia. More than one decade later, about 6 kilometres to the north, the Nusplingen Plattenkalk was discovered by F. A. QUENSTEDT, and both occurrences were often correlated. However, from this Kolbingen Plattenkalk macrofossils have never been recorded, and also its exact stratigraphical position remained unknown. New investigations have been carried out but the results are still unpublished; a preliminary version is presented here.

Within the quarry area, the Kolbingen Plattenkalk overlies a massive, reddish- to violet-coloured dedolomite (Massenkalk Formation). The total thickness of this member is about 5 metres. It consists of 2 different lithologies, of which today only the lower one is still visible in the field. It is made up of a 2.5 metres thick, finely laminated yellowish to brownish limestone (Fig. 6). Macrofossils have not been found yet, only some round, irregularly scattered structures on bedding planes may be interpreted as ichnofossils. At some places the laminates were overlain by a grey micritic limestone, which weathers into thin plates. This pseudo-lithographic fabric results from shear planes crossing at very low angles. The cleavage surfaces of the plates are covered with typical plumose structures. Both GWINNER (1976) and TEMMLER (1966) erroneously interpreted these plumose structures (Fig. 7) as a complex system of sloping scratches. It cannot be excluded that the cleavage follows a hidden cross bedding which is not discernible because of the fine micritic material.

350 metres SW of the quarry area, towards the Lippach valley, there is a well-exposed outcrop of the Upper Jurassic. The section starts with the Obere Felsenkalk Formation, bedded micritic limestones with horizontal layers of siliceous concretions. Within this formation several ammonite faunal horizons of Setatum Subzone of the late Kimmeridgian have been recognized. The Liegende Bankkalk Formation overlies the Obere Felsenkalk Formation with a sharp disconformity. The Liegende Bankkalk Formation consists of thinner-bedded greyish micritic limestones interfingering with sponge-microbial rock-forming mounds. Near the top of the section there is a thicker marl bed which can be classified as belonging to the Zementmergel Formation. At the top there is a brownish dolomitic bed which grades into a finely laminated, not recrystallized Kolbingen Plattenkalk. Several trace fossils, small-

scale cross bedding and so-called ‘*Kinneya* ripples’ on bedding planes indicate a very shallow environment. Interestingly the bedding planes are strongly dipping to the west, whereas the Kolbingen Plattenkalk in the quarry area lies almost horizontal. This shows that the Kolbingen Plattenkalk was not deposited in a basin as the nearby Nusplingen Plattenkalk, but on a high structure. Looking again at the lamination of the Kolbingen Plattenkalk in the quarry area its cyclicity strongly resembles that of tidal laminates (cf. ARCHER & FELDMAN 1994, TESSIER et al. 1995, BUATOIS et al. 1997). Within the section about 100 cyclothems can be counted. This very rapid sedimentation rate and an environment unfavourable for nektonic animals like ammonites or belemnites are probable reasons for the lack of macrofossils.

Biostratigraphically the uppermost fossiliferous bed little below the laminates belong to the Ulmense Subzone of the late Kimmeridgian (*zio-wepferi* Horizon beta, see SCHWEIGERT 1998). This means that the formation of laminates started little earlier in the nearby Nusplingen Plattenkalk than here at the Kolbingen site. But unless the Kolbingen Plattenkalk is an approximate time equivalent to the Nusplingen Plattenkalk, it differs strongly both in lithology and genesis.

The outcrops of the Kolbingen Plattenkalk and the Nusplingen Plattenkalk are separated from each other by a zone



Figure 6: Rock sample from the laminated lower part of the Kolbingen Plattenkalk.




Figure 7: Plumose structures occurring in the micritic upper part of the Kolbingen Plattenkalk.

of intensive diagenetic recrystallization (“Zuckerkörniger Lochfels”). Possibly this recrystallization results from an intra-Jurassic freshwater diagenesis occurring in the subcrop of small islands. Additional hints for this assumption are dedolomite clasts in turbidites of the Nusplingen Plattenkalk (BANTEL et al. 1999), some insects, many land plants and plant debris, several pterosaurs, and the recent discovery of a piece of fossil charcoal.

Second day (Sunday 18th, 2005)

The field trip starts in Beuron and follows the Bärental valley northwards to the village of Nusplingen. The outcrops are located W of the village on the “Westerberg” hill. Along the road from Nusplingen to the “Westerberg” an Upper Jurassic is exposed mainly consisting of spongiolithic limestones of Oxfordian to Kimmeridgian age. In an abandoned quarry the strong palaeorelief developed in this area is well visible.

Geological setting of the Nusplingen Plattenkalk

The Nusplingen Plattenkalk, also known in literature as “Nusplingen Lithographic Limestone”, represents a ‘Fossil Lagerstätte’ similar to the famous lower Tithonian Solnhofen Lithographic Limestone in Bavaria. It is located in the western part of the Swabian Alb, about 12 kilometres N of the Upper Danube Valley (Fig. 1). It is the only occurrence of fossiliferous laminated limestones in Swabia. These laminated limestones were deposited in a more or less anoxic environment of a lagoon surrounded by sponge/microbial mounds. Some of them were partly tectonically uplifted over the sea level as small islands. The palaeoenvironment of the Nusplingen lagoon is reconstructed in Fig. 8.

Spectacular fossils from the “Nusplinger Plattenkalk” were reported since the middle of the 19th century. Several excavation campaigns took place, at first with more commercial aspects




Figure 8: Reconstruction and palaeogeographic setting of the Nusplingen lagoon – without scale. From DIETL & SCHWEIGERT (2004).

as prospecting for limestones to be used for lithography or for roof tiles, later restricted to scientific interests. The Nusplingen Plattenkalk Limestone is of late Kimmeridgian age (Fig. 9) and thus approximately 0.5 my older than the lithographic limestones of Solnhofen and Eichstätt. Since 1983 the whole area of the Nusplingen Plattenkalk is a Protected Excavations Area because of its extraordinary fossils. The Nusplingen Lithographic Limestone is actually well exposed and excavated in two small quarries (Nusplingen and Egesheim quarries). The total thickness of the Nusplingen Plattenkalk is between 10.5 and 17 metres (Fig. 10). Detailed sections are described in DIETL et al. (1998) and in BANTEL et al. (1999). General overviews on this fossil site are given in DIETL & SCHWEIGERT (1999, 2001, 2004), for a bibliography of older publications see SCHWEIGERT (1997).

New excavations by the Natural History Museum of Stuttgart in the above mentioned two quarries have proliferated new data on sedimentology, palaeoecology, taphonomy, biostratigraphy, and especially on the fossil content of the Nusplingen Lithographic Limestone. Since 1993 more than 8.000 fossils were recovered. They belong to 320 taxa, many of them are new. They tell us interesting stories about sea life in or around the Nusplingen lagoon and from the surrounding shallow sea of the Upper Jurassic in SW Germany. In contrast to Solnhofen Nusplingen had a pelagic situation far away from larger continental islands like the Rhenish and Bohemian Massifs in the N and in the NE. Another interesting aspect is the preservation of the fossils within the Nusplingen Plattenkalk. Some of the fossils which are embedded in bituminous layers occurring in the upper part of the succession exhibit an extraordinary preservation of organic matter.

Fossil content

The Nusplingen Plattenkalk yields a significant amount of debris from the nearby reefs and from the water column. Among the microfossils the coccolithophorids are of great importance, because they characterize the laminates as marine alginites. Some bedding planes exhibit structures pointing to a temporary development of microbial mats. More than 60 species of radiolarians were identified by dissolving the limestone with acid.

In the higher part of the succession, especially in the bituminous layers, many land plants occur. They are derived from nearby uplifted islands and still preserved with their organic tissue showing finest anatomical details like cell-structures and stomata. Some years ago we were even able to identify amber. This amber occurs in situ within the resin vessels of araucariacean cone scales.

The most frequent invertebrate fossils of the Nusplingen Plattenkalk Limestone are ammonites and their aptychi. Some ammonites are very spectacular because they are preserved with their complete jaw apparatuses and stomach contents. The lower jaw is the well-known calcitic aptychus, whereas the upper jaw which was originally chitinous is preserved in carbon. Also nautilids have been found with both jaws still in the body chamber, sometimes even together with stomach-content. The belemnites occur frequently with a single species. Their

guards are very often bitten by predators. Teuthoids are often preserved with their ink sacs. Some years ago we discovered the first teuthoids with their beaks still in original position.

Among the arthropods, marine polychaetes sometimes exhibit soft tissue preservation. The horseshoe crab *Mesolimulus* is quite rare. One of the specimens, however, shows its compound eyes and even the muscles preserved, together with microbial layers covering the phosphatic muscle tissue (BRIGGS et al., in press). The newly excavated fossils include the first record of a Jurassic centipede. Other remarkable fossils are several insects like beetles and dragonflies. Most of these specimens are preserved in organic matter. Decapod crustaceans, especially natants, are typical of the Nusplingen Plattenkalk, and many of them are well-preserved. About 20 different taxa have been recorded. One of them still shows its compound eyes. At the end of the annual campaign of 2004 a marine isopod has been found, also with its preserved compound eyes. Other benthic organism such as echinoids, brittle stars and crinoids occur only sporadically. Especially the echinoids are usually bitten resulting from the activity of predators. The most frequent echinoderm, however, is the small planktonic crinoid *Saccocoma*. The isolated ossicles of this crinoid also occur within the widespread coprolites known as *Lumbricaria*, which were produced by ammonites – as it is demonstrated from crop contents.

Among the vertebrate fauna the fishes predominate. They are represented by Selachians, holocephalans and numerous actinopterygians. Most typical of the Nusplingen Plattenkalk

is the endemic angel shark *Squatina acanthoderma* which is preserved with its complete skeleton and the outline of the skin.

Reptiles rarely occur. They are represented by three genera of pterosaurs and at least two genera of marine crocodiles. In summer 2004 the excavation team of the Stuttgart Museum recovered an almost complete skeleton of a *Geosaurus* with remains of the skin and its stomach content.

In contrast to the typical Lithographic Limestone of Solnhofen, the ichnofauna of the Nusplingen Plattenkalk is much more diverse, but mostly restricted to special layers. Coprolites are generally frequent indicating marine life in the water column above the anoxic sea floor.

Stop 1: Type locality of the “Nusplingen Plattenkalk”

Geological Map 1:25 000 sheet 7819 Meßstetten,
R 3491520, H 5331810

The type locality of the Nusplingen Plattenkalk is a small outcrop from the first half of the 19th century. For the first time the famous Jurassic geologist F. A. QUENSTEDT from the University of Tübingen visited this locality in 1839 and described the “Nusplingen Plattenkalk” in 1843 from here. Although




Figure 9: General stratigraphical position of the Nusplingen Plattenkalk Limestone (Nuspl. Pl. Fm.) within the Upper Jurassic of Swabia (S Germany). From DIETL & SCHWEIGERT (2004).

QUENSTEDT was not able to find many fossils at that time, he concluded that the Nusplingen Plattenkalk is much richer in fossils than the Lithographic Limestones of Solnhofen. The small quarry goes back to the activity of a local farmer which was characterized by QUENSTEDT as an “industrial farmer”.

Stop 2: Nusplingen Plattenkalk in the Egesheim quarry

Geological Map 1:25 000 sheet 7819 Meßstetten,
R 3490540, H 5331580

The Egesheim quarry was exposed for extraction of road material in 1979/80. Because of intervention of the State Museum of Natural History Stuttgart the quarrying stopped at the end of 1980 because of the danger to destroy important fossils. Here are the lowermost 5 metres of the Nusplingen Plattenkalk Formation exposed. In this outcrop the excavations reached the bottom of the plattenkalk lagoon formed by the so called Liegende Bankkalk Formation, a highly bioturbated micritic limestone thus contrasting the nearly non-bioturbated lithographic facies. The bottom of the lagoon is dipping with about




Figure 10: Detailed section of the Nusplingen Plattenkalk and its stratigraphical position within the upper Kimmeridgian. The thickness of the Nusplingen Plattenkalk in the figured section is of about 10.5 m. From DIETL & SCHWEIGERT (2004).

15 degrees. In other outcrops the submarine relief can reach until 45 degrees. The western border of the lagoon consists of sponge/microbial mounds. They are exposed directly behind the blockhouse. Some years ago there was isolated slumping fold exposed within the laminated facies of this quarry.

Stop 3: Nusplingen Plattenkalk in the Nusplingen quarry

Geological Map 1:25 000 sheet 7819 Meßstetten,
R 3490630, H 5331690

The exploitation of the limestones in the Nusplingen quarry goes back until the middle of the 19th century. Since that time the Nusplingen Plattenkalk has been excavated sporadically for commercial purposes. At the same time the first important fossils were recovered, but lacking any detailed data. The first scientific excavation started in 1929 under the supervision of the University of Tübingen. But it took not much longer than one summer. Two other scientific excavations (1935, 1962) carried out again by the Geological Institute of the University of Tübingen were also restricted to a single summer campaign. The State Museum of Natural History in Stuttgart started new excavations in 1994. Since that time the excavations are going on with great success.

During the new excavations only the uppermost five metres of the Nusplingen Plattenkalk were exposed. The total thickness of this formation within this quarry area is about 10.5 metres. Especially in the upper part of the section some bituminous layers are remarkable in respect of their extraordinary preservation of fossils. They are easy to identify due to their blue-grey colour. Within the succession of the laminated limestones, several turbidites with graded bedding (so-called “allogenic limestones”) and several thicker bioturbated beds indicating a better oxygenation are intercalated. The turbidites yield angular lithoclasts and other components like calcareous ooids, sponge spicules, and even some fragments of hermatypic corals from shallow areas surrounding the lagoon. With the help of these marker beds it is possible to correlate the outcrops in the two quarries. Also some silica layers occur throughout the basin. In the Nusplingen quarry it is visible that the deposition of laminated limestones stopped abruptly. From the nearby sponge/microbial reefs, a lot of boulders had been broken off, gliding into the central parts of the lagoon as olistoliths, where they were deposited directly above the laminated limestone. This olistolith cover reaches a thickness of several meters. The allochthonous nature of these boulders is indicated by geopetal fillings in hollow fossils (e. g. brachiopods) and several ammonites significantly older than those from the underlying laminates. Below the boulder beds, the original stratification of the laminates is strongly disturbed by folding, boudinage, lateral compression, and repetition by stapling. We presume a strong seaquake being the reason for this dramatic event. Because of Cenozoic erosion no data about the younger Upper Jurassic of this area are available.

References

- ARCHER, A. W. & FELDMAN, H. R. (1994): Tidal rhythmites in fine-grained carboniferous limestones, U.S.A. – *Geobios, Mémoire spéciale*, **16**: 275-281.
- BANTEL, G., SCHWEIGERT, G., NOSE, M. & SCHULZ, H.-M. (1999): Mikrofazies, Mikro- und Nannofossilien aus dem Nusplinger Plattenkalk (Ober-Kimmeridgium, Schwäbische Alb). – *Stuttgarter Beiträge zur Naturkunde, B*, **279**: 1-55.
- BRIGGS, D. E. G., MOORE, R., SHULTZ, J. W. & SCHWEIGERT, G. (2005, in press): Mineralization of soft-part anatomy and invading microbes in the horseshoe crab *Mesolimulus* from the Upper Jurassic Lagerstätte of Nusplingen, Germany. – *Proceedings of the Royal Society of London, B*.
- BUATOIS, L. A., MANGANO, M. G. & MAPLES, C. G. (1997): The paradox of nonmarine ichnofaunas in tidal rhythmites: Integrating sedimentologic and ichnologic data from the Late Carboniferous of Eastern Kansas, USA. – *Palaios*, **1997**: 467-481.
- DIETL, G. & SCHWEIGERT, G. (1999): Nusplinger Plattenkalk – Eine tropische Lagune der Jura-Zeit. – *Stuttgarter Beiträge zur Naturkunde, C*, **45**: 1-62.
- DIETL, G. & SCHWEIGERT, G. (2001): Im Reich der Meerengel – Fossilien aus dem Nusplinger Plattenkalk; Munich (Pfeil), 114 pp.
- DIETL, G. & SCHWEIGERT, G. (2004): The Nusplingen Lithographic Limestone – A “Fossil Lagerstaette” of late Kimmerigian age from the Swabian Alb (Germany). – *Rivista Italiana de Paleontologia e Stratigrafia*, **110/1**: 303-309.
- DIETL, G., SCHWEIGERT, G., FRANZ, M. & GEYER, M. (1998): Profile des Nusplinger Plattenkalks (Oberjura, Schwäbische Alb). – *Stuttgarter Beiträge zur Naturkunde, B*, **265**: 1-37.
- FRANZ, M., SELG, M. & MAUS, H. (1997): Der Beuroner Sandstein: eine pliozäne Donauablagerung als Indiz kretazischer Sedimentation in SW-Deutschland. – *Jahreshefte des Geologischen Landesamtes von Baden-Württemberg*, **26**: 125-152.
- GWINNER, M. P. (1976): Origin of the Upper Jurassic Limestones of the Swabian Alb (Southwest-Germany). – *Contributions to Sedimentology*, **5**: 1-69.
- HEISS, G. (1990): Mikrofazies und Diagenese des Malm delta 3/4 im Steinbruch Thiergarten (Donautal). – Unpublished diploma thesis, University of Heidelberg, 56 pp.
- HUMBOLDT, A. V. (1823): *Essai géognostique sur le gisement des roches dans les deux hémisphères*; Paris (Levrault), 379 pp.
- KOCH, R., SENOWBARI-DARYAN, B. & STRAUSS, H. (1994): The Late Jurassic “Massenkalk Fazies” of Southern Germany: Calcareous sand piles rather than organic reefs. – *Facies*, **31**: 179-208.
- PAWELLECK, T. & AIGNER, T. (2003): Apparently homogeneous “reef”-limestones built by high-frequency cycles: Upper Jurassic, SW-Germany. – *Sedimentary Geology*, **160**: 259-284.
- QUENSTEDT, F. A. (1843): *Das Flözgebirge Württembergs. Mit besonderer Rücksicht auf den Jura*; Tübingen (Laupp), 558 pp.
- SCHALLER, J. & KOCH, R. (1996): Der Massenkalk des Schaufelsen im Oberen Donautal. Peloid-Lithoklast-Ooid Krustenkalk oder Schwammriff-Komplex? (Malm delta 4 und Malm epsilon). – *Jahresberichte und Mitteilungen des Oberrheinischen Geologischen Vereins, Neue Folge*, **78**: 309-358.
- SCHWEIGERT, G. (1997): Bibliographie des Nusplinger Plattenkalks von 1823–1997. – *Profil*, **11**: 341-349.
- SCHWEIGERT, G. (1998): Die Ammonitenfauna des Nusplinger Plattenkalks (Ober-Kimmeridgium, Beckeri-Zone, Ulmense-Subzone, Schwäbische Alb). – *Stuttgarter Beiträge zur Naturkunde, B* **267**: 1-61.
- TEMMLER, H. (1966): Über die Nusplinger Fazies des Weißen Jura der Schwäbischen Alb (Württemberg). – *Zeitschrift der Deutschen Geologischen Gesellschaft*, **116**: 891-907.
- TESSIER, B., ARCHER, A. W., LANIER, W. P. & FELDMAN, H. R. (1995): Comparison of ancient tidal rhythmites (Carboniferous of Kansas and Indiana, USA) with modern analogues (the Bay of Mont-Saint-Michel, France). – *Special Publications of the International Association of Sedimentologists*, **24**: 259-271.
- VOLK, H., KOCH, R. & FRANZ, M. (2001): Fazieskartierung in „Massenkalken“ des Weißjura delta 1 bis zeta 1 (ki.2.1 bis ki4). Ein Beispiel aus dem Oberen Donautal (Westliche Schwäbische Alb). – *Geologische Blätter für Nordost-Bayern*, **51**: 247-332.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Zitteliana Serie B](#)

Jahr/Year: 2005

Band/Volume: [26](#)

Autor(en)/Author(s): Schweigert Günter, Dietl Gerd, Koch Roman

Artikel/Article: [Field Trip D The Nusplingen Plattenkalk and Other Fossil Sites in the Western Swabian Alb \(SW Germany\) 87-95](#)