Twenty-one species were enumerated as represented in the collection, of which the following were described as new:—Crocidura Whitakeri, Mus peregrinus, and Lepus atlanticus.—P. L. Sclater, Secretary.

2. Linnean Society of New South Wales.

November 24th, 1897.—1)—3) Botanical.—4) On some Australian Electrinae. Part ii. By J. Douglas Ogilby. Five additional species of Australian electrins are described in this paper, namely—Carassiops (Caulichthys) Guentheri; Carassiops (Austrogobio) Galii, sp.nov.; Krefftius adspersus; Electris fuscus and Ophiorrhinus angustifrons, sp. nov. - 5) On two new Australian Fishes. By J. Douglas Ogilby. The two species described are Harengula stereolepis, from Torres' Straits, and Decapterus leptosomus, a Mackerel-Scad which annually visits Port Jackson but has hitherto escaped notice. - 6) A Contribution to the Zoology of New Caledonia. By J. Douglas Ogilby. After alluding to the meagreness of our knowledge of the biology of the island, the author gives a list of the fresh-water fishes referable to six species brought back by Mr. Hedley, with observations thereon. He considers that the Kuhlia is fairly seperable from the typical rupestris as a subspecies which he proposes to name Hedleyi, while for Gobius crassilabris, a herbivorous form with setaceous pharyngeals, he establishes the new genus Trichopharynx. - 7) On Australian Termitidae. Part iii. By W. W. Froggatt. Eleven species of Termes, of which ten are new, and seven species of Eutermes, of which five are new, are described. In most cases some account of the nests is also given. - 8) On new Marine Mollusca from the Solomon Islands and Australia. By John Brazier, C.M.Z.S. The Volute described at last Meeting from an imperfect specimen is redescribed from a perfect example which subsequently became available. Descriptions also are given of a new Cone from Flinders, Victoria, of two from the Solomon Islands, and of species of Axinaea from the Gippsland Lakes Entrance, Victoria. - 9) Observations on Papuan Land and Freshwater Shells, with Descriptions of new Species from New Guinea and Western Australia. By C. F. Ancey. (Communicated by C. Hedley.) From Western Australia the new species Trachia Froggatti, T. orthocheila and T. monogramma are described and figured from material collected by Mr. W. Froggatt. New Papuan shells from German New Guinea are Sulcobasis leptocochlea and Chloritis Möllendorffi, with which is figured Pupina Beddomei, described in an earlier volume of the Society's Proceedings. Critical observations on sundry Australian and Papuan shells conclude the paper. - Mr. Edgar R. Waite exhibited 1) examples of Typhlops aluensis, Blgr., from Wai Obi, Vuna Pi, Fiji, where they are known to the natives as "Naota." This species was previously known only from the Solomon Islands, and the new record supplies further evidence of the similarity of the faunas of the two Archipelagoes. 2) A New Zealand Fish (Neptotichthys violaceus, Hutton) recently caught in Port Jackson and the first recorded occurrence in Australian waters. 3) Two photographs, taken at Layson Island (Hawaiian Islands); one exhibits an immense concourse of Albatrosses (identified by Mr. A. J. North as Diomedea immutabilis, Rothschild) incubating their eggs.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Zoologischer Anzeiger

Jahr/Year: 1898

Band/Volume: 21

Autor(en)/Author(s): Ogilby Douglas

Artikel/Article: Linnean Society of New South Wales. 72