unter rechtem Winkel abstehend, gerade oder bogenförmig gekrümmt, ihre Enden sind stumpf abgerundet. Achse weich, biegsam, mit einer Spur von Kalk. Coenenchym dünn, mit sehr großen plattenförmigen Spicula in der oberen Schicht, die mit einem langen, stumpf kammförmig gezackten Rand versehen sind, mit dem sie über die Oberfläche hervorragen, und ihr eine rauhe Form verleihen; außerdem sind Spicula von keulenförmiger Gestalt vorhanden. Die innere Coenenchymschicht besteht aus spindel- oder sternförmigen Spicula. Polypen über die ganze Oberfläche vertheilt, vollkommen retractil, als kleine Poren, stellenweise als ganz niedrige Wärzchen über der Oberfläche der Äste markiert.

## 5. On a Peculiarity of the Cerebral Commissures in certain Marsupialia, not hitherto recognised as a Distinctive Feature of the Diprotodontia.

By G. Elliot Smith, M.D., Ch.M., Professor of Anatomy, Egyptian Government School of Medicine, Cairo.

> (Communicated to the Royal Society of London. March 20, 1902.) (With 5 figs.)

> > eingeg. 6. Juni 1902.

It has been known for a considerable time that some of the fibres of the ventral commissure of the cerebrum in certain Marsupials dissociate themselves from the rest of the commissure as soon as they have



Fig. 1. Trichosurus vulpecula. The mesial aspect of the right cerebral hemisphere. > 2.

crossed the mesial plane; and that, instead of passing bodily into the external capsule, which is the usual course of the fibres of the ventral or anterior commissure, they form an aberrant bundle which associates

585

itself with the internal capsule so as to reach the dorsal area of the neopallium by a shorter and slightly less circuitous course (fig. 2).

This peculiarity was represented in the drawings of sections through the brains of *Macropus* and *Phascolomys*, in 1865, by the late W. H. Flower<sup>1</sup>. It was more distinctly shown in a diagram<sup>2</sup> illustrating a coronal section through the brain of a Derbian Wallaby which was published 27 years later by Johnson Symington. Two years later I placed on record the observation upon it, that "in *Phalangista* [*Trichosurus vulpecula*] a bundle of anterior commissure fibres proceeds to the cortex vià the internal capsule, in addition to the external cap-



Fig. 2. Transverse section through the two cerebral hemispheres of the same in the plane a, b (fig. 1). >3. c.d., commissura dorsalis. c.v., commissura ventralis. cp. e., capsula externa. f.a., fasciculus aberrans.

sule"<sup>3</sup>, and in the same place noted an analogous arrangement in various species of *Macropus*.

In 1897 Theodor Ziehen recorded 4 the presence of such fibres in

<sup>&</sup>lt;sup>1</sup> On the Commissures of the Cerebral Hemispheres of the Marsupialia and Monotremata, as compared with those of the Placental Mammals. Phil. Trans., Vol. 155 (1865) p. 633.

<sup>&</sup>lt;sup>2</sup> The Cerebral Commissures in the Marsupialia and Monotremata. Journal of Anatomy and Physiology, Vol. 27, 1892, fig. 3, p. 81.

<sup>&</sup>lt;sup>3</sup> Preliminary Observations on the Cerebral Commissures. Proc. Linn. Soc. of N.S.W., 1894. p. 647-648.

<sup>&</sup>lt;sup>4</sup> Das Centralnervensystem der Monotremen und Marsupialia (Semon's Zoologische Forschungs-Reisen in Australien). Denkschr. Medic.-naturwiss. Gesellsch. Jena. Vol. 6. I.f. II and IV, 1897-1901.

Macropus, Aepyrymnus, and Phascolarctus; but, like Flower and Symington before him, he did not venture on any explanation of them.

The investigations for my memoir of 1894 were carried out chiefly on the brains of Ornithorhynchus, Perameles, Trichosurus, and Macropus. In the Monotreme and the smaller Marsupial (Perameles) the common Mammalian relationship of the ventral commissure to the external capsule was found to obtain, but in the two larger Marsupials some fibres of the ventral commissure were found to pursue the aberrant course indicated above. It was perhaps not unnatural to suppose (as I did in that early attempt at interpreting this peculiarity) that the increased size of the neopallium in Trichosurus and Macropus was wholly responsible for the presence of this aberrant bundle. For it seemed that since the commissural fibres of the neopallium had become too abundant to be wholly accommodated by the path provided by the



Fig. 3. Thylacinus cynocephalus. The mesial aspect of the right cerebral hemisphere. Nat. size.

external capsule, they, so to speak, had overflowed into the internal capsular route.

Upon examining a much larger series of Marsupials than were available when my memoir of 1894 was written, I soon became convinced that the explanation of the causation of this peculiarity which I then suggested could not be regarded as alone sufficient. I found the aberrant bundle in all members of the genera Macropus, Halmaturus, Hypsiprymnus, Dendrolagus, Trichosurus, Petaurus, Phascolarctus, and Phascolomys, quite irrespective of the size of the brain and of the extent of the neopallium. On the other hand, I sought in vain for it in Perameles, Sarcophilus, Dasyurus, Sminthopsis, Didelphys, Myrmecobius, and Notoryctes, even though many of these genera possess larger brains than some of the Diprotodonts.

These facts seemed to suggest that the aberrant bundle was essentially a distinctive feature of the Diprotodont Marcupiala and it or

Marsupials, and it appeared to me that the crucial test of this hypothesis would be afforded by the examination of the brain of Thylacinus, which although that of a Polyprotodont, is almost, if not quite, as large as the brain of the largest Macropod, and considerably larger than those of all other living Diprotodonts. I accordingly submitted the cerebrum of Thylacinus to the test, and found no trace of the aberrant bundle (figs. 3 and 4), wherefore it is



Fig. 4. Transverse section of the two cerebral hemispheres of the same in the plane *a*, *b* (fig. 3). *cp. i*, capsula interna.

clear that the presence of this aberrant fasciculus of the ventral commissure is distinctive of the Diprotodontia.

If we compare the brain of the *Diprotodontia* with that of the other three Mammalian groups: Monotremata, *Polyprotodontia*, and *Eutheria*, the meaning of the aberrant bundle becomes, I believe, fairly obvious.

A study of the structure of the brain in the Monotremes and the Polyprotodont Marsupials shows that in the progenitor of the Mammalia all the commissural fibres of the neopallium must have passed into the ventral commissure vià the external capsule (fig. 4).

The most pronounced growth tendency in the earliest Mammals must have been the enormous increase of the extent of the neopallium, for while at the beginning of the Eocene period this was almost as insignificant as it is in the Reptilia, in most recent Mammals it attains a bulk which far exceeds that of the whole of the rest of the nervous system. This sudden expanse of the neopallium would lead to the development of an enormous mass of fibres which must find some outlet from the pallium. There are only three possible routes for commissural fibres of the neopallium to the mesial plane. There is first of all the external capsule, which chiefly consists in all Mammals of such fibres passing to the ventral commissure: we find the second route in the path mapped out by the internal capsule from the dorsolateral neopallial area to it; and the third route can only involve the invasion of the alveus of the hippocampus.

These three routes, by which a fibre coming from the dorsal neopallium in the region x (fig. 5) may attain the region y in the other hemisphere, are indicated schematically in the following diagram: —

All the neopallial commissural fibres in the *Polyprotodontia* and some only of these in the *Diprotodontia* and *Eutheria* follow the first,



Fig. 5. A scheme of a transverse section in the same plane as that represented in fig. 2 and fig. 4 to show the three routes a, b, and c by which a commissural fibre may pass from the point x in one hemisphere to the region y in the other in different mammalian brains.

which is also the primitive, route (a). The commissural fibres, which spring from the dorso-lateral region of the neopallium in the *Dipro*todontia seem to be crowded out, as it were, of the first route and pursue the second route (b). In the *Eutheria* the neopallial commissural fibres from the dorso-lateral region of the hemisphere forsake both the first and second routes and break through the hippocampal formation (c), or, in other words, invade the alveus so as to form a new dorsally situated neopallial commissure which is the corpus callosum.

This hypothesis of the origin of the corpus callosum I have previously stated in my memoir of 1894 (vide supra) and I discussed it more fully in 1887<sup>5</sup>.

<sup>&</sup>lt;sup>5</sup> The Origin of the Corpus Callosum. Trans. Linn. Soc. of London, 2nd series, Zoology, Vol. 7. part 3. June. 1597. p. 61.

I refer to the matter now, merely to point out that the same determining cause which in the Eutheria calls the "corpus callosum" into being is probably functional in bringing into existence the "aberrant bundle" in the Diprotodontia.

When the relations of these commissural bundles in the four divergent mammalian groups-Monotremata, Polyprotodontia, Diprotodontia, and Eutheria-are carefully studied we are able to appreciate one-and by no means the least-of the reasons why the Eutheria have attained such a pronounced ascendency over the other three groups.

Their brain is that which has retained that particular modification of the commissural arrangement which not only furnishes the shortest and most direct path of communication (c) between the two hemispheres, but also permits of an unimpeded expansion (which is so freely exercised by the corpus callosum). In the other three groups, in which all the neopallial commissural fibres pass through the ventral commissure, the undue expansion of the latter would produce considerable disturbance in the surrounding structures, which in turn would exercise a restraining influence upon any marked increase in size in the commissure itself.

The development of any such commissural mass as the corpus callosum of the more highly organised Mammalia in the position occupied by its homologous fibres (fig. 3, a and b) in the Monotremes and Marsupials would cause the most profound disruptions of the corpus striatum, optic thalamus, and the basal region of the brain, and the complete disorganisation of its whole.

For these various reasons the development of the corpus callosum gives the Eutherian brain a great advantage in the struggle for supremacy, which must have exercised a considerable if not predominant influence in making the Eutheria the highest Mammals.

## II. Mittheilungen aus Museen, Instituten etc.

## 74. Versammlung deutscher Naturforscher und Ärzte in Karlsbad 21.-27. Sept. 1902.

10. Abtheilung: Zoologie, einschl. Entomologie. Einführende: Dr. Nagl (Karlsbad). Prof. Dr. v. Lendenfeld (Prag).

Prof. Dr. Cori (Triest).

Schriftführer: Assist. Freund, Demonstrator Mascha (Prag).

Sitzungslocal: Gymnasium, I. Stock, Tertia.

Frühstücks- und Mittagslocal: »Bayrischer Hof«.

## **ZOBODAT - www.zobodat.at**

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Zoologischer Anzeiger

Jahr/Year: 1901

Band/Volume: 25

Autor(en)/Author(s): Smith G. Elliot

Artikel/Article: On a Peculiarity of the Cerebral Commissures in certain Marsupialia, not hitherto recognised as a Distinctive Feature of the Diprotodontia. 584-589