

Während nach der Sarsschen Gattungsdiagnose die Innenäste des 1. und 4. Paares 2gliedrig, des 2. und 3. Paares 1gliedrig sind, haben wir bei *Parastenocaris* am 1. Fuß einen 2gliedrigen, am 2. Fuß einen 1gliedrigen Innenast. Interessant ist der Bau des Innenastes des 4. Fußes (auch des 3.) bei beiden Gattungen. Bei *Stenocaris* sehen wir zwei langgestreckte Glieder mit einem am Ende gezähnelten Dorn, bei *Parastenocaris* einen Dorn, der zwei Schwellungen besitzt, die ganz sicher beiden Gliedern bei *Stenocaris* entsprechen, und dessen letzter Teil den befiedert ist.

Auch im Bau des 5. Fußes lassen sich Beziehungen finden. *Stenocaris* besitzt zwei starke, breite Dornen nach innen zu, 7 Borsten nach außen zu, *Parastenocaris* einen breiten Dorn und 3 Borsten. Hier ist also die Reduktion am stärksten, sie spricht sich aber deutlich auch an den andern Fußpaaren aus, die man nicht mehr, wenigstens das 3. und 4. Paar, als Schwimmfüße bezeichnen kann, sondern nur als Kriech- oder Stelzfüße, eine Anpassung an ihre Lebensweise in feuchten Moospolstern. Dafür sprechen auch die geraden, starken Apicalborsten der Außenäste.

Wie wir gesehen haben, hat die neue Gattung *Parastenocaris* Beziehungen zu den Gattungen *Cylindropsyllus* Brady und *Stenocaris* Sars; zu der letzteren sind die Beziehungen aber viel größer.

II. Mitteilungen aus Museen, Instituten usw.

Sixth List of Generic names (Birds) under Consideration in Connection with the Official List of Zoological Names.

By Dr. C. W. Stiles.

21) At the Monaco (1913) meeting of the International Zoological Congress, the Secretary of the Commission on Nomenclature was requested to resubmit to special committees the proposed official Lists of Generic Names. This is now being done, and the attention of zoologists is invited to the fact that said lists have been published as follows:

First list, paragraphs 1—3: Ixodoidea; Science, N. Y., Dec. 8, 1911, p. 812; Zool. Anz., Dec. 19, 1911, pp. 589—590.

Second list, paragraphs 4—8: Trematoda; Science, Jan. 26, 1912, p. 146; Zool. Anz., Jan. 3, 1912, pp. 62—63.

Third list, paragraphs 9—14: Cestoda, Nematoda, Gordiacea, and Acanthocephala of man, and rejected names of Trematoda, Nematoda, Gordiacea and Acanthocephala; Science, Mar. 29, 1912, pp. 507—508; Zool. Anz., May 21, 1912, pp. 557—560.

Fourth list, paragraphs 15—18: Diptera; Entomological News, May, 1912, pp. 229—232; Science, May 10, 1912, pp. 751—752; Wiener Ent. Zeitung, June 15, 1912, pp. 155—158; Zool. Anz., June 14, 1912, pp. 653—656.

Fifth list, paragraphs 19—20: Mammals (rejected); Zool. Anz., Dec. 10, 1912, pp. 90—96.

22) Zoologists are cordially invited to reexamine these lists and to send to the Secretary any objections, they may have to the action proposed in connection with said names. Shortly, after January 1, 1914, the Secretary hopes to lay lists 1 to 4 before the Commission for final action, in the form of an Opinion.

23) A special committee of ornithologists, consisting of J. A. Allen (N. Y.), E. Hartert (Tring), C. E. Hellmayr (Münich), H. C. Oberholser (Washington), C. W. Richmond, Secretary (Washington), R. Ridgeway (Washington), L. Stejneger (Washington), W. Stone (Philadelphia), has unanimously agreed upon the following list of names of bird genera as valid and to be accepted:

- Aeryllium* Gray, List Gen. Birds, 1840, 61. Type *Numida vulturina* Hardwicke. (Monotypy and orig. desig.)
- Echmophorus* Coues, Proc. Phil. Acad., 1862, 229. Type *Podiceps occidentalis* Lawrence. (Monotypy and orig. desig.)
- Egithina* Vieillot, Analyse, 1816, 44. Type *Sylvia leucoptera* Vieillot = *Motacilla tiphia* Linn. (Monotypy.)
- Egotheles* Vigors & Horsfield, Tr. Linn. Soc. Lond., XV, i, 1827, 194. Type *Caprimulgus norae-hollandiae* Latham = *C. cristatus* White. (Monotypy.)
- Epyornis* I. Geoffroy St.-Hilaire, Ann. Sci. Nat. (Zool.), (3), XIV, 1851, 209. Type *E. maximus* I. Geoffroy St.-Hilaire. (Monotypy.)
- Aix* Boie, Isis, 1828, 329. Type *Anas sponsa* Linn. (Subs. desig., Eyton, 1838; Gray, 1840.)
- Alauda* Linnaeus, Syst. Nat., ed. 10, I, 1758, 165. Type *Alauda arvensis* Linn. (Subs. desig., Swainson, 1827.)
- Alca* Linnaeus, Syst. Nat., ed. 10, I, 1758, 130. Type *Alca torda* Linn. (Tautonymy, Opinion 16.)
- Alle* Link, Nat.-Samml. Rostock, I, 1806, 17. Type *Alle nigricans* Link = *Alca alle* Linn. (Monotypy.)
- Amandara* Blyth, in White, Nat. Hist. Selborne, 1836, 44. Type *Amandara punctata* Blyth = *Fringilla amandara* Linn. (Monotypy.)
- Anas* Linnaeus, Syst., Nat., ed. 10, I, 1758, 122. Type *Anas boschas* Linn. = *A. platyrhynchos* Linn. (Subs. desig., Lesson, 1828.)
- Anhima* Brisson, Orn., I, 1760, 48 (Cf. V, 518). Type "*Anhima*" = *Palamcdea cornuta* Linn., 1766. (Monotypy and tautonymy.)
- Anhinga* Brisson, Orn., I, 1760, 60 (Cf. VI, 476). Type "*Anhinga*" = *Plotus anhinga* Linn., 1766. (Monotypy and tautonymy.)
- Anser* Brisson, Orn., I, 1760, 58 (Cf. VI, 261). Type "*Anser domesticus*" = *Anas anser* Linn. (Tautonymy.)
- Apaloderma* Swainson, Zool. Illustr., (2), III, 1832, pl. 107. Type *Trogon narina* Stephens. (Orig. desig., and monotypy.)
- Aptenodytes* Miller, Various Subjects Nat. Hist., Nr. 4, 1778, pl. 23. Type *Aptenodytes patagonica* Miller. (Monotypy.)

- Apteryx* Shaw, Nat. Misc., XXIV, 1813, pll. 1057—1058. Type *Apteryx australis* Shaw. (Monotypy.)
- Aquila* Brisson, Orn., I, 1760, 28 (Cf. p. 419). Type "*Aquila*" = *Falco chrysaetos* L. (Tautonymy; also subs. desig., Jameson, 1838.)
- Aramus* Vieillot, Analyse, 1816, 58. Type *Ardea scolopacea* Gmelin. (Monotypy.)
- Ardea* Linnaeus, Syst. Nat., ed. 10, I, 1758, 141. Type *Ardea cinerea* Linn. (Subs. desig., Gray, 1840.)
- Arenaria* Brisson, Orn., I, 1760, 48 (Cf. V, 132). Type "*Arenaria*" = *Tringa interpres* Linn. (Tautonymy.)
- Argusianus* Gray (ex Rafinesque), Cat. Gen. and Subgen. Birds, 1855, 103. Type *Phasianus argus* Linn. (Monotypy.)
- Asio* Brisson, Orn., I, 1760, 28 (Cf. p. 477). Type "*Asio*" = *Strix otus* Linn. (Tautonymy.)
- Astrapia* Vieillot, Analyse, 1816, 36. Type *Paradisea nigra* Gmelin. (Monotypy.)
- Astrarchia* Meyer, Zeitschr. Ges. Orn., II, 1885, 378. Type *A. stephaniac* Meyer. (Monotypy.)
- Asturina* Vieillot, Analyse, 1816, 24. Type *Asturia cinerea* Vieillot = *Falco nitidus* Latham. (Monotypy.)
- Aulacorhynchus* Gould, Mon. Ramphast., 1834, Addenda to Introd. (Cf. also text to pl. of *Pter. sulcatus*.) Type *Pteroglossus sulcatus* Swainson. (Orig. desig.)
- Balaeniceps* Gould, Proc. Zool. Soc. Lond., 1852, 1. Type *B. rex* Gould. (Monotypy.)
- Balcarica* Brisson, Orn., I, 1760, 48 (Cf. V, 511). Type "*Balearica*" = *Ardea paronina* Linn. (Monotypy.)
- Batrachostomus* Gould, Icones Avium, Pt. II, 1838, pl. xvii. Type *Podargus auritus* Gray or Vigors. (Monotypy.)
- Branta* Scopoli, Annus I, 1769, 67. Type *Anas bernicla* Linn. (Subs. desig., Bannister, 1870.)
- Brotogeris* Vigors, Zool. Journ., II, 1825, 400. Type *Psittacus pyrrhopterus* Latham. (Monotypy.)
- Bubo* Duméril, Zool. Analyt., 1806, 34. Type "*Les Ducs*" = *Strix bubo* Linn. (Monotypy; taut., and subs. desig., Froriep, 1806; Opinion 46.)
- Bucco* Brisson, Orn., I, 1760, 42 (Cf. IV, 91). Type "*Bucco*" = *Bucco capensis* Linn., 1766. (Tautonymy.)
- Buphagus* Brisson, Orn., I, 1760, 32 (Cf. II, 436). Type "*Buphagus*" = *Buphaga africana* Linn., 1766. (Monotypy.)
- Burhinus* Illiger, Prodrömus, 1811, 250. Type *Charadrius magnirostris* Latham = *C. grallarius* Latham. (Monotypy.)
- Cairina* Fleming, Philos. Zoology, II, 1822, 260. Type *Anas moschata* Linn. (Monotypy.)
- Campephaga* Vieillot, Analyse, 1816, 39. Type *C. nigra* Vieillot = *C. flava* Vieillot. (Monotypy.)
- Capito* Vieillot, Analyse, 1816, 27. Type *Bucco niger* Müller. (Monotypy.)
- Caprimulgus* Linnaeus, Syst. Nat., ed. 10, I, 1758, 193. Type *C. europaeus* Linn. (Tautonymy; Opinion 16.)
- Carduelis* Brisson, Orn., I, 1760, 36 (Cf. III, 53). Type "*Carduelis*" = *Fringilla carduelis* Linn. (Tautonymy.)
- Cariama* Brisson, Orn., I, 1760, 48 (Cf. V, 516). Type "*Cariama*" = *Palamedea cristata* Linn., 1766. (Monotypy.)
- Casuarinus* Brisson, Orn., I, 1760, 46 (Cf. V, 10). Type "*Casuarinus*" = *Struthio casuarinus* Linn. (Monotypy.)
- Cathartes* Illiger, Prodrömus, 1811, 236. Type *Vultur aura* Linn. (Subs. desig., Vigors, 1825; Swainson, 1836.)
- Centrocercus* Swainson, Fauna Bor.-Amer., II, 1831 (1832), 358, 496. Type *Tetrao urophasianus* Bonaparte. (Monotypy.)

- Cephalopterus* Geoffroy St.-Hilaire, Ann. Mus. Hist. Nat., XIII, 1809, 238. Type *C. ornatus* Geoffroy St. Hilaire. (Monotypy.)
- Cepphus* Pallas, Spic. Zool., Fasc. 5, 1769, 33. Type *C. lactcolus* Pallas = *Alca grylle* Linn. (Monotypy.)
- Cereopsis* Latham, Suppl. Ind. Orn., 1801, lxvii. Type *C. norae hollandiae* Latham. (Monotypy.)
- Certhia* Linnaeus, Syst. Nat., ed. 10, I, 1758, 118. Type *Certhia familiaris* Linn. (Tautonymy, Opinion 16; also subs. desig., Jardine, 1839.)
- Charadrius* Linnaeus, Syst. Nat., ed. 10, I, 1758, 150. Type *C. hiaticula* Linn. (Tautonymy, Opinion 16.)
- Chauna* Illiger, Prodromus, 1811, 253. Type *Parra chacaria* Linn. (Monotypy.)
- Chrysolophus* Gray, Illustr. Ind. Zool., II, 1833—34, pl. 41, f. 2. Type *Phasianus pictus* Linn. (Monotypy.)
- Chunga* Burmeister, Proc. Zool. Soc. Lond., 1860, 335. Type *Dicholophus burmeisteri* Hartlaub. (Monotypy.)
- Ciccinnurus* Vieillot, Analyse, 1816, 35. Type *Paradisaea regia* Linn. (Monotypy.)
- Ciconia* Brisson, Orn., I, 1760, 48 (Cf. V. 361). Type "*Ciconia alba*" = *Ardea ciconia* Linn. (Tautonymy.)
- Cinclus* Borkhausen, Deutsche Fauna, I, 1797, 300. Type *Cinclus hydrophilus* Borkh. = *Sturnus cinclus* Linn. (Monotypy and taut.)
- Circæus* Vieillot, Analyse, 1816, 23. Type *Falco gallicus* Gmelin. (Monotypy.)
- Clamator* Kaup, Nat. Syst., 1829, 53. Type *Cuculus glandarius* Linn. (Monotypy.)
- Coccothraustes* Brisson, Orn., I, 1760, 36 (Cf. III, 218). Type "*Coccothraustes*" = *Loxia coccothraustes* Linn. (Tautonymy.)
- Coccyzus* Vieillot, Analyse, 1816, 28. Type *Cuculus americanus* Linn. (Monotypy.)
- Cochlearius* Brisson, Orn., I, 1760, 48 (Cf. V, 506). Type "*Cochlearius*" = *Canceroma cochlearia* Linn., 1766. (Monotypy and taut.)
- Coereba* Vieillot, Ois. Amér. Sept., II, 1807 (1809?), 70. Type *Certhia flareola* Linn. (Monotypy.)
- Colaptes* (Swainson MS.) Vigors, Trans. Linn. Soc. Lond., XIV, iii, 1825, 457. Type *Cuculus auratus* Linn. (Monotypy.)
- Colius* Brisson, Orn., I, 1760, 36 (Cf. III, 304). Type "*Colius capitis bonae spei*" = *Loxia colius* Linn., 1766. (Tautonymy.)
- Colluricincla* Vigors & Horsfield, Trans. Linn. Soc. Lond., XV, 1827, 213. Type *C. cinerea* Vigors & Horsfield. (Monotypy.)
- Columba* Linnaeus, Syst. Nat., ed. 10, I, 1758, 162. Type *C. oenas* Linn. (Subs. desig., Vigors, 1825.)
- Colymbus* Linnaeus, Syst. Nat., ed. 10, I, 1758, 135. Type *C. cristatus* Linn. (Subs. desig., Baird, Brewer & Ridgway, 1884.)
- Corvus* Linnaeus, Syst. Nat., ed. 10, I, 1758, 105. Type *Corvus corax* Linn. (Tautonymy, Opinion 16.)
- Cotinga* Brisson, Orn., I, 1760, 32 (Cf. II, 339). Type "*Cotinga*" = *Ampelis cotinga* Linn., 1766. (Tautonymy.)
- Coturnix* Bonnaterræ, Tabl. Enc. Méth., I, 1791, lxxxvii; 1792, 216. Type *C. communis* Bonn. = *Tetrao coturnix* Linn. (Tautonymy.)
- Crotophaga* Linnaeus, Syst. Nat., ed. 10, I, 1758, 105. Type *C. ani* Linn. (Monotypy.)
- Cuculus* Linnaeus, Syst. Nat., ed. 10, I, 1858, 110. Type *C. canorus* Linn. (Tautonymy, Opinion 16; also subs. desig., Swainson, 1837.)
- Diomedea* Linnaeus, Syst. Nat., ed. 10, I, 1758, 132. Type *D. exulans* Linn. (Subs. desig., Gray, 1840.)
- Dromas* Paykull, K. Vet. Akad. Nya Handl., XXVI, 1805, 188. Type *Dromas ardeola* Paykull. (Monotypy.)
- Dromiceius* Vieillot, Analyse, 1816, 54. Type *Casuarinus norae hollandiae* Latham. (Monotypy.)
- Ectopistes* Swainson, Zool. Journ., III, 1827, 362. Type *Columba migratoria* Linn. (Only certain species, and subs. desig., same vol., p. 614.)

- Egretta* Forster, Synop. Cat. Brit. Birds, 1817, 59. Type *Ardea garzetta* Linn. (Monotypy.)
- Elanus* Savigny, Descr. Égypte, I, 1809, 69, 97. Type *Elanus caesi* Savigny = *Falco melanopterus* Daudin. (Monotypy.)
- Eurynorhynchus* Nilsson, Orn. Suecica, II, 1821, 29. Type *E. griseus* Nilsson = *Platalia pygmaea* Linn. (Monotypy.)
- Eurylaimus* Horsfield, Trans. Linn. Soc. Lond., XIII, i, 1821, 170. Type *Eurylaimus javanicus* Horsfield. (Monotypy.)
- Eurypyga* Illiger, Prodr. Rom., 1811, 257. Type *Ardea helias* Pallas. (Monotypy.)
- Fratereula* Brisson, Orn., I, 1760, 52 (Cf. VI, 81). Type "*Fratereula*" = *Alca arctica* Linn. (Monotypy and tautonymy.)
- Fringilla* Linnaeus, Syst. Nat., ed. 10, I, 1758, 179. Type *F. coelebs* Linn. (Tautonymy, Opinion 16.)
- Fulica* Linnaeus, Syst. Nat., ed. 10, I, 1758, 152. Type *F. atra* Linn. (Tautonymy, Opinion 16.)
- Fulmarus* Stephens, Shaw's General Zool., XIII, i, 1826, 233. Type *Procellaria glacialis* Linn. (Subs. desig., Palmer, 1836.)
- Gallinago* Koch, Syst. Baier. Zool., 1816, 312. Type *G. media* of Koch = *Scolopax gallinago* Linn. (Tautonymy.)
- Gallinula* Brisson, Orn., I, 1760, 50 (Cf. VI, 2). Type "*Gallinula*" = *Fulica chloropus* Linn. (Tautonymy.)
- Gallus* Brisson, Orn., I, 1760, 26 (Cf. p. 164). Type "*Gallus domesticus*" = *Phasianus gallus* Linn. (Tautonymy.)
- Gampsonyx* Vigors, Zool. Journ., II, 1825, 69. Type *G. swainsonii* Vigors. (Monotypy.)
- Glarcola* Brisson, Orn., I, 1760, 48 (Cf. V, 141). Type "*Glarcola*" = *Hirundo pratincola* Linn., 1766. (Tautonymy.)
- Goura* Stephens, Shaw's General Zool., XI, i, 1819, 119. Type *Columba coronata* Linn., 1766 = *C. cristata* Pallas, 1764. (Subs. desig., Gray, 1840.)
- Gypæus* Storr, Alpenreise, I, 1784, 69. Type *G. grandis* Storr = *Vultur barbatus* Linn. (Monotypy.)
- Haematopus* Linnaeus, Syst. Nat., ed. 10, I, 1758, 152. Type *H. ostralegus* Linn. (Monotypy.)
- Haliæctus* Savigny, Descr. Égypte, I, 1809, 68, 85. Type *H. nesus* Savigny = *Falco albicilla* Linn. (Monotypy.)
- Haliastur* Selby, Cat. Gen. and Subgen. Types Birds, 1840, 3. Type "*F. pondicerianus* auct." = *Falco indus* Bodd. (Monotypy and orig. desig.)
- Heliornis* Bonnatere, Tabl. Enc. Méth., I, 1791, lxxxiv, 65. Type *Heliornis fulicarius* Bonn. = *Colymbus fulica* Bodd. (Monotypy.)
- Himantopus* Brisson, Orn., I, 1760, 46 (Cf. V, 33). Type "*Himantopus*" = *Charadrius himantopus* Linn. (Tautonymy.)
- Ibidorhyncha* Vigors, Proc. Comm. Sci. Zool. Soc. Lond., I, 1832, 174. Type *I. struthersi* Vigors. (Monotypy.)
- Jynx* Linnaeus, Syst. Nat., ed. 10, I, 1758, 112. Type *J. torquilla* Linn. (Monotypy.)
- Lagopus* Brisson, Orn., I, 1760, 26 (Cf. p. 181). Type "*Lagopus*" = *Tetrao lagopus* Linn. (Tautonymy.)
- Lanius* Linnaeus, Syst. Nat., ed. 10, I, 1758, 93. Type *Lanius excubitor* Linn. (Subs. desig., Swainson, 1824.)
- Leistes* Vigors, Zool. Journ., II, 1825, 191. Type *Oriolus americanus* Gmelin = *Emberiza militaris* Linn. (Orig. desig.)
- Loxia* Linnaeus, Syst. Nat. ed. 10, I, 1758, 171. Type *Loxia curvirostra* Linn. (Tautonymy, Opinion 16.)
- Lunda* Pallas, Zoogr. Rosso-Asiatica, II, 1811 (1826?), 363. Type *Alca cirrhata* Pallas. (Subs. desig., Gray, 1840.)
- Manucodia* Boddart, Table Pl. Enl., 1783, 39. Type *M. chalybea* Bodd. = *Paradisca chalybeata* Forster, 1781. (Monotypy.)

- Mcleagris* Linnaeus, Syst. Nat., ed. 10, I, 1758, 156. Type *M. gallopavo* Linn. (Tautonymy, Opinion 16.)
- Mergus* Linnaeus, Syst. Nat., ed. 10, I, 1758, 129. Type *Mergus merganser* Linn. (Subs. desig., Gray, 1840.)
- Merops* Linnaeus, Syst. Nat., ed. 10, I, 1758, 117. Type *Merops apiaster* Linn. (Tautonymy, Opinion 16; also Subs. desig., Swainson, 1820.)
- Momotus* Brisson, Orn., I, 1760, 44 (Cf. IV, 465). Type "*Momotus*" = *Ramphastos momota* Linn., 1766. (Tautonymy.)
- Motacilla* Linnaeus, Syst. Nat., ed. 10, I, 1758, 184. Type *M. alba* Linn. (Tautonymy, Opinion 16.)
- Musophaga* Isert, Schriften Gesellsch. nat. Freunde Berlin, IX, 1788, 17. Type *M. violacea* Isert. (Monotypy.)
- Neophron* Savigny, Descr. Égypte, I, 1809, 68, 75. Type *Vultur perenopterus* Linn. (Monotypy.)
- Notornis* Owen, Trans. Zool. Soc. Lond., III, v, 1848, 366. Type *Notornis mantelli* Owen. (Monotypy.)
- Numenius* Brisson, Orn., I, 1760, 48 (Cf. V, 311). Type "*Numenius*" = *Scelopax arquatus* Linn. (Tautonymy.)
- Numida* Linnaeus, Mus. Ad. Frid., 1764, 27. Type *Phasianus mcleagris* Linn., 1758. (Monotypy.)
- Nucifraga* Brisson, Orn., I, 1760, 30 (Cf. II, 58). Type "*Nucifraga*" = *Corvus caryocatactes* Linn. (Tautonymy and Monotypy.)
- Nyctea* Stephens, Shaw's General Zool., XIII, ii, 1826, 62. Type *Strix erminea* Shaw = *S. nyctea* Linn. (Tautonymy; only certain species; also monotypy.)
- Oedienemus* Temminck, Man. d'Orn., 1815, 321. Type *Æ. crepitans* Temm. = *Charadrius oedienemus* Linn. (Monotypy and tautonymy.)
- Opisthocomus* Illiger, Prodrömus, 1811, 239. Type *Phasianus cristatus* Gmelin = *P. hoazin* Müller. (Monotypy.)
- Oriolus* Linnaeus, Syst. Nat., ed. 12, I, 1766, 160. Type *Oriolus galbula* Linn. = *Coracias oriolus* Linn., 1758. (Tautonymy.)
- Otis* Linnaeus, Syst. Nat., ed. 10, I, 1758, 154. Type *Otis tarda* Linn. (Tautonymy, Opinion 16, also Subs. desig., Jardine & Selby, 1828.)
- Otus* Pennant, Indian Zool., 1769, 3. Type *Otus bakkamoena* Pennant. (Monotypy.)
- Pachycephala* Vigors, Trans. Linn. Soc. Lond., XIV, iii, 1825, 444. Type *Muscicapa pectoralis* Latham = *Turdus gutturalis* Latham. (Monotypy and Orig. desig.)
- Pandion* Savigny, Descr. Égypte, I, 1809, 69, 95. Type *Pandion fluvialis* Savigny = *Falco haliastur* Linn. (Monotypy.)
- Paradisaea* Linnaeus, Syst. Nat., ed. 10, I, 1758, 110. Type *P. apoda* Linn. (Tautonymy, Opinion 16; also Subs. desig., Gray, 1840.)
- Parotia* Vieillot, Analyse, 1816, 35. Type *Paradisaea sciflata* Forster, 1781. (Monotypy.)
- Parus* Linnaeus, Syst. Nat., ed. 10, I, 1758, 189. Type *Parus major* Linn. (Subs. desig., Gray, 1840.)
- Pavo* Linnaeus, Syst. Nat., ed. 10, I, 1758, 156. Type *P. cristatus* Linn. (Tautonymy, Opinion 16.)
- Pelecanus* Linnaeus, Syst. Nat., ed. 10, I, 1758, 132. Type *Pelecanus onocrotalus* Linn. (Tautonymy, Opinion 16.)
- Perdix* Brisson, Orn., I, 1760, 26 (Cf. p. 219). Type "*Perdix cinerea*" = *Tetrao perdix* Linn. (Tautonymy.)
- Pezoporus* Illiger, Prodrömus, 1811, 201. Type *Psittacus formosus* Latham = *P. terrestris* Shaw. (Monotypy.)
- Phæthön* Linnaeus, Syst. Nat., ed. 10, I, 1758, 134. Type *P. aethereus* Linn. (Subs. desig., Gray, 1840.)

- Phalacrocorax* Brisson, Orn., I, 1760, 60 (Cf. VI, 511). Type "*Phalacrocorax*" = *Pelicanus carbo* Linn. (Tautonymy.)
- Phalaropus* Brisson, Orn., I, 1760, 50 (Cf. VI, 12). Type "*Phalaropus*" = *Tringa fulicaria* Linn. (Tautonymy.)
- Pharomachrus* de la Llave, Registro Trimestre, I, 1832, 48. Type *P. mocinno* de la Llave. (Monotypy.)
- Phasianus* Linnaeus, Syst. Nat., ed. 10, I, 1758, 158. Type *P. colehius* Linn. (Tautonymy, Opinion 16.)
- Phoenicopterus* Linnaeus, Syst. Nat., ed. 10, I, 1758, 139. Type *P. ruber* Linn. (Monotypy.)
- Picus* Linnaeus, Syst. Nat., ed. 10, I, 1758, 112. Type *P. viridis* Linn. (Subs. desig., Swainson, 1820.)
- Platalea* Linnaeus, Syst. Nat., ed. 10, I, 1758, 139. Type *P. leucorodia* Linn. (Subs. desig., Gray, 1840.)
- Platycercus* Vigors, Zool. Journ., I, 1825, 527. Type *Psittacus pennanti* Latham = *P. elegans* Gmelin. (Orig. desig.)
- Plautus* Brünnich, Zool. Fund., 1771, 78. Type "*Brillefuglen*" = *Alca impennis* Linn. (Monotypy.)
- Pluvialis* Brisson, Orn., I, 1760, 46 (Cf. V, 42). Type "*Pluvialis aurea*" = *Charadrius apricarius* Linn. (Tautonymy.)
- Pluvianus* Vieillot, Analyse, 1816, 55. Type *Charadrius aegyptius* Linn. (Monotypy.)
- Polyplectron* Temminck, Cat. Syst., 1807, 149. Type *Polyplectron argus* Temm. = *Pavo bicalcaratus* Linn. (Monotypy.)
- Porphyrio* Brisson, Orn., I, 1760, 48 (Cf. V, 522). Type "*Porphyrio*" = *Fulica porphyrio* Linn. (Tautonymy.)
- Porzana* Vieillot, Analyse, 1816, 61. Type *Rallus porzana* Linn. (Monotypy.)
- Psittacus* Linnaeus, Syst. Nat., ed. 10, I, 1758, 96. Type *P. erithacus* Linn. (Subs. desig., Gray, 1840.)
- Psophia* Linnaeus, Syst. Nat., ed. 10, I, 1758, 154. Type *P. crepitans* Linn. (Monotypy.)
- Pteroglossus* Illiger, Prodrum, 1811, 202. Type *Ramphastos aracari* Linn. (Subs. desig., Swainson, 1821.)
- Ptiloris* Swainson, Zool. Journ., I, 1825, 479. Type *P. paradiseus* Swainson. (Monotypy.)
- Puffinus* Brisson, Orn., I, 1760, 56 (Cf. VI, 130). Type "*Puffinus*" = *Procellaria puffinus* Brünnich, 1764. (Tautonymy.)
- Querula* Vieillot, Analyse, 1816, 37. Type *Muscicapa cruenta* Bodd. = *M. purpurata* Müller. (Monotypy.)
- Rallus* Linnaeus, Syst. Nat., ed. 10, I, 1758, 153. Type *R. aquaticus* Linn. (Subs. desig., Fleming, 1821.)
- Raphus* Brisson, Orn., I, 1760, 46 (Cf. V, 14). Type "*Raphus*" = *Struthio cucullatus* Linn. (Monotypy.)
- Recurvirostra* Linnaeus, Syst. Nat., ed. 10, I, 1758, 151. Type *R. arosetta* Linn. (Monotypy.)
- Rhea* Brisson, Orn., I, 1760, 46 (Cf. V, 8). Type "*Rhea*" = *Struthio americanus* Linn. (Monotypy.)
- Rynchops* Linnaeus, Syst. Nat., ed. 10, I, 1758, 138. Type *R. nigra* Linn. (and *R. fulca* = *nigra*). (Monotypy.)
- Sarcoramphus* Duméril, Zool. Analytique, 1806, 32. Type *Vultur papa* Linn. (Cf. Froriep, Anal. Zool., 1806, 33). (Subs. desig., Vigors, 1825.)
- Scolopax* Linnaeus, Syst. Nat., ed. 10, I, 1758, 145. Type *S. rusticola* Linn. (Tautonymy, Opinion 16.)
- Scopus* Brisson, Orn., I, 1760, 48 (Cf. V, 503). Type "*Scopus*" = *S. umbretta* Gmelin, 1789. (Monotypy.)

- Sericulus* Swainson, Zool. Journ., I, 1825, 476. Type *Meliphaga chrysocephala* Lewin. (Monotypy.)
- Sitta* Linnaeus, Syst. Nat., ed. 10, I, 1758, 115. Type *Sitta europaea* Linn. (Monotypy.)
- Somateria* Leach, Annals of Philos., XIII, 1819, 61. Type *Anas mollissima* Linn. (Monotypy.)
- Sphenorhynchus* (Hempr. & Ehrenb. MS.) Lichtenstein, Verz. Doubl., 1823, 76. Type *Ciconia abdimii* Licht. (Monotypy.)
- Spindalis* Jardine & Selby, Ill. Orn., N. S., 1837, pl. IX. Type *S. bilineatus* J. & S. = *Tanagra nigricephala* Jameson. (Monotypy.)
- Stercorarius* Brisson, Orn., I, 1760, 56 (Cf. VI, 149). Type "*Stercorarius*" = *Larus parasiticus* Linn. (Tautonymy.)
- Sterna* Linnaeus, Syst. Nat., ed. 10, I, 1758, 137. Type *S. hirundo* Linn. (Tautonymy, Opinion 16.)
- Strigops* Gray, Gen. Birds, II, 1845, 426. Type *S. habroptilus* Gray. (Monotypy.)
- Strix* Linnaeus, Syst. Nat., ed. 10, I, 1758, 92. Type *S. stridula* Linn. = *S. aluco* Linn. (Tautonymy, Opinion 16.)
- Struthio* Linnaeus, Syst. Nat., ed. 10, I, 1758, 155. Type *S. camelus* Linn. (Tautonymy, Opinion 16.)
- Sturnella* Vieillot, Analyse, 1816, 34. Type *Alanda magna* Linn. (Monotypy.)
- Sturnus* Linnaeus, Syst. Nat., ed. 10, I, 1758, 167. Type *Sturnus vulgaris* Linn. (Tautonymy, Opinion 16; also Subs. desig., Stephens, 1817.)
- Sula* Brisson, Orn., I, 1760, 60 (Cf. VI, 494). Type "*Sula*" = *Pelecanus piscator* Linn. (Tautonymy.)
- Surnia* Duméril, Zool. Analytique, 1806, 34. Type *Strix caparoch* Müller. (Subs. desig., Froriep, 1806, and Monotypy.)
- Syrrhaptes* Illiger, Prodromus, 1811, 243. Type *Tetrao paradoxa* Pallas. (Monotypy.)
- Tachyphonus* Vieillot, Analyse, 1816, 33. Type *Tanagra rufa* Boddaert. (Monotypy.)
- Talgalla* Lesson, Man. d'Orn., II, 1828, 185. Type *T. cucieri* Less. (Monotypy.)
- Thamnophilus* Vieillot, Analyse, 1816, 40. Type *Lanius dolius* Linn. (Subs. desig., Swainson, 1824.)
- Todus* Brisson, Orn., I, 1760, 44 (Cf. IV, 528). Type "*Todus*" = *Alcedo todus* Linn. (Tautonymy.)
- Trichoglossus* (Vigors MS.) Stephens, General Zool., XIV, i, 1826, 129. Type *T. swainsoni* Jardine & Selby, 1831 = *haematopus* Stephens (nec *haematodus* Linn.) = *Psittacus novae hollandiae* Gmelin. (Subs. desig., Swainson, 1832.)
- Tringa* Linnaeus, Syst. Nat., ed. 10, I, 1758, 148. Type *Tringa oerophus* Linn. (Tautonymy, Opinion 16.)
- Troglodytes* Vieillot, Ois. Amér. Sept., II, 1807 (1809?), 52. Type *T. acdon* Vieillot. (Subs. desig., Baird, 1858.)
- Trogon* Brisson, Orn., I, 1760, 42 (Cf. IV, 164). Type "*Trogon cayanensis cinereus*" = *T. strigilatus* Linn., 1766. (Subs. desig., Allen, 1907.)
- Turdus* Linnaeus, Syst. Nat., ed. 10, I, 1758, 168. Type *T. viscivorus* Linn. (Subs. desig., Gray, 1840.)
- Upupa* Linnaeus, Syst. Nat., ed. 10, I, 1758, 117. Type *U. epops* Linn. (Tautonymy, Opinion 16.)
- Uratelornis* Rothschild, Novit. Zool., II, 1895, 479. Type *U. chimaera* Rothschild. (Orig. desig., and Monotypy.)
- Vanga* Vieillot, Analyse, 1816, 41. Type *Lanius curvirostris* Linn. (Monotypy.)
- Virco* Vieillot, Ois. Amér. Sept., I, 1807 (1808?), 83. Type *V. musicus* Vieillot = *Tanagra grisea* Bodd. (Subs. desig., Gray, 1840.)
- Vultur* Linnaeus, Syst. Nat., ed. 10, I, 1758, 86. Type *V. gryphus* Linn. (Subs. desig., Allen, 1907.)

24) Zoologists are cordially invited to examine the list and to submit to the Secretary of the Commission, before Jan. 1, 1914, any objection they have to the incorporation of any of the foregoing names in the official (accepted) list. Unless circumstances indicate that delay is advisable, the Secretary will call for the vote of the Commission early in 1914.

25) The permanent address of the Secretary is Hygienic Laboratory 25th & E. Sts., N. W., Washington, D. C. His temporary address for 1913 and 1914 is U. S. Marine Hospital, Wilmington, N. C. A letter directed at any time to either address will reach him.

III. Personal-Notizen.

J. S. Kingsley of Tufts College goes to the University of Illinois at Urbana, Ill. as Professor of Zoology in charge of Vertebrates.



ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Zoologischer Anzeiger](#)

Jahr/Year: 1913

Band/Volume: [42](#)

Autor(en)/Author(s): Stiles Charles Wardell

Artikel/Article: [Mitteilungen aus Museen, Instituten usw. 520-528](#)