

# Distribution pattern and phylogeography of tree rats *Chiromyscus* (Rodentia, Muridae) in eastern Indochina

Alexander E. Balakirev<sup>1,2</sup>, Alexei V. Abramov<sup>1,3</sup>, Viatcheslav V. Rozhnov<sup>1,2</sup>

<sup>1</sup> Joint Russian-Vietnamese Tropical Research and Technological Centre, 63 Nguyen Van Huyen, Nghia Do, Cau Giay, Hanoi, Vietnam

<sup>2</sup> A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Leninskii pr. 33, Moscow 119071, Russia

<sup>3</sup> Zoological Institute, Russian Academy of Sciences, Universitetskaya nab. 1, Saint Petersburg 199034, Russia

<http://zoobank.org/DD5BE6F7-403E-4E16-BA21-D23F7C06CF7A>

Corresponding author: Alexander E. Balakirev ([alexbalakirev@mail.ru](mailto:alexbalakirev@mail.ru))

Academic editor: M.T.R. Hawkins ♦ Received 11 August 2020 ♦ Accepted 12 January 2021 ♦ Published 3 February 2021

## Abstract

The study combines available data on species distribution in eastern Indochina to investigate the phylogeographical genetic and morphological diversity of tree rats (*Chiromyscus*, Rodentia, Muridae) and to specify their natural ranges. We examined the diversity and distribution of tree rats over its range, based on recent molecular data for mitochondrial (*Cyt b*, *COI*) and nuclear (*IRBP*, *RAG1* and *GHR*) genes. The study presents the most complete and up-to-date data on the distribution and phylogeography of the genus in eastern Indochina. As revealed by mitochondrial genes, *C. langbianis* splits into at least four coherent geographically-distributed clades, whereas *C. thomasi* and *C. chiropus* form two distinctive mitochondrial clades each. *Chiromyscus langbianis* and *C. chiropus* show significant inconsistency in nuclear genes, whereas *C. thomasi* shows the same segregation pattern as can be traced by mitochondrial markers. The Northern and Southern phylogroups of *C. thomasi* appear to be distributed sympatrically with northern phylogroups of *C. langbianis* in most parts of eastern Indochina. The mitochondrial clades discovered are geographically subdivided and divergent enough to suspect independent subspecies within *C. langbianis* and *C. thomasi*. However, due to the insufficiency of obvious morphological traits, a formal description is not carried out here. The processes of recent fauna formation, species distribution patterns, dispersion routes and possible natural history in Indochina are discussed.

## Key Words

biodiversity, Indochina, Southeast Asia, taxonomy, tree rats, Vietnam

## Introduction

Genus *Chiromyscus* Thomas, 1925 is currently assigned to the *Dacnomys* division of the tribe Rattini (Musser and Carleton 1993, 2005). It was first described, based on *Mus chiropus* (Thomas, 1891) from East Burma (= Myanmar) and for a long time was considered monotypic. Its close relationships with *Niviventer langbianis* (Robinson, Kloss, 1922) and *N. cremoriventer* (Miller, 1900) were initially suspected and discussed by Musser (Musser 1973, 1981; Musser and Carleton 1993, 2005). The genus was recently re-established by Balakirev et al. (2014) as comprising at least three recent species: *C. chiropus*

(Thomas, 1891), *C. langbianis* and *C. thomasi* Balakirev, Abramov & Rozhnov, 2014.

To date, Thomas' tree rat, *C. thomasi*, is known to be distributed from southwest China (Lan et al. 1994; Chen et al. 1995; Wang 2002) to eastern Myanmar and northern Thailand (Marshall 1977), Vietnam (Dang et al. 1994; Lunde and Nguyen 2001; Can et al. 2008, Balakirev et al. 2014) and Laos (Musser 1981; Corbet and Hill 1992; Aplin et al. 2003; Aplin and Lunde 2016), where it has long been known under the name *C. chiropus*. The Dalat tree rat *C. langbianis* was described from the Dalat Plateau in southern Vietnam (Robinson and Kloss 1922) and is currently recorded throughout Vietnam and Laos

(Musser 1973; Dang et al. 1994; Lunde et al. 2003; Balakirev et al. 2011; Lu et al. 2015) and southern China (Lu et al. 2015), including Hainan Island (Ge et al. 2018). Fea's tree rat *C. chiropus* is known from Myanmar and easternmost India (Musser 1973) and southern Vietnam (Balakirev et al. 2014).

The evolution of the genus *Chiromyscus* was affected by the natural history of this region. Continuous forest cover in Indochina existed during a considerable proportion of the Pliocene and Pleistocene (Meijaard and Groves 2006), enabling the direct contact, dispersion and genetic exchange of western and eastern Indochinese populations, which are currently mostly interrupted by the extensive deforestation in areas of central Thailand and southern Cambodia. During Pleistocene glaciations, the forest edge was located at lower elevations (Cox and Moore 2000) and substantial forest contraction happened during the last glacial periods, as evidenced in Peninsular Malaysia and Palawan (Wurster et al. 2010). Cranbrook (2000) and Gathorne-Hardy et al. (2002) stressed that most areas in this region were covered with savannah vegetation unfavourable to arboreal species during long periods of the Quaternary epoch. However, the most recent surveys on the biogeography and paleoenvironments of the Sunda Shelf have suggested that the interactions between climate and sea level and their effects on the distribution patterns of the fauna and flora are more complex (van den Bergh et al. 2001; Meijaard and Groves 2006) and environmental conditions changed repeatedly during the Pleistocene. These global processes of ecosystems change were likely to have had an impact on the recent genetic structure of *Chiromyscus* species. This group of rodents is still quite rare in museum collections, so little information is available about its natural diversity, distribution and phylogenetic relationships and only a few specimens have been genetically characterised. In this paper, we combine available data, including novel data, on species distribution in eastern Indochina to investigate the phylogeography and diversity of these species, both genetic and morphological and specify their natural ranges.

## Material and methods

### Specimens and samples

A great number of *Chiromyscus* specimens, obtained in Vietnam during a series of field expeditions of the Joint Russian-Vietnamese Tropical Research and Technological Centre between 2007 and 2018, were sampled for genetic analysis in full agreement with current Vietnamese regulations in the field of Nature Protection and Biodiversity Conservation. We followed the guidelines of the American Society of Mammalogists during the collection and handling of the animals used in this survey (Gannon et al. 2011). The museum specimens were kept in the Zoological Museum, Moscow State University, Moscow, Russia (ZMMU) and the Zoological Institute, Russian

Academy of Sciences, Saint-Petersburg, Russia (ZIN); genetic samples are part of collection of Joint Russian–Vietnamese Tropical Research and Technological Centre, Hanoi, Vietnam.


New samples were combined with sequences available in GenBank, including our sequences previously submitted (Balakirev and Rozhnov 2010; Balakirev et al. 2011, 2012, 2014; Rowe et al. 2008; Pages et al. 2010; Zhang et al. 2016). In total, 79 specimens were investigated (47 *C. langbianis*, 15 *C. thomasi* and 17 *C. chiropus*). The geographic scope of the survey included 23 localities (Fig. 1, Table 1; see also Suppl. material 1: Table S1) scattered over China, Laos, Vietnam and Cambodia and constitute the known species distribution. Most of these sample specimens were collected personally by the authors (AVA, AEB).

### DNA extraction

Small pieces of liver or muscle tissue were sampled in the field and stored in 96% ethanol. Total genomic DNA was extracted using a routine phenol/chloroform/proteinase K protocol (Kocher et al. 1989; Sambrook et al. 1989). The DNA was further purified either by a DNA Purification Kit (Fermentas, Thermo Fisher Scientific Inc., Pittsburgh, PA, USA) or by direct ethanol precipitation. We targeted five genes that were previously used for the phylogenetic analysis of *Chiromyscus* and were available for comparative analyses in GenBank. These genes included a complete Cytochrome *b* (*Cyt b*, 1140 bp); the 5'-proximal portion (680 bp) of subunit I of the Cytochrome Oxidase (*COI*), which is generally used for species diagnoses and for DNA barcoding for a number of mammals (Hebert et al. 2003); a portion of the first exon of the Interphotoreceptor Retinoid Binding Protein gene (*IRBP*, also known as *Rbp3*, up to 1233 bp); the first exon of the Recombination Activation Factor gene (*RAG1*, 1244 bp); and a portion of exon 10 of the Growth Hormone Receptor gene (*GHR*, 815 bp).

### PCR amplification and sequencing

*Cyt b* was amplified using H15915R, CytbRglu (Kocher et al. 1989; Irvin et al. 1991) and CytbRCb9H primers (Robins et al. 2007). The *COI* gene was amplified using the universal conservative primers BatL 5310 and R6036R (Kocher et al. 1989; Irwin et al. 1991). The following universal PCR protocol was used to amplify mtDNA fragments: initial denaturation for 1 min 30 sec at 95 °C, 35 cycles of denaturation for 30 sec at 95 °C, annealing for 1 min at 52 °C and elongation for 30 sec at 72 °C, followed by terminal elongation for 2 min at 72 °C. The PCR was performed in a 30–50 µl volume that contained 2.5–3 µl 10 x standard PCR buffer, 50 mM of each dNTP, 2 mM MgCl<sub>2</sub>, 10 pmol of each primer, 1 unit of Taq DNA polymerase (Fermentas, Thermo



**Figure 1.** Scattering of *Chiromyscus* spp. genetic samples in eastern Indochina. Type localities of *C. chiroplus*, *C. langbianis*, and *C. thomasi* are shown by black circles.

Fisher Scientific Inc., Pittsburgh, PA, USA) and 0.5 µl (25–50 ng) of total DNA template per tube. The reaction was performed using a Tercik (DNK-Tehnologia, Novosibirsk, Russia) thermocycler. *IRBP* (1000–1610 bp in length) was amplified using the *IRBP125f*, *IRBP1435r*, *IRBP1125r* and *IRBP1801r* primers according to Stanhope et al. (1992). A nested PCR technique was applied to amplify *GHR* in accordance with Jansa et al. (2009). An approximately 1.0-kb portion of exon 10 from the *GHR* gene was amplified using the primers *GHRF1* and *GHRendAlt*. This PCR product was re-amplified using the nested *GHRF1* primer paired with *GHR750R* and the

*GHRF50* primer paired with *GHRendAlt*. A 1244 bp portion of the *RAG1* gene was obtained using primers *S70* and *S71*, as described by Stepan et al. (2004).

The PCR products were purified using a DNA Purification Kit (Fermentas, Thermo Fisher Scientific Inc., Pittsburgh, PA, USA). The resulting double-stranded DNA products were directly sequenced in both directions using the Applied Biosystems 3130 Genetic Analyzer with the BigDye Terminator Cycle Sequencing Ready Reaction Kit (Applied Biosystems, Waltham, Massachusetts, USA). All obtained sequences were deposited in GenBank ([www.ncbi.nlm.nih.gov/Genbank](http://www.ncbi.nlm.nih.gov/Genbank); [MK957014–](https://doi.org/10.1093/oxford/abz014)



**Table 1.** List of geographical localities of *Chiromyscus* specimens used for genetic and morphological analyses.

| No. (Fig. 1) | Species              | DNA lineage       | Locality                                   | Elevation (m asl) | Latitude / Longitude |
|--------------|----------------------|-------------------|--|-------------------|----------------------|
| 1            | <i>C. langbianis</i> | N lineage         | China, Yunnan, Xishuangbanna               |                   | 22.0°N, 100.8°E      |
| 5            | <i>C. langbianis</i> | N lineage         | Vietnam, Tuyen Quang, Khong May            | 102               | 22.383°N, 105.339°E  |
| 6            | <i>C. langbianis</i> | N lineage         | Vietnam, Vinh Phuc, Tam Dao                | 850               | 21.452°N, 105.636°E  |
| 7            | <i>C. langbianis</i> | N lineage         | Vietnam, Lang Son, Huu Lien                | 230               | 21.661°N, 106.362°E  |
| 8            | <i>C. langbianis</i> | N lineage         | Vietnam, Nghe An, Pu Hoat                  | 840               | 19.756°N, 104.796°E  |
| 13           | <i>C. langbianis</i> | N lineage         | Laos, Khammouane                           |                   | 17.5°N, 105.33°E     |
| 13           | <i>C. langbianis</i> | N lineage         | Laos, Khammouane, Pha Deng                 |                   | 17.57°N, 105.23°E    |
| 14           | <i>C. langbianis</i> | N lineage         | Vietnam, Quang Binh, Le Thuy, Sa Khia      | 156               | 17.068°N, 106.601°E  |
| 15           | <i>C. langbianis</i> | N lineage         | Vietnam, Kon Tum, Kon Plong                | 1030              | 14.722°N, 108.316°E  |
| 16           | <i>C. langbianis</i> | N lineage         | Vietnam, Kon Tum, Kon Chu Rang             | 1020              | 14.505°N, 108.541°E  |
| 17           | <i>C. langbianis</i> | N lineage         | Vietnam, Gia Lai, Kon Ka Kinh              | 900               | 14.203°N, 108.315°E  |
| 19           | <i>C. langbianis</i> | S lineage         | Vietnam, Lam Dong, Bi Doup-Nui Ba          | 1400–1800         | 12.179°N, 108.679°E  |
| 11           | <i>C. langbianis</i> | Hainan lineage    | China, Hainan, Jianfengling                |                   | 18.74°N, 108.85°E    |
| 12           | <i>C. langbianis</i> | Hainan lineage    | China, Hainan, Baoting                     |                   | 18.641°N, 109.775°E  |
| 18           | <i>C. langbianis</i> | Cambodian lineage | Cambodia, Kaoh Kong, Thmar Bang, Tatai Leu |                   | 11.961°N, 103.303°E  |
| 3            | <i>C. thomasi</i>    | N lineage         | Vietnam, Lao Cai, Bat Xat, Y Ty            | 1830              | 22.624°N, 103.629°E  |
| 4            | <i>C. thomasi</i>    | N lineage         | Vietnam, Son La, Muong Coi                 | 547               | 21.343°N, 104.749°E  |
| 5            | <i>C. thomasi</i>    | N lineage         | Vietnam, Tuyen Quang, Khong May            | 102               | 22.383°N, 105.339°E  |
| 2            | <i>C. thomasi</i>    | S lineage         | Laos, Houay Sai, Houay Khot Station        |                   | 20.267°N, 100.4°E    |
| 9            | <i>C. thomasi</i>    | S lineage         | Vietnam, Nghe An, Xoong Con                | 141               | 19.252°N, 104.318°E  |
| 10           | <i>C. thomasi</i>    | S lineage         | Vietnam, Nghe An, Pu Mat                   | 200               | 18.957°N, 104.686°E  |
| 13           | <i>C. thomasi</i>    | S lineage         | Laos, Khammouane, Pha Deng                 |                   | 17.57°N, 105.23°E    |
| 15           | <i>C. thomasi</i>    | S lineage         | Vietnam, Kon Tum, Kon Plong                | 1030              | 14.722°N, 108.316°E  |
| 20           | <i>C. chiropus</i>   |                   | Vietnam, Lam Dong, Bao Loc                 | 650               | 11.837°N, 107.64°E   |
| 21           | <i>C. chiropus</i>   |                   | Vietnam, Dong Nai, Ma Da                   | 75                | 11.381°N, 107.062°E  |
| 22           | <i>C. chiropus</i>   |                   | Vietnam, Tay Ninh, Lo Go Xa Mat            |                   | 11.583°N, 105.933°E  |
| 23           | <i>C. chiropus</i>   |                   | Vietnam, Ba Ria-Vung Tau, Binh Chau        | 68                | 10.55°N, 107.483°E   |


**MK957137**). *Niviventer* spp., *Rattus norvegicus* and *Mus musculus* were used as outgroups.

## Molecular data analyses

Individual sequences were edited manually using BioEdit v. 7.1.11 (Hall 1999) and aligned by Clustal W software incorporated into BioEdit and MEGA 6. The basic sequence parameter calculations and the best-fitting evolution models and inter- and intrapopulation divergence evaluations were performed using MEGA 6 (Tamura et al. 2013). No pseudogenes were detected for the mitochondrial genes. The optimal substitution models and their parameters are summarised in Suppl. material 1: Table S2. Genetic distances ( $d$ ) between groups under Tamura-Nei gamma distributed invariant sites including (TN93+G+I) or Tamura 3-parameter (T3P) models (Tamura et al. 2004), (depending on the best model determined) were calculated, based on the *Cytb* and *COI* genes in MEGA 6. Bayesian phylogenetic trees were inferred using MrBayes v.3.2. (Huelsenbeck and Ronquist 2001; Ronquist and Huelsenbeck 2003), two MCMCs for four chains with the default heating value and with a burn-in parameter equal to 25% of the initial number of runs. We applied  $6 \times 10^6$  generations for the *Cytb* dataset,  $2 \times 10^6$  for the *COI*,  $4 \times 10^6$  for the *RAG1* datasets and  $5 \times 10^6$  for both *GHR* and *IRBP* datasets until the average standard deviation of split frequencies dropped below the level of 0.0025 after the runs for all datasets investigated. We used a flat Dirichlet prior for the relative nucleotide frequencies and for the relative rate parameters, a discrete uniform prior for the topologies and an exponential distribution for the gamma shape

parameter and all branch lengths. The gamma shape parameters for Bayesian Inference were evaluated directly from a general dataset by MrBayes v.3.2. A burn-in period of one million generations was determined graphically using TRACER v.1.4 (Rambaut and Drummond 2007) to ensure convergence. Consensus trees were built from the last 25% of trees obtained ( $15 \times 10^4$ ,  $15 \times 10^4$ ,  $50 \times 10^3$ ,  $10 \times 10^3$  and  $12.5 \times 10^3$  trees for *Cytb*, *COI*, *RAG1*, *GHR* and *IRBP*, respectively) during the MCMC procedure by MrBayes v.3.2. The five individual genes were concatenated using the software SequenceMatrix v1.7.6 (Vaidya et al. 2011) to create a master alignment of 5,199 bp total (5,208 bp including three triplet insertions in *Mus musculus GHR* gene). A restricted dataset, including all species included in this study and consisting of samples with a complete data matrix, were used for concatenated sequences analyses. A total of  $5 \times 10^6$  generations was applied during the MCMC procedure by MrBayes v.3.2. for concatenated alignment until the average standard deviation value dropped to 0.0077. TREEROT v.3 (Sorenson and Franzosa 2007) was used to examine tree-bisection-reconnection branch-swapping (PBS) to assess the contribution of each data partition in the combined analysis (Baker and DeSalle 1997). This analysis was performed to test the sustainability of the primary internal nodes for the different gene analyses. The robustness of the trees was assessed by posterior probabilities (PP). Trees were visualised and prepared by FigTree v.1.4.3 (Rambaut 2012).

Divergence time approximation was performed by Mega X (Kumar et al. 2018), a time tree inferred using the Reltime method (Tamura et al. 2012; 2018) and the General Time Reversible model and branch lengths evaluated by MrBayes v.3.2. for the concatenated dataset. The



**Figure 2.** The phylogenetic tree (*Cyt b*, Bayesian inference) for *Chiromyscus* genetic lineages radiation. The posterior probability values are presented at the nodes, and the branch lengths (scale bar at the bottom) are indicated above the nodes. The sample labels and locality numeration are indicated as in Fig. 1 and Suppl. material 1: Table S1.

timetree was computed using one calibration constraint chosen as the divergence event between *Mus* and *Rattus* genera which are known to happen within 12.3–11.0 Mya (95% CI) (Benton and Donoghue 2007 with correction of Kimura et al. 2015). Discrete Gamma distribution was used to model evolutionary rate differences amongst sites (five categories (+G, parameter = 0.9185)). The rate variation model allowed for some sites to be evolutionarily invariable ([+I], 54.21% sites). The analysis involved all 82 nucleotide sequences and 5208 nucleotide positions.

### Morphological data analyses

In total, 63 intact skulls of adult *Chiromyscus* (15 *C. chiropus*, 35 *C. langbianis* and 13 *C. thomasi*) obtained from 19 genetically-investigated localities in Vietnam (see Table 1 and Suppl. material 1: Tables S1 for references) were measured for morphometric comparison and analyses. Age was assessed by tooth wear and closure of cranial sutures. Due to the limited sampling, sexual differences were not especially tested; the possible sexual bias was compensated for by equalisation of representatives of different sexes in the sample. The sex ratio did not exceed 15 percentage points.

Twenty measurements were taken from each skull by means of digital calipers to the nearest 0.01 mm: greatest length of skull (ONL), braincase breadth (BBC), braincase height (HBC), zygomatic breadth (ZB), interorbital breadth (IB), length of rostrum (LR), breadth of rostrum

(BR), breadth of zygomatic plate (BZP), diastema length (LD), length of foramina incisive (LIF), breadth of foramina incisive (BIF), length of bony palate (LBP), breadth across the palatal bridge at the level of the first molar (BBP), distance from the anterior edge of the premaxillary to the posterior edge of the palatine (= postpalatal length, PPL), breadth of the mesopterygoid fossa (BMF), length of the bulla (LB), upper molar row length (CLM1-3), first upper molar breadth (BM1), first lower molar breadth (Bm1) and lower molar row length (CLm1-3). The cranial measurements followed Musser et al. (2006) and Balakirev et al. (2011), Suppl. material 1: Fig. S1. The measurements dataset is available from AEB by request.

Principal components analysis (PCA) and canonical discriminant function analysis (DFA) were used to evaluate “distinctiveness” amongst the samples. A one-way analysis of variance (ANOVA) was performed to test the differences amongst groups on all cranial variables. The software programme Statistica 8.0 (StatSoft Inc., Tulsa, OK, USA) was used for all analytical procedures.

## Results

### Phylogenetic subdivision and relationships

The most representative tree was constructed for 66 *Cyt b* sequences. The trees were well supported (PP = 1) (Fig. 2). Different geographical populations of

*C. chiropus* formed a single homogeneous cluster, while *C. thomasi* and *C. langbianis* populations were represented by a few geographically-segregated clusters. *C. thomasi* split into two clusters, which we conventionally called the Northern and Southern phylogroups. The haplotypes forming the Northern cluster were distributed locally over the north-western part of Vietnam, whereas Southern haplotypes spread substantially more widely, stretching to the south to the Tay Nguyen Plateau (also known as the Central Highlands) and westwards to the extreme northwest of Laos. *Chiromyscus langbianis* formed three coherent geographically-distributed clusters. The first from the Dalat Plateau, the *terra typica* of this species; named hereafter as the Southern lineage. Samples originating from continental Indochina, southern Yunnan and north-eastern Vietnam in the north to the Tay Nguyen Plateau in the south formed a second cluster, the Northern lineage. Another cluster, which was sister to the Northern lineage, contained samples from Hainan Island (Figure 2). The genetic divergence within and between these groups is shown in Tables 2, 3. The estimated genetic distances ( $d$ ) for intraspecific *C. langbianis* and *C. thomasi* phylogroups were not very high and fell within the limits of 0.036 and 0.063.

Another tree constructed using the mitochondrial *COI* gene of 43 samples revealed another additional specific phylogenetic lineage (Suppl. material 1: Figure S2), with divergence levels as high as those for the groups described above. (Table 3). It was formed by a single sample from the Cardamom Mountains, Cambodia. Unfortunately, this was the only sample of this group and, so far, there are no data about the distribution of this genetic lineage in southern Indochina. The clustering pattern of *C. thomasi* samples was similar to that obtained using *Cyt b*, with

*C. chiropus* demonstrating two reliable subclusters, combining the animals from the Dalat Plateau foothills (Lam Dong Province) and lowland populations (Tay Ninh, Dong Nai and Ba Ria-Vung Tau Provinces), respectively. Thus, mitochondrial genes supported *C. langbianis* split into four geographically-distributed phylogroups, whereas *C. thomasi* and *C. chiropus* formed two distinctive phylogroups each.


Phylogenetic reconstructions, based on nuclear genes, did not allow us to clarify the relationships and the taxonomic rank of the distinctive phylogroups identified. Thus, only species-level clusters were reliably traced by the *RAG1* gene tree constructed for the 26 available samples (Suppl. material 1: Fig. S3). The overall level of divergence was low and genetic distances did not exceed 0.01. The same species-level groups were identified by the *GHR* gene (Suppl. material 1: Fig. S4), of which we included 52 samples. Within *C. langbianis*, complex soft polytomy without notable geographic segregation was traced, whereas within *C. thomasi*, two clusters corresponding to the mitochondrial phylogenetic lineages mentioned above were clearly demonstrated. At the same time, the considerable length of the branches was apparent for *C. thomasi* and for some specimens of *C. langbianis* from the Dalat Plateau. These branches were significantly longer than those characteristic of *C. chiropus* and most of the *C. langbianis* samples, a trait that may indicate a special pattern of its evolutionary history and, in particular, the longer evolutionary age of these populations. Species-level clusters may also be traced in the *IRBP* gene tree, of which we had 49 samples (Suppl. material 1: Fig. S5). Within the *C. langbianis* cluster, no geographical segregation was traced, which may indicate incomplete sorting of lineages; on the other hand, *C. thomasi*

**Table 2.** Genetic distances ( $d$ , TN93+G+I, gamma = 1.48) for geographic populations and species of *Chiromyscus* as calculated based on the *Cyt b* gene sequence (1140 bp). Standard error (S.E.) estimates are shown above the diagonal.

|                                | <i>C.langbianis</i><br>(Hainan) | <i>C.langbianis</i><br>(Northern) | <i>C.langbianis</i><br>(Southern) | <i>C.chiropus</i> | <i>C.thomasi</i><br>(Northern) | <i>C.thomasi</i><br>(Southern) |                         |        |
|--------------------------------|---------------------------------|-----------------------------------|-----------------------------------|-------------------|--------------------------------|--------------------------------|-------------------------|--------|
| between group distances        |                                 |                                   |                                   |                   |                                |                                | within groups distances |        |
| $d$ (TN93+G+I, Tamura-Nei)     |                                 |                                   |                                   |                   |                                |                                | S.E.                    |        |
| <i>C.langbianis</i> (Hainan)   |                                 | 0.005                             | 0.008                             | 0.012             | 0.015                          | 0.016                          | 0.0061                  | 0.0013 |
| <i>C.langbianis</i> (Northern) | 0.036                           |                                   | 0.007                             | 0.011             | 0.015                          | 0.015                          | 0.0097                  | 0.0015 |
| <i>C.langbianis</i> (Southern) | 0.063                           | 0.052                             |                                   | 0.013             | 0.016                          | 0.016                          | 0.0080                  | 0.0017 |
| <i>C.chiropus</i>              | 0.129                           | 0.123                             | 0.130                             |                   | 0.014                          | 0.013                          | 0.0074                  | 0.0015 |
| <i>C.thomasi</i> (Northern)    | 0.179                           | 0.188                             | 0.187                             | 0.166             |                                | 0.008                          | 0.0039                  | 0.0013 |
| <i>C.thomasi</i> (Southern)    | 0.186                           | 0.191                             | 0.192                             | 0.162             | 0.063                          |                                | 0.0071                  | 0.0016 |

**Table 3.** Genetic distances ( $d$ ; T3P, T92+I) for geographic populations and species of *Chiromyscus* as calculated based on the *COI* gene sequence (680 bp). Standard error (S.E.) estimates are shown above the diagonal.

|                                | <i>C.langbianis</i><br>(Northern) | <i>C.langbianis</i><br>(Southern) | <i>C.langbianis</i><br>(Cambodia) | <i>C.chiropus</i><br>(Lam_Dong) | <i>C.chiropus</i><br>(others) | <i>C.thomasi</i><br>(Northern) | <i>C.thomasi</i><br>(Southern) |        |        |
|--------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|---------------------------------|-------------------------------|--------------------------------|--------------------------------|--------|--------|
| between group distances        |                                   |                                   |                                   |                                 |                               |                                | within groups distances        |        |        |
| $d$ (T3P: GTR)                 |                                   |                                   |                                   |                                 |                               |                                | S.E.                           |        |        |
| <i>C.langbianis</i> (Northern) |                                   | 0.0106                            | 0.0100                            | 0.0176                          | 0.0243                        | 0.0289                         | 0.0305                         | 0.0045 | 0.0020 |
| <i>C.langbianis</i> (Southern) | 0.035                             |                                   | 0.0128                            | 0.0183                          | 0.0243                        | 0.0296                         | 0.0278                         | 0.0022 | 0.0021 |
| <i>C.langbianis</i> (Cambodia) | 0.033                             | 0.044                             |                                   | 0.0190                          | 0.0242                        | 0.0264                         | 0.0292                         |        |        |
| <i>C.chiropus</i> (Lam_Dong)   | 0.083                             | 0.089                             | 0.093                             |                                 | 0.0121                        | 0.0262                         | 0.0277                         | 0.0050 | 0.0024 |
| <i>C.chiropus</i> (others)     | 0.124                             | 0.130                             | 0.127                             | 0.040                           |                               | 0.0294                         | 0.0307                         | 0.0022 | 0.0021 |
| <i>C.thomasi</i> (Northern)    | 0.156                             | 0.161                             | 0.138                             | 0.155                           | 0.172                         |                                | 0.0131                         | 0.0110 | 0.0036 |
| <i>C.thomasi</i> (Southern)    | 0.163                             | 0.153                             | 0.154                             | 0.160                           | 0.178                         | 0.049                          |                                | 0.0037 | 0.0026 |




**Figure 3.** A. The phylogenetic time tree (*Cyt b/COI/RAG1/GHR/IRBP* genes, concatenated analyses Bayesian inference) for *Chiromyscus* genetic lineages radiation. The posterior probability values and average divergence time (Mya, in brackets) are presented at the nodes. Branches lengths are indicated above the branches. B. The position of *Chiromyscus* among most closely relative groups of rodents of SE Asia, marked by arrow (see Pages et al. 2016 for details). **Footnote:** The sample labels and locality numeration are indicated as in Fig. 1 and Suppl. material 1: Table S1.

showed deep trichotomy. The samples corresponding to the Southern cluster of mitochondrial lineages were represented here by two independent branches, which formed populations from the Tay Nguyen Plateau and populations distributed further to the north. Similar to the *GHR* gene tree (Suppl. material 1: Fig. S4), the inequality of the phylogenetic branch lengths should garner attention. However, in contrast to the *GHR* gene, all samples of *C. langbianis* from the Dalat Plateau recovered longer branches. In general, it can be concluded that *C. langbianis* and *C. chiropus* showed significant homogeneity in nuclear genes, whereas *C. thomasi* had the same pattern of nuclear gene variation as traced by mitochondrial markers.

In addition to low support levels for many nuclear gene clades, tree-bisection-reconnection branch-swapping (PBS) analysis indicates an existence of conflicting phylogenetic signals, especially for segments within *C. langbianis*. In general, the low posterior probability values for internal branches and the conflicting phylogenetic signals in many lineages can be explained by a significantly slower evolution rate of nuclear genes (generally weak phylogenetic signal) and incomplete lineage sorting that may be the result of symplesiomorphy. The tree which constructed the concatenated sequence (Fig. 3) is consistent with nuclear gene trees, but posterior probabilities values for some internal nodes are lower, mainly





**Figure 4.** Results of the multivariate analyses of *Chiromyscus* spp. from eastern Indochina. **A.** Ungrouped morphometric separation (PCA analysis); the data were drawn from 20 craniodental measurements. **B.** Grouped morphometric separation (DFA analysis) drawn from the same specimens and measurements.

inside *C. langbianis*. The Hainan cluster is not monophyletic, as one of the samples was recovered in the Northern continental cluster.

## Morphological analyses

The descriptive statistics of the craniodental measurements for phylogenetic lineages of *C. langbianis* (two of four phylogroups discovered were available) and *C. thomasi* that were identified by the abovementioned analyses are summarised in Suppl. material 1: Tables S3, S4. Craniodental measurements for *C. chiropus* are given in Suppl. material 1: Table S5. As revealed by the intergroup F-test, the populations under study demonstrated notable peculiarities of cranial morphology. The samples were significantly different ( $p < 0.05$  or lower) from each other in 18 and 7 cranial characters for *C. langbianis* and *C. thomasi*, respectively.

In a principal components analysis (PCA) drawing on 20 craniodental measurements, the first two axes captured 60.9% (mainly reflecting general size) and 6.4% of the total variation, respectively. ONL, ZB, IB, LD, PPL and CLM1-3 were the six measurements that had the highest correlations with PC 1 (Suppl. material 1: Table S6). In the PCA of skull measurements, all three species of *Chiromyscus* overlapped and *C. langbianis* showed the largest range of variation amongst these species (Fig. 4A). Discriminant function analysis (DFA), which drew on the same variables, provided another means of illuminating the morphometric distinctions and the first two axes captured 53.6% and 25.4% of the variation (Suppl. material 1: Table S3). The DFA yielded moderate to high discrimination amongst all species and genetic lineages (Fig. 4B).

## Discussion

### Taxonomic implications

The concordance of morphological and genetic traits and a good separation of samples in 3D factor space indicate the morphological specificity of the studied populations. On the other hand, the concordant pattern of morphological, genetic and clear geographic subdivision of the mitochondrial phylogroups allow us to question the taxonomic status of these populations; in particular, they allow us to attribute the observed genetic lineages to distinct taxa.

The Northern genetic lineage of *C. langbianis* must be undoubtedly assigned to subspecies *C. l. indosinicus* Osgood, 1932. This taxon was described as *Rattus indosinicus* by Osgood (1932) from Sapa in northern Vietnam, Lao Cai Province. The Northern genetic lineage of *C. thomasi* has to be attributed to the nominotypical subspecies *C. t. thomasi* (this phylogroup includes the holotype of *C. thomasi*). The appropriate name for the Southern genetic lineage of *C. thomasi* is debatable. It may be associated with *Rattus indosinicus vientianensis* Bourret, 1942, described from the surroundings of Vientiane, Laos. However, Musser (1973) treated *vientianensis* as a younger synonym of *langbianis* and, in our previous survey where the most recent genus revision has been made (Balakirev et al. 2014), we also supposed that nomen *vientianensis* should be associated with *C. langbianis*. Unfortunately, we have no specimens from the neighbouring Vientiane and we cannot identify which of the species is distributed there. The holotype of *vientianensis* is unavailable.

The genetic distances between the two phylogroups of *C. thomasi* correspond minimally to the subspecific level (Baker 2006). However, despite their considerable ages,



**Table 4.** Estimated time to most recent common ancestor (Mya) for *Chiromyscus* based on Reltime method and the General Time Reversible model. A discrete Gamma distribution was used to model evolutionary rate differences among sites (5 categories (+G, parameter = 0.9185); The rate variation model allowed for some sites to be evolutionarily invariable ([+I], 54.21% sites). The time tree was computed using 1 calibration constraints.

|              | Calibration<br><i>Mus/Rattus</i><br>divergence<br>point | Clade A<br><i>Chiromyscus/</i><br><i>Niviventer</i> common<br>ancestor | Clade B<br><i>C. thomasi</i><br>divergence<br>point | Clade C<br><i>C. chiropus</i><br>divergence<br>point | Clade D<br>Northern/Southern<br>lineages of <i>C. thomasi</i><br>divergence point | Clade E<br>Cambodian lineage<br>of <i>C. langbianis</i><br>divergence point | Clade F<br>Southern lineage<br>of <i>C. langbianis</i><br>divergence point | Clade G<br>Hainan lineage<br>of <i>C. langbianis</i><br>divergence point |
|--------------|---|--|---|--|---|---|--|--|
| Mean         | 11.65   | 4.85   | 4.09  | 2.70   | 1.19  | 1.248   | 0.950  | 0.621  |
| 95% CI lower | 11.0  | 2.89   | 1.78  | 0.73   | 0.545   | 0.041   | 0.032  | 0.020  |
| 95% CI upper | 12.3  | 7.02   | 5.90  | 3.58   | 3.49  | 2.88  | 2.01   | 1.39   |

these groups have no visually remarkable morphological differences (see Suppl. material 1: Table S3), so we cannot give here formal description of a new subspecies and assert only that the southern populations of *C. thomasi* belong to a unique monophyletic genetic lineage.

### Phylogeography and recent fauna formation

Tree rats are usually confined to forest environments and their dispersal is restricted by the forest edge. They usually do not spread beyond these limits and never cross wide deforested areas as they do not feel confident on open ground surface (Musser 1981; Corbet and Hill 1992; our personal observations). The main kind of natural event that contributed to the dispersal, population segregation and speciation of mammalian fauna is, most probably, repeated disjunction-reconnection events of natural populations that were associated with areas covered by tropical forest during the late Miocene-Holocene (Hall 1998). The distribution pattern and genetic diversification of the genetic lineages revealed in the genus *Chiromyscus* shed light on the natural history and range formation of these species. Based on genetic data, the population of *C. langbianis* inhabiting the Dalat Plateau is apparently older than other recent continental populations. It can be assumed that the Dalat Plateau served as its main refugium during the Holocene climate oscillations (see also Meschersky et al. 2016; Abramov et al. 2017 for another rodent species). Judging by observed genetic distances and the homogeneity of the Northern genetic lineage of *C. langbianis*, its expansion beyond the Plateau occurred fairly quickly. Moreover, there is a reason to suppose multiple refugia, as supported by the fact that the most northern Hainanese cluster occupies a basal position in relation to the continental lineages. Noteworthy, to date, the Hainanese haplogroup may occur amongst continental populations. This may indicate both incomplete lineage sorting in this pair of clusters and an ancient hybridisation event between insular and continental populations. Multiple reconnection events that occurred during the Pleistocene make the second scenario possible. This gives some reason to believe that colonisation of the Island initiated from a different population, not the one that inhabits the Dalat highlands. Instead, it probably originated from an additional northern refugium. The colonisation of

Hainan by *C. langbianis* might have happened simultaneously with those of other Muridae (*Niviventer* and *Rattus*), which are currently represented by distinct insular populations (Pan et al. 2007; Li et al. 2008; Smith and Xie 2008). This event could be dated back to the Late Miocene. According to Voris et al. (2000), Hainan had been connected to the mainland when the sea level was 120–75 m below the current level, which has happened many times, with the longest connections occurring at approximately 0.25, 0.15 and 0.017 Mya. However, judging by the estimated time of species level genetic lineage divergence (over 1 Mya, Fig. 3, Table 4), all of them were formed much earlier than these dates and cannot be associated with recent insularisation. It should be noticed that estimates, evaluated for divergence time for Muridae, are slightly higher than proposed earlier (Rowe et al. 2011; Pages et al. 2016); however, the genus *Chiromyscus* was represented there by only a single individual. Our finding provides evidence in support of more complex patterns of its evolutionary history. In any case, even if our estimates are closer to the higher limit of generic age determined earlier (Fabre et al. 2013; Pages et al. 2016), these timings for group split points are significantly older than the last events of the Hainan-Mainland reconnection. The latter supports the hypothesis of their formation in the continental refugia during the Late Miocene. On the other hand, the occurrence of another original genetic lineage in southern Cambodia, which is an even more ancient separation than the Hainanese, indicates that there may have been several insularisation and resettlement events and that “distribution waves” originated from the Dalat and any other refugia during the Pleistocene.

The split of *C. thomasi* into the Northern and Southern phylogroups happened before the split of the corresponding *C. langbianis* phylogroups and apparently is associated with antecedent global natural factor fluctuations. However, the recent distribution pattern of these species indicates that their natural history differs significantly amongst the populations that originate from different dispersion centres/refugia. As far as can be traced by the data available, *C. thomasi* does not reach the Dalat Plateau and more southern regions inhabited by *C. chiropus* and the Southern lineage of *C. langbianis*. At the same time, *C. thomasi* (both Northern and Southern phylogroups) appears to be distributed sympatrically with the Northern phylogroup of *C. langbianis* in most of eastern

Indochina. This indicates that their possible migration routes alongside the Annamite Range occurred in two opposite directions, with *C. thomasi* moving northwards and *C. langbianis* moving southwards. The fact that *C. thomasi* did not participate in mammal fauna formation on Hainan Island supports the recent natural area expansion of *C. langbianis* and are probably explained by ecological factors. Namely, these phenomena may reflect the ecological preferences of this species. *C. thomasi* is known to be more strictly associated with mountain forest formations than *C. langbianis*, showing greater habitat versatility, which apparently allowed the latter to spread much further eastwards along the plains of eastern Indochina. On the other hand, the significant genetic homogeneity of *C. chiropus*, which inhabits forest formations everywhere in the extreme south of Indochina and its basal position in relation to the genetic lineages of *C. langbianis*, may indicate that these recent populations diverged significantly earlier. This finding also indicates that forest refugia remained at the southern part of Indochina throughout the Holocene and even earlier. They could be associated not only with the Dalat Plateau, but also with the Cardamom Mountains, Bolaven Plateau and probably also with some of the offshore islands on the shelf of the Gulf of Siam.

The distribution pattern of *Chiromyscus* species in the region also raises the problem of the initial intrusion and distribution of *C. chiropus* in eastern Indochina. The *terra typica* for this species is the Karen Mountains in eastern Myanmar, where the species inhabit mountainous forests. As we pointed out earlier (Balakirev et al. 2014), Burmese specimens do not demonstrate noticeable morphological differences from the southern Vietnamese populations and these are treated as conspecific. Unfortunately, there are still no genetic data from Burmese populations that would allow direct comparison of their genetic identity. However, the wide distribution of this species to the east in eastern Indochina through the Yunnan and Annamite Ranges is hampered by the wide distribution of another species, namely, *C. thomasi*, which populates these regions. No cases of sympatry are currently documented, which may suggest a competitive exclusion in this pair of species. At the same time, the existence of a direct connection between the Malacca and southern Indochina in the Holocene by a forest corridor cannot be excluded. Based on data of Meijaard et al. (2003) on tropical forest persistence and the distribution of forest-dependent species on islands of the South China Sea and a forest connection between southern Indochina and Malacca, a southern expansion route is probable. Nevertheless, there are no records on the current distribution of this species in the lowland areas in central Indochina to the west from 105°E.

## Conclusions

We show that the genetic distances between phylogroups of *C. langbianis* and *C. thomasi* correspond to the subspecific level at least. However, these phylogenetic groups do

not demonstrate obvious univocal diagnostic differences in cranial features suitable for species diagnoses without special statistical analysis. Our study shows that the recent phylogenetic structure of *C. langbianis* is the most recent within the genus and appears within several independent refugia that remained isolated throughout the Pleistocene. In turn, the phylogroups of *C. thomasi* are likely older than those of *C. langbianis*. Environmental factors and species preferences followed recent natural ecological shifts which drove allopatry. However, *C. chiropus* demonstrates the greatest age; the ways of formation of the area of this species still remain obscure and are likely to be associated with changes in forest cover in Indochina and Malacca Peninsula during the Pleistocene. The possibility of competitive interaction of these species in the process of formation of their recent natural areas also cannot be excluded.

## Acknowledgements

This study was realised with the support of the Joint Russian-Vietnamese Tropical Research and Technological Center, Hanoi, Vietnam. We thank Dr. Sergei V. Kruskop (Zoological Museum of Moscow State University, Moscow, Russia) and Olga V. Makarova (Zoological Institute of Russian Academy of Sciences, Saint Petersburg, Russia) for giving access to the collections under their care. We are grateful to Dr. Viktor V. Sunstov and Dr. German V. Kuznetsov, whose field collections of skulls and skins we used to investigate morphology. We thank Dr. Nguyen Dang Hoi, Dr. Bui Xuan Phuong, Tran Quang Tien, Le Xuan Son and Tran Huu Coi (all from the Joint Russian-Vietnamese Tropical Research and Technological Center, Hanoi, Vietnam), who put considerable effort into the expedition's preparations. We also thank the administrations of Huu Lien, Ke Go, Kon Chu Rang, Kon Plong, Vinh Cuu Ma Da, Pu Mat, Pu Hoat, Bi Doup-Nui Ba, Lo Go Xa Mat and Binh Chau National Parks and Nature Reserves for their help with managing our research. We are also very grateful to Dr. Miguel Camacho Sanchez (Estación Biológica de Doñana, Sevilla, Spain) and Dr. Melissa T. R. Hawkins (Smithsonian Institution, National Museum of Natural History, Washington, USA) for their helpful and constructive comments on an earlier version of the manuscript. The study was partly supported by the programme of the Ministry of Science and Higher Education of the Russian Federation (project AAAA-A19-119082990107-3).

All authors participated in samples collection, AEB did the genetic analyses and wrote the main part of paper, AEB and AVA together performed the morphological analyses and prepared illustrations; VVR provided funding and coordinated all our surveys in Vietnam.

The study was performed in full agreement with current Vietnamese regulations in the field of Nature Protection and Biodiversity Conservation. We followed the guidelines of the American Society of Mammalogists during the collection and handling of the animals.

## References

- Abramov AV, Balakirev AE, Rozhnov VV (2017) New insights into the taxonomy of the marmoset rats *Hapalomys* (Rodentia: Muridae). *Raffles Bulletin of Zoology* 65: 20–28.
- Aplin KP, Brown PR, Jacob J, Krebs CJ, Singleton GR (2003) Field methods for rodent studies in Asia and the Indo-Pacific. ACIAR Monograph 100, ACIAR, Canberra.
- Aplin K, Lunde D (2016) *Chiromyscus chiropus* (errata version published in 2017). The IUCN Red List of Threatened Species 2016: e.T4669A115069693. <https://doi.org/10.2305/IUCN.UK.2016-3.RLTS.T4669A22453964.en> [Accessed on 20 Apr 2019]
- Baker RH, DeSalle R (1997) Multiple sources of character information and the phylogeny of Hawaiian *Drosophila*. *Systematic Biology* 46: 654–673. <https://doi.org/10.1093/sysbio/46.4.654>
- Balakirev AE, Rozhnov VV (2010) Species composition in the genus *Niviventer* (Rodentia, Muridae) based on studies of the cytochrome *b* gene of mtDNA. *Moscow University Biological Sciences Bulletin* 4: 170–173. <https://doi.org/10.3103/S0096392510040139>
- Balakirev AE, Abramov AV, Rozhnov VV (2011) Taxonomic revision of *Niviventer* (Rodentia, Muridae) from Vietnam: a morphological and molecular approach. *Russian Journal of Theriology* 10(1): 1–26. <https://doi.org/10.15298/rusjtheriol.10.1.01>
- Balakirev AE, Abramov AV, Tikhonov AN, Rozhnov VV (2012) Molecular phylogeny of the *Dacnomys* division (Rodentia, Muridae): the taxonomic positions of *Saxatilomys* and *Leopoldamys*. *Doklady Biological Sciences* 445(1): 251–254. <https://doi.org/10.1134/S0012496612040096>
- Balakirev AE, Abramov AV, Rozhnov VV (2014) Phylogenetic relationships in the *Niviventer-Chiromyscus* complex (Rodentia, Muridae) inferred from molecular data, with description of a new species. *ZooKeys* 451: 109–136. <https://doi.org/10.3897/zookeys.451.7210>
- Benton MJ, Donoghue PCJ (2007) Paleontological evidence to date the tree of life. *Molecular Biology and Evolution* 24: 26–53. <https://doi.org/10.1093/molbev/msl150>
- van den Bergh GD, de Vos J, Sondaar PY (2001) The late quaternary palaeogeography of mammal evolution in the Indonesian Archipelago. *Palaeogeography, Palaeoclimatology, Palaeoecology* 171: 385–408. [https://doi.org/10.1016/S0031-0182\(01\)00255-3](https://doi.org/10.1016/S0031-0182(01)00255-3)
- Bourret R (1942) Sur quelques petits Mammifères du Tonkin et du Laos. *Comptes Rendus des seances du Conseil de recherches scientifiques de l'Indochine*, 2 semestre: 27–30.
- Can DN, Endo H, Son NT, Oshida T, Canh LX, Phuong DH, Lunde DP, Kawada S-I, Hayashida A, Sasaki M (2008) Checklist of wild mammal species of Vietnam. Institute of Ecology and Biological Resources, Hanoi.
- Chen ZP, Jiang XL, Wang YX (1995) Karyotype and banding patterns of fea's tree rat (*Chiromyscus chiropus*). *Caryologia* 48: 9–16. <https://doi.org/10.1080/00087114.1995.10797313>
- Corbet GB, Hill JE (1992) *Mammals of the Indo-Malayan Region: a Systematic Review*. Oxford University Press, Oxford.
- Cox CB, Moore PD (2000) *Biogeography: an ecological and evolutionary approach* (6th ed.). Blackwell Science Ltd., Oxford.
- Cranbrook E (2000) Northern Borneo environments of the past 40,000 years: archaeozoological evidence. *Sarawak Museum Journal* 55: 62–109.
- Dang HH, Dao DVT, Cao VS, Pham TA, Hoang MK (1994) Checklist of mammals in Vietnam. Publishing House "Science and Technics", Hanoi.
- Gannon WL, Sikes RS (2011) American Society of Mammalogists. Guidelines of the American Society of Mammalogists for the use of wild animals in research. *Journal of Mammalogy* 92(1): 235–253. <https://doi.org/10.1644/10-MAMM-F-355.1>
- Gathorne-Hardy FJ, Syaekani, Davies RG, Eggleton P, Jones DT (2002) Quaternary rainforest refugia in south-east Asia: using termites (Isoptera) as indicators. *Biological Journal of the Linnean Society* 75: 453–466. <https://doi.org/10.1046/j.1095-8312.2002.00031.x>
- Ge D, Lu L, Xia L, Du Y, Wen Z, Cheng J, Abramov AV, Yang Q (2018) Molecular phylogeny, morphological diversity, and systematic revision of a species complex of common wild rat species in China (Rodentia, Murinae). *Journal of Mammalogy* 99(6): 1350–1374. <https://doi.org/10.1093/jmammal/gyy117>
- Fabre P-H, Pagès M, Musser GG, Fitriana YS, Fjeldså J, Jennings A, Jonsson KA, Kennedy J, Michaux J, Semiadi G, Supriatna N, Helgen KM (2013) A new genus of rodent from Wallacea (Rodentia: Muridae: Murinae: Rattini), and its implication for biogeography and Indo-Pacific Rattini systematics. *Zoological Journal of the Linnean Society* 169(2): 408–447. <https://doi.org/10.1111/zoj.12061>
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98. <https://doi.org/10.1021/bk-1999-0734.ch008>
- Hebert PDN, Cywinska A, Ball SL, DeWaard JR (2003) Biological identifications through DNA barcodes. *Proceedings of the Royal Society B* 270: 313–321. <https://doi.org/10.1098/rspb.2002.2218>
- Hall R (1998) The plate tectonics of Cenozoic Southeast Asia and the distribution of land and sea. In: Hall R, Holloway JD (Eds) *Biogeography and geological evolution of SE Asia*. Backhuys, Leiden.
- Huelsenbeck JP, Ronquist F (2001) MRBAYES: Bayesian inference of phylogeny. *Bioinformatics* 17: 754–755. <https://doi.org/10.1093/bioinformatics/17.8.754>
- Irwin D, Kocher TD, Wilson AS (1991) Evolution of the cytochrome *b* gene of mammals. *Journal of Molecular Evolution* 32: 128–144. <https://doi.org/10.1007/BF02515385>
- Jansa SA, Giarla TC, Lim BK (2009) The phylogenetic position of the rodent genus *Typhlomys* and the geographic origin of Muroidea. *Journal of Mammalogy* 90: 1083–1094. <https://doi.org/10.1644/08-MAMM-A-318.1>
- Kimura Y, Hawkins MTR, McDonough MM, Jacobs LL, Flynn LJ (2015) Corrected placement of *Mus-Rattus* fossil calibration forces precision in the molecular tree of rodents. *Scientific Reports* 5: e14444. <https://doi.org/10.1038/srep14444>
- Kocher TD, Thomas WK, Meyer A, Edwards SV, Paabo S, Villablanca F, Wilson A (1989) Dynamics of mitochondrial DNA evolution in animals: amplification and sequencing with conserved primers. *PNAS* 86: 6196–6200. <https://doi.org/10.1073/pnas.86.16.6196>
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: Molecular Evolutionary Genetics Analysis across computing platforms. *Molecular Biology and Evolution* 35: 1547–1549. <https://doi.org/10.1093/molbev/msy096>
- Lan H, Chen Z, Wang Y, Shi L (1994) The mitochondrial DNA restriction map of the Fea's tree mouse (*Chiromyscus chiropus*) and its evolutionary relationship with mouse and rat. *Zoological Research* 15: 39–44.
- Li Y, Wu Y, Harada M, Lin LK, Motokawa M (2008) Karyotypes of three rat species (Mammalia: Rodentia: Muridae) from Hainan Island, China, and the valid specific status of *Niviventer lotipes*. *Zoological Science* 25: 686–692. <https://doi.org/10.2108/zsj.25.686>



- Marshall JT (1977) Family Muridae: rats and mice. In: Lekagul B, McNeely JA (Eds) Mammals of Thailand. Association for the Conservation of Wildlife, Bangkok.
- Lu L, Chesters D, Zhang W, Li G, Ma Y, Ma H, Song X, Wu H, Meng F, Zhu C, Liu Q (2012) Mammal Investigation in Spotted Fever Focus with DNA-Barcoding and Taxonomic Implications on Rodents Species from Hainan of China. PLoS ONE 7(8): e43479. <https://doi.org/10.1371/journal.pone.0043479>
- Lu L, Ge DY, Chesters D, Ho SYW, Ma Y, Li GC, Wen ZX, Wu YJ, Wang J, Xia L, Liu JL, Guo TY, Zhang XL, Zhu CD, Yang QS, Liu QY (2015) Molecular phylogeny and the underestimated species diversity of the endemic white-bellied rat (Rodentia: Muridae: *Niviventer*) in southeast Asia and China. Zoologica Scripta 44: 475–494. <https://doi.org/10.1111/zsc.12117>
- Lunde D, Nguyen TS (2001) An identification guide to the rodents of vietnam. New York: American Museum of Natural History Center for Biodiversity and Conservation.
- Lunde DP, Musser GG, Nguyen TS (2003) A survey of small mammals from Mt. Tay Con Linh II, Vietnam, with the description of a new species of *Chodsigoa* (Insectivora: Soricidae). Mammal Study 28: 31–46. <https://doi.org/10.3106/mammalstudy.28.31>
- Meijaard E (2003) Mammals of south-east Asian islands and their Late Pleistocene environments. Journal of Biogeography 30: 1245–1257. <https://doi.org/10.1046/j.1365-2699.2003.00890.x>
- Meijaard E, Groves CP (2006) The geography of mammals and rivers in mainland Southeast Asia. In: Lehman SM, Fleagle JG (Eds) Primate Biogeography. Springer, New York.
- Meschersky IG, Abramov AV, Lebedev VS, Chichkina AN, Rozhnov VV (2016) Evidence of a complex phylogeographic structure in the Indomalayan pencil-tailed tree mouse *Chiropodomys gliroides* (Rodentia: Muridae) in eastern Indochina. Biochemical Systematics and Ecology 65: 147–157. <https://doi.org/10.1016/j.bse.2016.02.015>
- Musser GG (1973) Species limits of *Rattus cremoriventer* and *Rattus langbianis*, murid rodents of southeastern Asia and the Greater Sunda Islands. American Museum of Natural History Nov 2525: 1–65.
- Musser GG (1981) Results of the Archbold expeditions. No.105. Notes on systematics of Indo-Malayan murid rodents, and descriptions of new genera and species from Ceylon, Sulawesi and the Philippines. Bulletin of the American Museum of Natural History 168: 225–334.
- Musser GG, Carleton MD (1993) Family Muridae. In: Wilson DE, Reeder DM (Eds) Mammal species of the world. A taxonomic and geographic reference (2<sup>nd</sup> edn.). Smithsonian Institution Press, Washington, 501–756.
- Musser GG, Carleton MD (2005) Family Muridae. In: Wilson DE, Reeder DM (Eds) Mammal species of the world. A taxonomic and geographic reference (3<sup>rd</sup> edn.). John Hopkins University Press, Baltimore, 894–1531.
- Musser GG, Lunde DP, Nguyen TS (2006) Description of a new genus and species of rodent (Murinae, Muridae, Rodentia) from the Tower Karst region of northeastern Vietnam. American Museum of Natural History Nov 3517: 1–41. [https://doi.org/10.1206/0003-0082\(2006\)3517\[1:DOANGA\]2.0.CO;2](https://doi.org/10.1206/0003-0082(2006)3517[1:DOANGA]2.0.CO;2)
- Osgood WH (1932) Mammals of the Kelley-Roosevelts and Delacour Asiatic expeditions. Field Museum of Natural History. Zoological series 18: 193–339. <https://doi.org/10.5962/bhl.title.2798>
- Pages M, Chaval Y, Herbretau V, Waengsothorn S, Cosson JF, Hugot JP, Morand S, Michaux J (2010) Revisiting the taxonomy of the Rattini tribe: a phylogeny-based delimitation of species boundaries. BMC Evolutionary Biology 10: e184. <https://doi.org/10.1186/1471-2148-10-184>
- Pages M, Fabre P-H, Chaval Y, Mortelliti A, Nicolas V, Wells K, Michaux JR, Lazzari V (2016) Molecular phylogeny of South-East Asian arboreal murine rodents. Zoologica Scripta 45(4): 349–364. <https://doi.org/10.1111/zsc.12161>
- Pan QH, Wang YX, Yan K (2007) A field guide to the mammals of China. China Forestry Publishing House, Beijing.
- Rambaut A, Drummond AJ (2007) Tracer, version 1.4. <http://tree.bio.ed.ac.uk/software/tracer> [accessed 15 October 2019]
- Rambaut A (2012) FigTree v1.4.3: Tree Figure Drawing Tool. <http://treebioedacuk/software/figtree/> [Accessed 2019 May 15]
- Robins JH, Hingston M, Matisoo-Smith E, Ross HA (2007) Identifying *Rattus* species using mitochondrial DNA. Molecular Ecology Notes 7: 717–729. <https://doi.org/10.1111/j.1471-8286.2007.01752.x>
- Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. Bioinformatics 19: 1572–1574. <https://doi.org/10.1093/bioinformatics/btg180>
- Rowe KC, Reno ML, Richmond DM, Adkins RM, Steppan SJ (2008) Pliocene colonization and adaptive radiations in Australia and New Guinea (Sahul): multilocus systematics of the old endemic rodents (Muroidea: Murinae). Molecular Phylogenetics and Evolution 47(1): 84–101. <https://doi.org/10.1016/j.ympev.2008.01.001>
- Rowe KC, Aplin KP, Baverstock PR, Moritz C (2011) Recent and rapid speciation with limited morphological disparity in the genus *Rattus*. Systematic Biology 60: 188–203. <https://doi.org/10.1093/sysbio/syq092>
- Rowe KC, Achmadi AS, Fabre P-H, Schenk JJ, Steppan SJ, Esselstyn JA (2019) Oceanic islands of Wallacea as a source for dispersal and diversification of murine rodents. Journal of Biogeography 46: 2752–2768. <https://doi.org/10.1111/jbi.13720>
- Sambrook J, Fritsch EF, Maniatis T (1989) Molecular cloning: a laboratory manual. Cold Spring Harbor Lab, Cold Spring Harbor.
- Sorenson MD, Franzosa EA (2007) *TreeRot, Version 3*. Boston University, Boston. <http://people.bu.edu/msoren/TreeRot.html> [accessed 15 October 2019]
- Smith AT, Xie Y (2008) A guide to the mammals of China. Princeton University Press, Princeton and Oxford.
- Stanhope MJ, Czelusniak J, Si J-S, Nickerson J, Goodman M (1992) A molecular perspective on mammalian evolution from the gene encoding interphotoreceptor retinoid binding protein, with convincing evidence for bat monophyly. Molecular Phylogenetics and Evolution 1: 148–160. [https://doi.org/10.1016/1055-7903\(92\)90026-D](https://doi.org/10.1016/1055-7903(92)90026-D)
- Steppan SJ, Storz BJ, Hoffmann RS (2004) Nuclear DNA phylogeny of the squirrels (Mammalia, Rodentia) and the evolution of arboreality from c-myc and RAG1. Molecular Phylogenetics and Evolution 30: 703–719. [https://doi.org/10.1016/S1055-7903\(03\)00204-5](https://doi.org/10.1016/S1055-7903(03)00204-5)
- Tamura K, Nei M, Kumar S (2004) Prospects for inferring very large phylogenies by using the neighbor-joining method. PNAS 101: 11030–11035. <https://doi.org/10.1073/pnas.0404206101>
- Tamura K, Stecher G, Peterson D, Filipiński A, Kumar S (2013) MEGA6: Molecular evolutionary genetics analysis version 6.0. Molecular Phylogenetics and Evolution 30: 2725–2729. <https://doi.org/10.1093/molbev/mst197>
- Tamura K, Battistuzzi FU, Billing-Ross P, Murillo O, Filipiński A, Kumar S (2012) Estimating Divergence Times in Large Molecular Phylogenies. PNAS 109: 19333–19338. <https://doi.org/10.1073/pnas.1213199109>

- Tamura K, Qiqing T, Kumar S (2018) Theoretical foundation of the RelTime Method for estimating divergence times from variable evolutionary rates. *Molecular Phylogenetics and Evolution* 35: 1770–1782. <https://doi.org/10.1093/molbev/msy044>
- Thomas O (1891) Diagnoses of three new mammals collected by Signor L. Fea in the Carin Hills, Burma. *Annali del Museo civico di storia naturale di Genova (Ser. 2)* 10: e884.
- Thomas O (1925) The mammals obtained by Mr. Herbert Stevens on the Sladen-Godman expedition to Tonkin. *Proceedings of the Zoological Society of London* 95: 495–506. <https://doi.org/10.1111/j.1096-3642.1925.tb01524.x>
- Vaidya G, Lohman, DJ, Meier R (2011) SequenceMatrix: concatenation software for the fast assembly of multigene datasets with character set and codon information. *Cladistics* 27(2): 171–180. <https://doi.org/10.1111/j.1096-0031.2010.00329.x>
- Voris HK (2000) Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time durations. *Journal of Biogeography* 27: 1153–1167. <https://doi.org/10.1046/j.1365-2699.2000.00489.x>
- Wang YX (2002) A complete checklist of mammal species and subspecies in China: a taxonomic and geographic reference. China Forestry Publishing House, Beijing.
- Wurster CM, Bird MI, Bull ID, Creed F, Bryant C, Dungait JA, Paz V (2010) Forest contraction in north equatorial southeast Asia during the last glacial period. *PNAS* 107: 15508–15511. <https://doi.org/10.1073/pnas.1005507107>
- Zhang B, He K, Wan T, Chen P, Sun G, Liu S, Nguyen TS, Lin L, Jiang X (2016) Multi-locus phylogeny using topotype specimens sheds light on the systematics of *Niviventer* (Rodentia, Muridae) in China. *BMC Evolutionary Biology* 16(1): e261. <https://doi.org/10.1186/s12862-016-0832-8>

## Supplementary material 1

### Tables S1–S6, Figures S1–S5

Authors: Alexander E. Balakirev

Data type: phylogenetic, morphological

Explanation note: Tables, samples and other reference materials.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zse.97.57490.suppl1>

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Zoosystematics and Evolution](#)

Jahr/Year: 2021

Band/Volume: [97](#)

Autor(en)/Author(s): Balakirev Alexander E., Abramov Alexei V., Rozhnov Viatcheslav V.

Artikel/Article: [Distribution pattern and phylogeography of tree rats \*Chiromyscus\* \(Rodentia, Muridae\) in eastern Indochina 83-95](#)