

**RECENT PLECOPTERA LITERATURE**  
**(CALENDAR YEAR 1998 AND EARLIER)**

**Note:** Unless otherwise indicated, references are to the original title in the language in which the article is written. If titles are given in two languages separated by a slash, the translated title was provided by the author(s) and there is also an abstract in that same language. If the second title is in English but appears in square brackets, an editorial translation of the title was included for the convenience of readers of PERLA, but there is no English abstract.

- ALBARIÑO, R.J.; BALSEIRO, E.G. (1998):** Larval size and leaf conditioning in the breakdown of *Notofagus pumilio* leaves by *Klapopteryx kuscheli* (Insecta, Plecoptera) in a South Andean stream. — Intern. Rev. Hydrobiol. 83: 397-404.
- ALFORD M.H. (1998):** New records of *Allocapnia* (Plecoptera: Capniidae) from Mississippi and Louisiana, with accompanying scanning electron micrographs. — Entomol. News 109 (3): 183-188.
- ANSORGE, J. (1996):** Insekten aus dem oberen Lias von Grimmen (Vorpommern, Norddeutschland) [Insects from the upper Liassic of Grimmen (Vorpommern, North Germany)]. — Neue Palaeontol. Abhandl. 2: 1-132, pls 1-17; FOSSILS; Plecoptera on pp. 33-36.
- ASHE, P.; O'CONNOR, J.P.; MURRAY, D.A. (1998):** A Checklist of Irish Aquatic Insects. — Occ. Publ. Ir. Biogeog. Soc. 3, 80 pp.
- BACK, C.; LEBLANC, A.; AUBIN, A. (1982):** Effets sur la dérive des insectes aquatiques d'un traitement au Temephos contre les larves de moustiques dans le Québec subarctique. — Can. Entomol. 115: 703-712.
- BAKER S.C.; SHARP H.F. JR. (1998):** Evaluation of the recovery of a polluted urban stream using the Ephemeroptera-Plecoptera-Trichoptera index. — J. Freshwat. Ecol. 13 (2): 229-234.
- BANNING, M., LEUCHS, H., RÜTTEN, M., SCHLEUTER, M., SCHÖLL, F., & TITTIZER, T. (1989):** Die Bundeswasserstraßen als Lebensraum für Tiere [Federal waterways as animal habitats.]. — Jber. Bundesanst. Gewkde., Koblenz, BfG-0592: 1-25.

- BARLET, J. (1986):** Contribution à la connaissance du squelette thoracique des Plécoptères (Insectes) [Contribution to the knowledge of the thoracic skeleton of the Plecoptera (Insects)]. — Bull. Soc. r. Sci. Liège, 55(5-6): 607-616.
- BARLET, J. (1987):** Contribution à la connaissance de la musculature thoracique des Plécoptères [Contribution to the knowledge of the thoracic musculature of the Plecoptera (Insects)]. — Bull. Anns Soc. r. belge Entomol. 123: 165-178.
- BELLSTEDT, R. (1980):** *Hydraena subimpressa* Rey (Col., Hydraenidae) neu für die Fauna der DDR [*Hydraena subimpressa* Rey (Col., Hydraenidae) new for the fauna of the DDR]. — Abh. Ber. Mus. Nat. Gotha, 1980: 61-62.
- BOURNAUD, M.; TACHET, H.; BERLY, A.; CELLOT, B. (1998):** Importance of microhabitat characteristics in the macrobenthos distribution of a large river reach. — Anns Limnol. 34(1): 83-98.
- BRINKMANN, R.; REUSCH, H. (1998):** Zur Verbreitung der aus dem norddeutschen Tiefland bekannten Ephemeroptera- und Plecoptera-Arten (Insecta) in verschiedenen Biotoptypen / Notes on the distribution of all known species of Ephemeroptera and Plecoptera (Insecta) in the lowland of North Germany from different biotopes. — Braunschw. naturkd. Schr. 5(3): 531-540.
- BRODERSEN, K.P.; DALL, P.C.; LINDEGAARD, C. (1998):** The fauna in the upper stony littoral of Danish lakes: macroinvertebrates as trophic indicators. — Freshwat. Biol. 39: 577-592.
- BUCKTON S.T.; BREWIN P.A.; LEWIS A.; STEVENS P.; ORMEROD S.J. (1998):** The distribution of dippers, *Cinclus cinclus* (L.), in the acid-sensitive region of Wales, 1984-95. — Freshwat. Biol. 39 (2): 387-396.
- CEREGHINO, R.; LAVANDIER, P. (1998):** Influence of hydropeaking on the distribution and larval development of the Plecoptera from a mountain stream. — Regulated Rivers: Res. Management 14(3): 297-309.
- CHESSMAN, B.C.; MCKOY, P.K. (1998):** Towards diagnostic biotic indices for river macroinvertebrates. — Hydrobiologia 364(2): 169-182 (1997/98).
- COLLIER, K.J.; WILCOCK, R.J.; MEREDITH, A.S. (1998):** Influence of substrate type and physico-chemical conditions on macroinvertebrate faunas and biotic indices in some lowland Waikato, New Zealand streams. — N. Z. J. Marine Freshwat. Res. 32(1): 1-19.

- DAHMEN, D.; DAHMEN, E.-G.; GELLERT, G. (1998):** Einfluß extremer Schwermetallbelastungen auf die Zusammensetzung der Makrobenthoszönose eines Mittelgebirgsbaches in einem ehemaligen Erzabbaugebiet des Raumes Overath (Bergisches Land) [Influence of extreme heavy metal loads on the composition of the macrobenthic coenosis in a foothill stream in a former ore mining area in the district of Schaffrath (Bergisches Land)]. — *Decheniana* 151: 173-182.
- DELK, J.K.; KILGORE, M.J.; STARK, B.P. (1998):** Comparison of epiproct structure of two closely related species, *Sweltsa fidelis* (Banks) and *S. revelstoka* (Jewett) (Plecoptera: Chloroperlidae). — *Gt Basin Natur.* 58(3): 282-284.
- DEWALT, R.E.; STARK, B.P.; HARRIS, M.A. (1998):** *Perlesta golconda* (Plecoptera: Perlidae), a new stonefly species from Illinois.—*Ent. News* 109(5):315-317.
- DINIZ FILHO J.A.F.; OLIVEIRA L.G.; SILVA M.M. (1998):** Explaining the beta diversity of aquatic insects in "Cerrado" streams from central Brazil using multiple Mantel test. — *Rev. Brasil. Biol.* 58 (2): 223-231.
- DU, Y. (1995):** A taxonomic study on the family Perlidae from China (Plecoptera: Perloidea). — PhD-Thesis, Northwestern Agricultural University, Yangling, Shaanxi, China, X + 254 pp. In Chinese, English abstract.
- DU, Y.; CHOU, L. (1998):** Taxonomic study of the genus *Agnentina* Klapálek (Plecoptera: Perlidae: Perlinae) from China. — *Entomotaxonomia* 20(2): 100-110. In Chinese, English summary.
- DUFFIELD, R.M.; NELSON, C.H. (1998):** Stoneflies (Plecoptera) in the diet of brook trout (*Salvelinus fontinalis* Mitchell) in Libby Creek, Wyoming, USA—*Hydrobiologia* 380:59-65.
- DUNKEL, F.V.; RICHARDS, D.C. (1998):** Effect of Azadirachtin formulation on six nontarget aquatic macroinvertebrates. — *Environm. Entomol.* 27 (3): 667-67
- EGGLETON, M.A.; MORGAN, E.L.; PENNINGTON, W.L. (1996):** Effects of liming on an acid-sensitive southern Appalachian stream. — *Restor. Ecol.* 4: 247-263.
- ERKINARO H.; ERKINARO J. (1998):** Feeding of Atlantic salmon, *Salmo salar* L., parr in the subarctic River Teno and three tributaries in northernmost Finland. — *Ecol. Freshwat. Fish* 7 (1): 13-24.

- ERMAN, N.A. (1998): Invertebrate richness and trichoptera phenology in Sierra Nevada (California, USA) cold springs: sources of Variation—In: BOTOSANEANU, L. (ed.). Studies in crenobiology: The biology of springs and springbrooks; Backhuys Publishers, Leiden, pp95-108**
- FABRE, E.; CHAUVET, E. (1998): Leaf breakdown along an altitudinal stream gradient. — Arch. Hydrobiol. 141: 167-179.**
- FINCK, P. (1998): Der Einfluß von Probenahmezeitpunkt und —häufigkeit auf die Erfassung von Makroinvertebraten in Mittelgebirgsbächen / The influence of sample timing and frequency on the recording of the macroinvertebrates in mountain brooks. — Lauterbornia 34: 245-254.**
- FISCHER, J.; FISCHER, F.; SCHNABEL, S.; WAGNER, R.; BOHLE, H.W. (1998): Die Quellenfauna der hessischen Mittelgebirgsregion. Besiedlungsstruktur, Anpassungsmechanismen und Habitatbindung der Makroinvertebraten am Beispiel von Quellen aus dem Rheinischen Schiefergebirge und der osthessischen Buntsandsteinlandschaft / Spring fauna of the Hessian Mittelgebirge — Population structure, adaptative strategies, and relations to habitats of the macroinvertebrates, as exemplified by springs in the the Rhenish metamorphic shield and in the East-Hessian sandstone plate. — In: BOTOSANEANU, L. (ed.): Studies in Crenobiology. The biology of springs and springbrooks; Backhuys Publishers, Leiden, 1998, pp. 183-199 (ISBN90-73348-04-8).**
- FLOOK P.K.; ROWELL C.H.F. (1998): Inferences about orthopteroid phylogeny and molecular evolution from small subunit nuclear ribosomal DNA sequences. — Insect Molecul. Biol. 7 (2): 163-178.**
- FOCHETTI, R.; DE BIASE, A.; BELFIORE, P.; AUDISIO, P. (1998): Faunistica e biogeografia regionale dei Plecotteri Italiani (Plecoptera) [Faunistics and regional biogeography of the Italian stoneflies (Plecoptera)]. — Mem. Soc. Ent. in Italian. 76: 3-19.**
- FRIBERG, N.; REBSDORF, A.; LARSEN, S.E. (1998): Effects of afforestation on acidity and invertebrates in Danish streams and implications for freshwater communities in Denmark. — Water, Air & Soil Pollution 101(1-4): 235-256.**
- GALDEAN N. (1997): Some considerations on the benthic communities of the Upper Tisa River. — Trav. Mus. Nat. Hist. Natur. "Grigore Antipa" 37: 255-286.**
- GARVEY, J.E.; MARSCHALL, E.A.; WRIGHT, R.A. (1998): From star charts to stoneflies: detecting relationships in continuous bivariate data. — Ecology 79: 442-447. -70-**

- GELWICK, F.P.; STOCK, M.S.; MATTHEWS, W.J. (1997):** Effects of fish, water depth, and predation risk on patch dynamics in a north-temperate river ecosystem. — *Oikos* 80: 382-398
- GEREBEN-KRENN, B.-A.; PASS, G. (1998):** Morphology and evolution of the circulatory organs of the caudal appendages in basic insects. — *Book of Abstracts, Proc. Vith Europ. Congr. Entomol.* (BRUNNHOFER, V.; SOLDÁN, T., eds), Ěeské Budejovice, 1: 62.
- GRIFFITH, M.B.; BARROWS, E.M.; PERRY, S.A. (1998):** Lateral dispersal of adult aquatic insects (Plecoptera, Trichoptera) following emergence from headwater streams in forested Appalachian catchments. — *Ann. Entomol. Soc. Amer.* 91(2): 195-201.
- GRUBAUGH, J.W.; WALLACE, J.B.; HOUSTON, E.S. (1997):** Production of benthic macroinvertebrate communities along a southern Appalachian river continuum. — *Freshwat. Biol.* 37(3): 581-596.
- HAAPALA, A.; MUOTKA, T. (1998):** Seasonal dynamics of detritus and associated macroinvertebrates in a channelized boreal stream. — *Arch. Hydrobiol.* 142: 171-189.
- HALL, R.O.; MEYER, J.L. (1998):** The trophic significance of bacteria in a detritus-based stream food web. — *Ecology* 79: 1995-2012.
- HARMAN W.N. (1997):** Changes in the Otsego Lake macrobenthos communities between 1935 and 1994. — *Lake & Reservoir Management* 13 (2): 160-169.
- HAWKINS, C.P.; HOGUE, J.N.; DEKKER, L.M.; FEMINELLA, J.W. (1997):** Channel morphology, water temperature, and assemblage structure of stream insects. — *J. N. Am. Benthol. Soc.* 16: 728-749.
- HELESIC, j.; KUBICEK, F.; ZAHRAĐKOVA, S. (1998):** The impact of regulated flow and altered temperature regime on river bed macroinvertebrates.—In: BRETSCHKO, G; HELESIC, J. (eds). *Advances in river bottom ecology; River Bottom IV Symposium, Brno, August 1996; Backhuys Publishers, Leiden, pp225-243.*
- HICKEY C.W.; CLEMENTS W.H. (1998):** Effects of heavy metals on benthic macroinvertebrate communities in New Zealand streams. — *Envir. Toxicol. Chemist.* 17 (11): 2338-2346.

- HOKE R.A.; ANKLEY G.T.; KOSIAN P.A.; COTTER A.M.; VANDERMEIDEN F.M.; BALCER M.; PHIPPS G.L.; WEST C.; COX J.S. (1997):** Equilibrium partitioning as the basis for an integrated laboratory and field assessment of the impacts of DDT, DDE and DDD in sediments. — *Ecotoxicol.* 6 (2): 101-125.
- HÜBNER, G.; KLOSE, S. (1998):** Erfolgskontrolle der Renaturierung eines Baches in der Lüneburger Heide anhand der Ephemeroptera, Plecoptera und Trichoptera. / Monitoring of the development of a restored stream in the Lüneburger Heide/North German Lowland by focussing on Ephemeroptera, Plecoptera and Trichoptera. — *Lauterbornia* 34: 53-66.
- HUNTSMAN, B.O.; BAUMANN, R.W.; KONDRATIEFF, B.C. (1999):** Stoneflies (Plecoptera) of the Black Hills of South Dakota and Wyoming, USA: Distribution and zoogeographic affinities.—*Gr. Basin Natur.* 59: 1-17.
- ILLYOVA M.; TOMAJKA J. (1998):** Interstitial community composition in several montane and submontane streams in the West Carpathians. — *Biologia (Bratislava)* 53 (2): 159-166.
- INADA, K. (1998):** Illustrated stonefly adults (Insecta, Plecoptera) of Hyôgo Prefecture, Honshu, Japan, Part 2, Perlidae (1). — *Biol. Inland Waters, Nara*, 13: 24-66. In Japanese.
- ISOBE, Y. (1997):** Egg morphology in Systellognatha (Insecta: Plecoptera)—68th Annual Meeting Zool. Soc. of Japan Nara, December 1997; *Zool. Sci. Tokyo* 14(Suppl), 40.
- ISOBE, Y. (1997):** The origin of the daily rhythmicity in insect behaviors and of aquatic insects. — *Biol. Sci. News* 312:12-16. In Japanese.
- JACOBSEN, D. (1998):** The effect of organic pollution on the macroinvertebrate fauna of Ecuadorian highland streams. — *Arch. Hydrobiol.* 143: 179-195.
- JACOBSEN, D.; ENCALADA, A. (1998):** The macroinvertebrate fauna of Ecuadorian highland streams in the wet and dry season. — *Arch. Hydrobiol.* 142: 53-70.
- JOOST, W.; KÜTTNER, R. (1998):** Beitrag zur Kenntnis der Steinfliegen Sachsens (Plecoptera) [Contribution to the knowledge of stoneflies of Saxony (Plecoptera)]. — *Entomol. Nachr. Ber.* 41 (4) (1997): 213-232.

- KLONOWSKA OLEJNIK M.; FIALKOWSKI W. (1998):** Ephemeroptera and Plecoptera as indicators of environmental changes caused by reservoir construction. — **BRETSCHKO, G.; HELESIC, J. (eds).** *Advances in river bottom ecology; River Bottom IV Symposium, Brno, August 1996; Backhuys Publishers, Leiden, pp. 215-223.*
- KNISPEL, S.; LUBINI, V.; VINÇON, G.; REDING, J.-P. (1998):** Four Plecoptera species new for the Swiss fauna. — *Mitt. schweiz. entomol. Ges. 71(1-2): 165-168.*
- KÖLLNER, T.; HAASE, P. (1998):** Über einige seltene Wasserinsekten des Westharzes (Insecta: Ephemeroptera, Plecoptera, Coleoptera, Trichoptera) / Contribution to some rare captured water-insects of the western Harz Mountains (Insecta: Ephemeroptera, Plecoptera, Coleoptera, Trichoptera). — *Göttinger Naturkundl. Schr. 4: 173-179 [1997].*
- KONDRATIEFF B.C.; KIRCHNER R.F. (1998):** A new species of *Sweltsa* (Plecoptera: Chloroperlidae) from eastern North America. — *Entomol. News 109 (4): 293-295.*
- KRNO I. (1998):** Influence of abiotic and biotic factors on the life cycles and production of stoneflies (Plecoptera) in an acidified spring area. — *Biologia (Bratislava) 53 (2): 195-204.*
- KÜTTNER, R. (1998):** *Protonemura austriaca* (Theischinger, 1976) in Sachsen (Plecoptera) [*Protonemura austriaca* (Theischinger, 1976) in Saxony (Plecoptera)]. — *Entomol. Nachr. Ber. 42(3): 173-174.*
- LANDA, V.; SOLDÁN, T.; ZAHRÁDKOVÁ, S.; HELESIC, J. (1998):** Composition of fauna of stoneflies (Plecoptera) of the Czech Republic from the biodiversity conservation point of view. — **Book of Abstracts, Proc. VIth Europ. Congr. Entomol. (BRUNNHOFER, V.; SOLDÁN, T., eds), ěeské Budejovice, 1: 412-413.**
- LEMLY, A.D. (1998):** Bacterial growth on stream insects: potential for use in bioassessment. — *J. N. Am. Benthol. Soc. 17(2): 228-238.*
- LINDEGAARD, C.; BRODERSEN, K.P.; WIBERG-LARSEN, P.; SKRIVER, J. (1998):** Multivariate analyses of macrofaunal communities in Danish springs and springbrooks. — In: **BOTOSANEANU, L. (ed.):** *Studies in Crenobiology. The biology of springs and springbrooks; Backhuys Publishers, Leiden, pp. 201-219.*

- LUZON-ORTEGA, J.M.; TIERNO DE FIGUEROA, J.M.; SANCHEZ-ORTEGA, A. (1998): Faunistic composition and phenology of the stoneflies (Insecta: Plecoptera) from the Sierra de Baza (Granada, Spain). — *Boln Asoc. Esp. Entomol.* 23(3-4): 125-138.
- LYON, M.L.; STARK, B.P. (1997): *Alloperla* (Plecoptera: Chloroperlidae) of Western North America. — *Entomol. News* 108(5): 321-334.
- MATTHAEI, C.D.; WERTHMÜLLER, D.; FRUTIGER, A. (1998): An update on the quantification of stream drift. — *Arch. Hydrobiol.* 143: 1-19.
- McLELLAN, I. D. (1998): A revision of *Acroperla* (Plecoptera: Zelandoperlinae) and removal of species to *Taraperla* new genus. — *N. Z. J. Zool.* 25: 185-203.
- McPEEK, M.A.; PECKARSKY, B.L. (1998): Life histories and the strengths of species interactions: combining mortality, growth, and fecundity effects. — *Ecology* 79: 867-879.
- MEEGAN, S.K.; PERRY, S.A.; PERRY, W.B. (1996): Detrital processing in streams exposed to acidic precipitation in the Central Appalachian Mountains. — *Hydrobiologia* 339: 101-110.
- MILLETT, M.C.; LEARNER, M.A. (1998): Plecoptera and Ephemeroptera species assemblages in the Usk river system, Wales. — *Arch. Hydrobiol.* 143: 307-334.
- MULLEN, G.R.; VERCAMMEN-GRANDJEAN, P.H. (1980): The Larval stage of Stygothrombiinea Thor, 1935 and Wandesiinae Schworbel, 1961 and erection of a new superfamily, Stygothrombioidea.—*Intern of Acarol.*6(1): 25-28.
- MÖBES-HANSEN, B.; WARINGER, J.A. (1998): The influence of hydraulic stress on microdistribution patterns of zoobenthos in a sandstone brook (Weidlingbach, Lower Austria). — *Intern. Rev. Hydrobiol.*, 83: 381-396.
- MOREIRA, G.R.P. (1998): Reproductive morphology and copulatory mechanism of the stonefly *Agnentina capitata* (Pictet)(Plecoptera, Perlidae). *Rev. Brasil. Entomol.* 41: 267-283.
- MURPHY, J.F.; GILLER, P.S.; HORAN, M.A. (1998): Spatial scale and the aggregation of stream macroinvertebrates associated with leaf packs. — *Freshwat. Biol.* 39: 325-337. s
- OLIVEIRA L.G.; BISPO P.D.C.; SA N C.D. (1997): Ecology of benthic insects communities (Ephemeroptera, Plecoptera and Trichoptera) in streams at Parque Ecologico de Goiania, Goias, Brazil. — *Rev. Brasil. Zool.* 14 (4): 867-876. In Portuguese.

- ORENDT, C. (1998): Macroinvertebrates and diatoms as indicators of acidification in forest spring brooks in a region of eastern Germany (Leipzig-Halle-Bitterfeld) highly impacted by industrial activities. — Arch. Hydrobiol. 143: 435-467.**
- PARK, S.J.; PARK, S.Y.; KIM, J.S.; BAE, Y.J. (1997): Altitudinal distribution of Ephemeroptera, Plecoptera, and Trichoptera in a mountain stream of Chiaksan. Entomol. Res. Bull. 23: 37-42.**
- PECKARSKY, B.L.; MCINTOSH, A.R. (1998): Fitness and community consequences of avoiding multiple predators. — Oecologia 113(4): 565—576.**
- PENNUTO C.M.; DENOYELLES F.JR.; CONRAD M.A.; VERTUCCI F.A.; DEWEY S.L. (1998): Winter macroinvertebrate communities in two Montane Wyoming streams. — Gt Basin Natur. 58 (3): 231-244.**
- PERRIN, C.P.; RICHARDSON, J.S. (1997): N and P limitation of benthos abundance in the Nechako River, British Columbia. — Can. J. Fish. Aquat. Sci. 54: 2574-2583.**
- PLAGUE G.R.; WALLACE J.B.; GRUBAUGH J.W. (1998): Linkages between trophic variability and distribution of *Pteronarcys* spp. (Plecoptera: Pteronarcyidae) along a stream continuum. — Amer. Midl. Natur. 139 (2): 224-234.**
- PRINGLE, C.M.; RAMIREZ, A. (1998): Use of both benthic and drift sampling techniques to assess tropical stream invertebrate communities along an altitudinal gradient, Costa Rica. — Freshwat. Biol. 39(2): 359-373.**
- RAVIZZA, C. (1998): Plecoptera Capniidae of the Italian Region. Insecta. — Atti Soc. it. Sci. nat. Mus. civ. Stor. nat. Milano 139(1): 73-95.**
- REUSCH, H.; BRINKMANN, R.; PINZ, K.; SPETH, S. (1996): Bemerkenswerte Eintagsfliegen (Ephemeroptera) aus dem norddeutschen Tiefland [Remarkable mayflies (Ephemeroptera) from the lowland of northern Germany]. — Lauterbornia 27: 21-33**
- REUSCH, H.; WEINZIERL, A. (1998): Rote Liste der Steinfliegen (Plecoptera) / Red list of stoneflies (Plecoptera). — In BINOT, M.; BLESS, R.; BOYE, P.; GRUTTKE, H.; PRETSCHER, P. (eds): Rote Liste gefährdeter Tiere Deutschlands, Schriftenreihe für Landschaftspflege und Naturschutz, 55; Bundesamt f. Naturschutz, Bonn-Bad Godesberg, pp. 255-259.**

- REYNOLDS E.; DE VILLIERS C.; DAVIES B.R. (1997): A comparison of the food sources of stoneflies (Plecoptera) from an open- and a closed-canopy headwater stream in South Africa using stable-isotope techniques. — *S. Afr. J. Aquat. Sci.* 23 (1): 3-13.
- ROBINSON, C.T.; GESSNER, M.O.; WARD, J.V. (1998): Leaf breakdown and associated macroinvertebrates in alpine glacial streams. — *Freshwat. Biol.* 40: 215-228.
- ROSCISZEWSKA E. (1997): Morphology of the accessory ovaries in adult male of the stone fly, *Perla marginata* (Panzer) (Plecoptera: Perlidae). — 22nd Conference on Embryology: Plants, Animals, Humans, Osieczany, Poland, May 14-17, 1997. *Acta Biol. Cracoviensia, Ser. Zool.* 39 (SUPPL. 1): 43.
- ROSCISZEWSKA E. (1997): Previtellogenesis and vitellogenesis in the panoistic ovary of *Perla* sp. (Plecoptera: Perlidae) stone-fly. — 22nd Conference on Embryology: Plants, Animals, Humans, Osieczany, Poland, May 14-17, 1997. *Acta Biol. Cracoviensia, Ser. Zool.* 39 (SUPPL. 1): 28.
- ROSCISZEWSKA, E. (1998): The structure of the egg capsules of the stoneflies (Plecoptera) — a comparative study. — *Book of Abstracts, Proc. Vth Europ. Congr. Entomol.* (BRUNNHOFER, V.; SOLDÁN, T., eds), Ěeské Budejovice, 1: 38. 76
- ROSCISZEWSKA, E.; SOLDÁN, T. (1998): Morphology of accessory ovaries in adult males of *Perla marginata* (Plecoptera: Perlidae). — *Book of Abstracts, Proc. Vth Europ. Congr. Entomol.* (BRUNNHOFER, V.; SOLDÁN, T., eds), Ěeské Budejovice, 1: 183-184.
- RUFFIEUX, L.; ÉLOUARD, J.-M.; SARTORI, M. (1998): Flightlessness in mayflies and its relevance to hypotheses on the origin of insect flight. — *Proc. R. Soc. Lond., B*, 265: 2135-2140.
- SANDBERG, J.B.; SZCZYTKO, S.W. (1997): Life cycle of *Isoperla lata* (Plecoptera: Perlodidae) in a central Wisconsin trout stream. — *Gt. Lakes Entomol.* 30 (4): 143-160.
- SANTAMARIA S. (1997): *Lancisporomyces*, a new genus of Trichomycetes with lance-shaped zygospores. — *Mycologia* 89 (4): 639-642,
- SCHÖLL, F.; BALZER, I. (1998): Das Makrozoobenthos der deutschen Elbe 1992-1997 / The macroinvertebrates in the German River Elbe. — *Lauterbornia* 32: 113-129.

- SCHÖLL, F.; BANNING, M. (1998):** Faunistische Untersuchungen (Aquatische Makrofauna) an der Donau (Kelheim — Jochenstein) zur Erfassung des ökologischen Ist—Zustandes [Faunistic studies on the Danube (Kelheim — Jochenstein) for documentation of the present ecological status]. — Bundesanstalt f. Gewässerkunde, BfG\_1128: 37 pp. and 9 unnumbered pp. in attachments.
- SHIMIZU, T. (1998):** The genus *Protonemura* in Japan (Insecta: Plecoptera: Nemouridae). — Species Diversity 3 (1): 133-154.
- SHIMIZU, T. (1998):** The Group of *Amphinemura flavostigma* (Plecoptera, Nemouridae). — Aquatic Insects 20(4): 203-208
- SIEBERT, M. (1998):** Wasserinsekten im Hyporhithral und Epipotamal der Fulda, einst und jetzt / Aquatic insects in the hyporhithral and epipotamal zones of the river Fulda, present and past. — Lauterbornia 33: 53-83.
- SINICHENKOVA N.D. (1998):** New Upper Mesozoic stoneflies from Central Transbaikalia (Insecta, Perlida equals Plecoptera). — Paleontol. Zh. 1998 (2): 64-69. In Russian,
- SINITSHENKOVA, N.D. (1998):** The first European Cretaceous stonefly (Insecta, Perlida = Plecoptera). — Cretaceous Research 19: 317-321. FOSSIL.
- SOLDÁN T. (1997):** The Ephemeroptera: Whose sister-group are they? — In: LANDOLT, P.; SARTORI, M. (eds), Ephemeroptera & Plecoptera Biology-Ecology-Systematics, Fribourg: Mauron + Tinguely & Lachat SA, 514-519.
- SOLDÁN, T.; PAPÁČEK, M.; NOVÁK, K.; ZELENÝ, J. (1996):** The Šumava Mountains: a unique biocentre of aquatic insects (Ephemeroptera, Odonata, Plecoptera, Megaloptera, Trichoptera and Heteroptera-Nepomorpha). — Silva Gabreta 1: 179-186.
- SOLDÁN, T.; ZHRÁDKOVÁ, S.; HELESIC, J.; LANDA, V. (1998):** Distributional and Quantitative Patterns of Ephemeroptera and Plecoptera in the Czech Republic: A Possibility of Detection of Long-term Environmental Changes of Aquatic Biota. — Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia 98, 305 pp.
- SOLUK, D.A.; RICHARDSON, J.S. (1997):** The role of stoneflies in enhancing growth of trout: a test of the importance of predator-predator facilitation within a stream community. — Oikos 80: 214-219.

- SOMERS, K.M.; REID, R.A.; DAVID, S.M. (1998): Rapid biological assessments: How many animals are enough: J. N. Am. Benthol. Soc. 17: 348-358.
- SPETH S. (1997): Ephemeroptera, Plecoptera and Trichoptera (Insecta) of the Dallbek (Lauenburger Geest, Schleswig-Holstein). — Faunist.-Ökol. Mitt. 7 (5-6): 213-219. In German.
- STARK, B.P. (1998): The *Anacroneuria* of Costa Rica and Panama (Insecta: Plecoptera: Perlidae). — Proc. Biol. Soc. Wash. 111(3): 551-603.
- STARK, B.P.; SIVEC, I. (1998): *Anacroneuria* of Peru and Bolivia (Plecoptera: Perlidae).—Scoplia 40: 1-64. (J. Of Scolenian Museum of Natur. Hist. Ljubljana. In Engl., with Engl. and Slovene abstracts).
- STARK, B.P.; STEWART, K.W.; SZCZYTKO, S.W.; BAUMANN, R.W. (1998): Common names of stoneflies (Plecoptera) from the United States and Canada. Ohio Biol. Surv. Notes 1: 1-18.
- STARK, B.P.; SZCZYTKO, S.W.; NELSON, C.R. (1998): American stoneflies: A photographic guide to the Plecoptera.—The Caddis Press. Columbus, Ohio. 128pp.
- STOLZENWALD, T.; SCHMIDT-BRÜCKEN, R. (1998): Das Makrozoobenthos der Schwabach und Trubach (Regnitz/Main). The Makrozoobenthos of the streams Schwabach and Trubach (Regnitz/Main) in North Bavaria. — Lauterbornia 32: 131-149.
- STONE, M.K.; WALLACE, J.B. (1998): Long-term recovery of a mountain stream from clear-cut logging: the effect of forest succession on benthic invertebrate community structure. — Freshwat. Biol. 39: 151-169.
- TAPPENBECK, L.; BÖHME, D. (1997): 4.3.2.5 Steinfliegen (Plecoptera). — pp. 176-181 in: Arten- und Biotopschutzprogramm Sachsen-Anhalt Landschaftsraum Harz. — Ber. Landesamt Umweltschutz Sachsen-Anhalt, Sonderheft 4.
- TAYLOR B.W.; ANDERSON C.R.; PECKARSKY B.L. (1998): Effects of size at metamorphosis on stonefly fecundity, longevity, and reproductive success. — Oecologia 114 (4): 494-502.
- TIERNO DE FIGUEROA, J.M.; LUZÓN-ORTEGA, J.M.; SÁNCHEZ-ORTEGA, A. (1998): Imaginal biology of the stonefly *Hemimelaena flaviventris* (Pictet, 1841) (Plecoptera: Perlodidae). — Ann. Zool. Fennici 35: 225-230.

- TIERNO DE FIGUEROA, J.M.; MORO PÉREZ, M.; SÁNCHEZ-ORTEGA, A. (1998):** Primeras citas de Plecópteros (Insecta: Plecoptera) en Ceuta /España, Norte de Africa. / First records of stoneflies (Insecta: Plecoptera) in Ceuta (Spain, North of Africa). — *Boln Asoc. Esp. Ent.* 22(3-4): 252-253. In Spanish.
- TIUNOVA, T.M.; TESLENKO, V.A.; MEDVEDEVA, L.A.; KOCHARINA, S.L. (1997):** The ecosystem of a small salmon river in the Far East of Russia. — pp. 99-105 In: WADA, E.; TIMOSHKIN, O.A. ; FUJITA, N.; TANIDA, K. (eds): *New scope on boreal ecosystems in East Siberia. DIWPA series, Vol. 2, 180 pp.,* Novosibirsk, Siberian Branch Russ. Acad. Sci.
- TONKOV, V. (1925):** Über den Bau der Rectaldrüsen bei Insekten [On the build of rectal glands of insects]. — *Z. Morph. Ök. Tiere* 4: 416-429.
- TOWNSEND, C.R.; SCARSBROOK, M.R. (1997):** The intermediate disturbance hypothesis, refugia, and biodiversity in streams. — *Limnol. Oceanogr.* 42: 938-949.
- TRAYLER, K.M.; DAVIS, J.A. (1998):** Forestry impacts and the vertical distribution of stream invertebrates in south-western Australia. — *Freshwat. Biol.* 40: 331-342. 79
- VINÇON, G.; PARDO, L. (1998):** Three New *Leuctra* Species from Tunisia (Plecoptera: Leuctridae). — *Aquatic Insects* 20(2): 109-123.
- WACHS, B. (1998):** Ökobewertung der Schwermetallbelastung von Fließgewässern mittels Pflanzen- und Zoobenthon-Arten [Eco-assessment of heavy metal contamination of running waters via species of plant and zoobenthos]. — *Münchner Beitr. Abwasser-, Fischerei- und Flußbiologie* 51 (Integrierte Ökologische Gewässerbewertung — Inhalte und Möglichkeiten): 534-585.
- WAGNER, F.. (1989):** Aufenthalt und Flugverhalten von Imagines merolimnischer Insekten mit dem Schwerpunkt Plecoptera in der Bach-Aue am Beispiel der Leutra (Thuringen). —Diplomarbeit, Friedrich-Schiller-Universität Jena (Dr. W. Schonborn, Dr. W. Joost), 58 pp
- WEINZIERL, A. (1988):** Plecoptera, Steinfliegen. – In: *Dienstbesprechung Biologisch-ökologische Gewässeruntersuchung 1998, Kurzfassungen der Beiträge; Bayerisches Landesamt für Wasserwirtschaft, Materialien* 77: 69-77.

- WHITING M.F.; CARPENTER J.C.; WHEELER Q.D.; WHEELER W.C.** (1997): The Strepsiptera problem: Phylogeny of the holometabolous insect orders inferred from 18S and 28S ribosomal DNA sequences and morphology. — *Syst. Biol.* 46 (1): 1-68.
- WIDDIG, T., SCHMIDT, T.** (1998): Rote Liste der Steinfliegen (Plecoptera) Hessens. — 23 pp. in: *Natur in Hessen*, ed. NaturHessisches Mineisterium des Innern und fur Landwirtschaft, Forsten und Naturschutz, Wiesbaden.
- WILLIAMS, D.D.; WILLIAMS, N.E.** (1998): Aquatic insects in an estuarine environment: densities, distribution, salinity tolerance. — *Freshwat. Biol.* 39: 411-421.
- WILLIAMS, D.D.; WILLIAMS, N.E.** (1998): Seasonal variation, export dynamics and consumption of freshwater invertebrates in an estuarine environment. — *Estuarine, Coastal and Shelf Science* 46(3): 393-410.
- WOLF K W.** (1997): The formation of accessory tubules in spermatids of the red firebug, *Pyrrhocoris apterus* (Hemiptera: Pyrrhocoridae). — *Eur. J. Entomol.* 94 (2): 263-270.  
#Ken, refers to *Isoperla*; delete this line once read!#
- WRIGHT, J.F.; FURSE, M.T.; ARMITAGE, P.D.** (1994): Use of macroinvertebrate communities to detect environmental stress in running waters. In: *SUTCLIFFE, D.W. (ed.): Water quality and stress indicators in marine and feshwater systems: linking levels of organisation.* — Windermere: *Freshwat. Biol. Assoc.*, 15-34.
- YOKUM, K.A.; ANGRADI, T.R.; TARTER, D.C.** (1996): Ecology of *Peltoperla arcuata* and *Tallaperla maria* (Plecoptera: Peltoperlidae) at the Fernow Experimental Forest, Tucker County, West Virginia. — *Psyche* 102(3-4): 151-168.
- YOSHIMURA, M.; ISOBE, Y.; OISHI, T.** (1997): The factors which lead to success of mating in stoneflies (Plecoptera)—68th Annual Meeting Zool. Soc. Of Japan, Nara, October 1997; *Zool. Sci. Tokyo* 14 (Suppl), 60.
- ZHILTOVA, L.A.; POTIKHA, E.V.** (1992): Additions to the fauna of stoneflies (Plecoptera) of Sikhote-Alinsky Biosphere Reserve. — *Entomol. Obozr.* 71(3): 569-573. In Russian.