

RECENT PLECOPTERA LITERATURE (CALENDAR YEAR 2003 AND EARLIER)

- ALBARÍÑO, R.; VILLANUEVA, V.D. (2003): Feeding plasticity and growth of two Plecopteran larvae in a low order Andean stream. – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 31-38.
- ANDRIKOVICS, S.; MURANYI, D. (2001): A checklist of stoneflies with remarks of published, undocumented species and two species new to the Hungarian fauna (Insecta: Plecoptera). – *Folia Ent. Hungarica* 62: 23-35. (correction of title in PERLA 21).
- ARNEKLEIV, J.V.; KJAERSTAD, G. (2003): Changes in density and intra-watercourse distribution of Plecoptera following hydropower regulation in a Central Norwegian river. – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 255-262.
- ARSCOTT, D.B.; TOCKNER, K.; WARD, J.V. (2003): Spatio-temporal patterns of benthic invertebrates along the continuum of a braided Alpine river. – *Arch. Hydrobiol.* 158: 431-460.
- BAILEY, W.J. (2003): Insect duets: Underlying mechanisms and their evolution. – *Physiological Entomology* 28(3): 157-174.
- BAUMGÄRTNER, D.; ROTHHAUPT, K.-O. (2003): Predictive length-dry mass regressions for freshwater invertebrates in a pre-alpine lake littoral. – *Internat. Rev. Hydrobiol.* 88: 453-463.
- BEAMISH, R.J.; NOAKES, D.J.; NOAKES, D.L.G.; BEAMISH, F.W.H. (2003): William Edwin Ricker, OC, FRSC, LLD, DSc. – *Can. J. Fisheries Aquat. Sci.* 60(1): iii-v.
- BEJAOUI, M.; BOUMAIZA, M. (2001): Morphological and ecological data on nymphs of six species of stoneflies of Tunisia (Insecta, Plecoptera). – *Bull. Soc. Zool. Fr.* 126(1-2): 224-225.
- BEJAOUI, M.; BOUMAIZA, M. (2002): Life cycle of *Eoperla ochracea* Kolbe, 1888 (Plecoptera, Perlidae) in Tunisia. – *Bull. Soc. Zool. Fr.* 127(2): 95-101.
- BELLSTEDT, R.; SAMIETZ, R. (2002): Katalog der in den Sammlungen des Museums der Natur in Gotha aufbewahrten Typen. Teil 1: Insekten. [Catalog of types in the collections of the Museum der Natur in Gotha. Part 1: Insects.] – *Abh. Ber. Museum d. Natur Gotha* 22: 187-195.
- BERLIN, A. (2003): Checkliste der Eintags- und Steinfliegen (Insecta: Ephemeroptera, Plecoptera) in Mecklenburg-Vorpommern [Checklist of Ephemeroptera and Plecoptera in Mecklenburg-Vorpommern, Germany]. – *Lauterbornia* 47: 5-11.
- BEUTEL, R.G.; GORB, S.N. (2001): Ultrastructure of attachment specializations of hexapods (Arthropoda): evolutionary patterns inferred from a revised ordinal phylogeny. – *J. Zool. Syst. Evol. Res.* 39(4): 177-207.

- BISPO, P.C.; FROEHLICH, C.G.; OLIVEIRA, L.G. (2002): Stonefly (Plecoptera) fauna in a mountainous area of Central Brazil: Abiotic factors and nymph density. – Revista Brasileira Zool. 19 (suppl.) 325-334.
- BOJSEN, B.H.; JACOBSEN, D. (2003): Effects of deforestation on macroinvertebrate diversity and assemblage structure in Ecuadorian Amazon streams. – Arch. Hydrobiol. 158: 317-342.
- BOULTON, A.J. (2003): Parallels and contrasts in the effects of drought on stream macroinvertebrate assemblages. – Freshwat. Biol. 48(7): 1173-1185.
- BOYERO, L. (2003): Multiscale patterns of spatial variation in stream macroinvertebrate communities. – Ecological Research 18(4): 365-379.
- BOYERO, L. (2003): The effect of substrate texture on colonization by stream macroinvertebrates. – Ann. Limnol. Int. J. Lim. 39: 211-218.
- BRAASCH, D. (2003): *Isoperla pawlowskii* Wojtas 1961 (Insecta, Plecoptera) wieder in Deutschland [*Isoperla pawlowskii* Wojtas 1961 (Plecoptera, Insecta) back in Germany]. – Lauterbornia 46: 121-124.
- BRAASCH, D. (2003): Rückkehr der Steinfliegen (Plecoptera) in Oder und Lausitzer Neiße [Return of stoneflies (Plecoptera) in Oder and Lausitzer Neiße]. – Lauterbornia 46: 93-101.
- BRETTFELD, R.; BELLSTEDT, R. (2003): Die Verbreitung der flussbewohnenden Steinfliege Brachyptera braueri (Klapálek, 1900) in Werra und Saale, zugleich ein Beitrag zur Situation der Flussinsekten in Thüringen. [Distribution of the river-inhabiting stonefly Brachyptera braueri (Klapálek, 1900) in Werra und Saale, and a contribution to the status of river insects in Thuringia]. – Veröffentlichungen Naturhist. Museum Schleusingen 18: 69-78.
- BRIERS, R.A.; CARISS, H.M.; GEE, J.H.R. (2003): Flight activity of adult stoneflies in relation to weather. — Ecological Entomology 28(1): 31-40.
- BRITTAIN, J.E.; CASTELLA, E.; KNISPEL, S.; LENCIOMI, V.; LODS-CROZET, B.; MAIOLINI, B.; MILNER, A.M.; SALTVEIT, S.J.; SNOOK, D.L. (2003): Ephemeropteran and Plecopteran communities in glacial rivers. – In: GAINO, E.(ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia 271-277.
- BURCHER, C.L.; SMOCK, L.A. (2002): Habitat distribution, dietary composition and life history characteristics of odonate nymphs in a blackwater costal plain stream. – Amer. Midland Natur. 148(1): 75-89.

- CALL, R.G.; BAUMANN, R.W. (2002): Stoneflies (Plecoptera) of southern Utah with an updated checklist of Utah species. — *Monographs of the Western North American Naturalist* 1: 65-89.
- CAO, T.K.T.; BAE, Y.J. (2002): Nymphs of two peltoperlid stoneflies (Plecoptera: Peltoperlidae) from Vietnam. — *Insecta Koreana* 19(3-4): 299-302.
- CAUCHIE, H.M. (2002): Chitin production by arthropods in the hydrosphere. — *Hydrobiologia* 470: 63-96.
- CEREGHINO, R.; GIRAUDEL, J.L.; COMPIN, A. (2001): Spatial analysis of stream invertebrates distribution in the Adour-Garonne drainage basin (France), using Kohonen self organizing maps. — *Ecol. Model.* 146(1-3): 167-180.
- CEREGHINO, R.; PARK, Y.-S.; COMPIN, A.; LEK, S. (2003): Predicting the species richness of aquatic insects in streams using a limited number of environmental variables. — *J. N. Amer. Benthol. Soc.* 22(3): 442-456.
- CHERTOPROOD, M.V. (2002): Stream macrobenthic fauna of Klin-Dmitroiv elevation. — *Biology of Inland Waters* (Russ. Akad. Sci., Mockwa): 16-24.
- COLLIER, K.J.; BURY, S.; GIBBS, M. (2002): A stable isotope study of linkages between stream and terrestrial food webs through spider predation — *Freshwat. Biol.* 47(9): 1651-1659.
- COLLIER, K.J.; QUINN, J.M. (2003): Land-use influences macroinvertebrate community response following a pulse disturbance. — *Freshwat. Biol.* 48(8): 1462-1481.
- COLLIER, K.J.; SMITH, B.J. (2003): Role of wood in pumice-bed streams II: Breakdown and colonisation. — *Forest Ecology & Management* 177(1-3): 261-276.
- COLLIER, K.J.; SMITH, B.J.; QUINN, J.M.; SCARSBROOK, M.R.; HALLIDAY, N.J.; CROKER, G.F.; PARKYN, S.M. (2000): Biodiversity of stream invertebrate faunas in a Waikato hill-country catchment in relation to land use. — *N. Z. Entomol.* 23: 9-22.
- COMPIN, A.; CEREGHINO, R. (2003): Sensitivity of aquatic insect species richness to disturbance in the Adour-Garonne stream System (France). — *Ecological Indicators* 3(2): 135-142.
- CORTES, R.M.V.; FERREIRA, M.T.; OLIVEIRA, S.V.; OLIVEIRA, D. (2002): Macroinvertebrate community structure in a regulated river segment with different flow conditions. — *River Res. Appl.* 18(4): 367-382.
- COURTNEY, G.W. (1993): Archaic black flies and ancient forests: conservation of *Parasimilium*-habitats in the Pacific Northwest. — *Aquatic Conservation: marine and Freshwater Ecosystems*, Vol.3: 361-373. John Wiley & Sons.

- DANGLES, O.; JONSSON, M.; MALMQVIST, B. (2002): The importance of detritivore species diversity for maintaining stream ecosystem functioning following the invasion of a riparian plant. – *Biological Invasions* 4(4): 441-446.
- DAVIS, S.; GOLLADAY, S.W.; VELLIDIS, G.; PRINGLE, C.M. (2003): Macroinvertebrate biomonitoring in intermittent coastal plain streams impacted by animal agriculture. – *J. Environmental Quality* 32(3): 1036-1043.
- DEATH, R.G. (2003): Spatial patterns in lotic invertebrate community composition: Is substrate disturbance actually important? – *Can. J. Fisheries Aquatic Sci.* 60(5): 603-611.
- DI GIOVANNI, M.V.; GORETTI, E.; CECCAGNOLI, D.; LA PORTA, G.; CHIAPPAFREDDO, U. (2003): Ephemeroptera and Plecoptera in the Chiascio river (Central Italy) since a dam's building. – In: GAINO, E. (ed.), *Research update on Ephemeroptera and Plecoptera*; University of Perugia; pp. 293-298.
- DIAMOND, J.M.; BRESSLER, D.W.; SERVEISS, V.B. (2002): Assessing relationships between human land uses and the decline of native mussels, fish, and macroinvertebrates in the Clinch and Powell River watershed, USA. – *Environ. Tox. Chem.* 21: 1147-1155.
- DOBRIN, M.; GIBERSON, D.J. (2003): Life history and production of mayflies, stoneflies, and caddisflies (Ephemeroptera, Plecoptera, and Trichoptera) in a spring-fed stream in Prince Edward Island, Canada: evidence for population asynchrony in spring habitats? – *Can. J. Zool.* 81: 1083-1095.
- DOHET, A. (2002): Are caddisflies an ideal group for the biological assessment of water quality in streams? – *Nova Supplementa Entomologica* 15: 507-520.
- DORN, A.; WEINZIERL, A. (2003): Ein Beitrag zur Kenntnis der Eintags- und Steinfliegen-Fauna (Ephemeroptera; Plecoptera) der niederbayerischen Isar [A contribution to the knowledge of the mayflies and stoneflies (Ephemeroptera; Plecoptera) from the river Isar in Bavaria inferior/Germany]. – *Lauterbornia* 47: 21-26.
- DUGGAN, I.C.; SCARSBROOK, M.R.; QUINN, J.M. (2003): Comparison of coded abundance and fixed count rapid assessment techniques for biomonitoring in New Zealand streams. – *N. Z. J. Marine Freshwater Research* 37(1): 23-29.
- EARLE, J.I. (2003): Stonefly (Plecoptera) emergence patterns and flight periods in Indian Creek Watershed, Pennsylvania, USA. – In: GAINO, E. (ed.), *Research update on Ephemeroptera and Plecoptera*; University of Perugia; pp. 167-175.
- EGGERT, S.L.; WALLACE, J.B. (2003): Litter breakdown and invertebrate detritivores in a resource-depleted Appalachian stream. – *Arch. Hydrobiol.* 156: 315-338.

- ELLIOTT, J.M. (2002): A continuous study of the total drift of freshwater shrimps, *Gammarus pulex*, in a small stony stream in the English Lake District. – Freshwat. Biol. 47(1): 75-86.
- ELLIOTT, J.M. (2003): A comparative study of the functional response of four species of carnivorous stoneflies. – Freshwat. Biol. 48: 191-202.
- ELLIOTT, J.M. (2003): A comparative study of the dispersal of 10 species of stream invertebrates. – Freshwat. Biol. 48(9): 1652-1668.
- ELLIOTT, J.M. (2003): Interspecific interference and the functional response of four species of carnivorous stoneflies. – Freshwat. Biol. 48(9): 1527-1539.
- ENTING, K. (2002): Beitrag zur Kenntnis der Steinfliegenfauna des Echtersbachtales bei Brecht (Südeifel) (Insecta: Plecoptera). [Contribution to the knowledge of the Plecoptera fauna in the valley of the stream Echtersbach near Brecht (Southern Eifel) (Insecta: Plecoptera)]. – Dendrocos 29: 43-47.
- ENTING, K. (2003): Die Epiprocte der außeralpinen mitteleuropäischen Arten der Gattung *Nemoura* Latreille 1833 (Insecta: Plecoptera) [Epiprocts of the non-alpine Central European species of the genus *Nemoura* Latreille 1833 (Insecta: Plecoptera)]. – Lauterbornia 47: 131-140.
- ERKINARO, H.; ERKINARO, J. (2003): Distribution and nymphal age of *Dinocras cephalotes* (Curtis) (Plecoptera: Perlidae) in northern Finland. – Entomologica Fennica 14: 125-128.
- FAUSTO, A.M.; BELARDINELLI, M.C., TIERNOW DE FIGUEROA, J.M.; FOCHETTI, R.; MAZZINI, M. (2003): Ultrastructural studies on Plecoptera spermatozoa: New data and comparative remarks. – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 431-436.
- FENOGLIO, S. (2003): Feeding habits of *Anacroneuria* nymphs (Plecoptera Perlidae). — Bollettino della Società Entomologica Italiana 135(1): 15-17.
- FENOGLIO, S.; TIERNOW DE FIGUEROA, J.M. (2003): Observations on the feeding of adults of some *Neoperla* and *Anacroneuria* species (Plecoptera: Perlidae). – African Entomology 11(1): 138-139.
- FENOGLIO, S.; MORISI, A. (2001): *Anacroneuria talamanca* Stark 1988, especie nueva para la fauna de Nicaragua (Plecoptera, Perlidae). [*Anacroneuria talamanca* Stark 1988, a new species for the fauna of Nicaragua (Plecoptera, Perlidae).] — Revista Nicaraguense de Entomología 55-58: 5-9.

- FIGUEROA, R.; VALDOVINOS, C.; ARAYA, E.; PARRA, O. (2003): Macroinvertebrados bentónicos como indicadores de calidad de agua de ríos del sur de Chile. [Benthic macroinvertebrates as indicators of water quality of southern Chile rivers.] – Revista Chilena de Historia Natural 76(2): 275-285.
- FISCHER, P.; GUSTIN, M.S. (2002): Influence of natural sources on mercury in water, sediment and aquatic biota in seven tributary streams of the East Fork of the Upper Carson River, California. – Water Air Soil Pollution 133: 283-295.
- FLASPOHLER; D.J.; FISHER, H.C.J.; BUB, B.R.; VAN DUSEN, P.J. (2002): Temporal patterns in aquatic and avian communities following selective logging in the Upper Great Lakes Region. – Forest-Science 48(2): 339-349.
- FLEITUCH, T. (2003): Structure and functional organization of benthic invertebrates in a regulated stream. – Internat. Rev. Hydrobiol. 88: 332-344.
- FOCHETTI, R.; AMICI, I.; ARGANO, R. (2003): Seasonal changes and selectivity in the diet of brown trout in the River Nera (Central Italy). – J. Freshwat. Ecology 18(3): 437- 444.
- FOCHETTI, R.; TIENRO DE FIGUEROA, J.M. (2002): Biochemical evidence of a species boundary between *Isoperla nevada* Aubert, 1952 and *Isoperla grammatica* (Poda, 1761) (Plecoptera, Perlodidae). – Bol. Asoc. esp. Ent., 26(3-4): 131-135.
- FOCHETTI, R.; TIENRO DE FIGUEROA, J.M. (2002): Redescription of *Leuctra costai* Aubert, 1953 and considerations about its morphological variability (Plecoptera, Leuctridae). – Aquat. Insects 24(2): 137-141.
- FROEHLICH, C.G. (2003): Stoneflies (Plecoptera: Perlidae) from the Brazilian Amazonia with the description of three new species and a key to *Macrogynopax*. – Studies Neotropical Fauna Environment 38: 129-134.
- FÜREDER, L.; WELTER, C.; JACKSON, J. K. (2003): Dietary and stable isotope ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$) analyses in alpine Ephemeroptera and Plecoptera. – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 39-46.
- FÜREDER, L.; WELTER, C.; JACKSON, J. K. (2003): Dietary and stable isotope ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$) analyses in alpine stream insects. – Internat. Rev. Hydrobiol. 88: 314-331.
- GALAS, J.; DUMNICKA, E. (2003): Organic matter dynamics and invertebrate functional groups in a mountain stream in the West Tatra Mountains, Poland. – Internat. Rev. Hydrobiol. 88: 3362-371.
- GAYRAUD, S.; PHILIPPE, M. (2003): Influence of bed-sediment features on the interstitial habitat available for macroinvertebrates in 15 French streams. – Intern. Rev. Hydrobiol. 88: 77-93.

- GJERLOV, C.; HILDREW , A.G.; JONES, J.I. (2003): Mobility of stream invertebrates in relation to disturbance and refugia: a test of habitat templet theory. – J. N. Am. Benthol Soc. 22: 207-223.
- GONZALEZ, J.M.; BASAGUREN, A.; POZO, J. (2002): Size-mass relationship of stream invertebrates in a northern Spain stream. — Hydrobiologia 489: 131-137.
- GONZÁLEZ, J.M.; BASAGUREN, A.; POZO, J. (2003): Macroinvertebrate communities along a third-order Iberian stream. – Ann. Limnol. - Int. J. Lim. 39: 287-296.
- GRAF, W.; HUTTER, G. (2002): [New data on the stonefly fauna (Insecta, Plecoptera) of Austria.] Neue Daten zur Steinfliegenfauna Oesterreichs (Insecta, Plecoptera). – Linzer Biologische Beitraege 34(2): 1085-1090.
- GRAF, W.; HUTTER, G. (2003): Recent findings of the endangered stonefly *Brachyptera trifasciata* (PICTET, 1832) in Austria with a description of its habitat . – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 305-307.
- GRAF, W.; WEINZIERL, W. (2003): Distribution of *Brachyptera starmachi* SOWA, 1966 and *Capnia vidua rilensis* RAUSER, 1962. – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 309-311.
- GRAF, W.; SIVEC, I.; KOVACS, T. (2003): *Perla pallida* Guerin, 1838, in Oesterreich, Slowenien und Ungarn. *Perla pallida* Guerin, 1838, in Austria, Slovenia and Hungary. — Lauterbornia 47: 33-39.
- GRASSER, U.; GRAF, W. (2003): Erste Erfahrungen mit dem Potamon-Typie-Index (Schöll & Haybach 2001) in Österreich [First experiences with the application of the Potamon-Type-Index (Schöll & Haybach 2001) in Austria]. – Lauterbornia 47: 153-172.
- GRUBBS, S.A.; BRIGHT, E. (2003): *Arcynopteryx compacta* (Plecoptera: Perlodidae), a holarctic stonefly confirmed from Lake Superior, with a review and first checklist of the stoneflies of Michigan. – Great Lakes Entomologist 34(2): 77-84.
- HADEN, G.A.; SHANNON, J.P.; WILSON, K.P.; BLINN, D.W. (2003): Benthic community structure of the Green and Colorado rivers through Canyonlands National Park, Utah, USA. – Southwestern Naturalist. [print] 48(1): 23-35.
- HAHN, H.J. (2003): Eignen sich Fallen zur repräsentativen Erfassung aquatischer Meiofauna im hyporheischen Interstitial und im Grundwasser? [Are traps suitable for representative sampling of aquatic meiofauna in both the hyporheic zone and in groundwater?. Limnologica 33: 138-146. German.

- HALL, D.L.; BERGTHOLD, B.S.; SITES, R.W. (2003): The influence of adjacent land uses on macroinvertebrate communities of prairie streams in Missouri. – *J. Freshw. Ecol.* 18: 55-68.
- HALL, M.J.; CLOSS, G.P.; RILEY, R.H. (2001): Relationships between land use and stream invertebrate community structure in a South Island, New Zealand, coastal stream catchment. – *New Zealand J. Mar. and Freshw. Res.* 35: 591-603.
- HALVORSEN, G.A.; HEEGAARD, E.; FJELLHEIM, A.; RADDUM, G.G. (2003): Tracing recovery from acidification in the western Norwegian Nausta watershed. – *Ambio* 32(3): 235-239.
- HAMADA, N.; COUCEIRO, S.R.M. (2003): An illustrated key to nymphs of Perlidae (Insecta, Plecoptera) genera in Central Amazonia, Brazil. — *Revista Brasileira de Entomologia* 47(3): 477-480.
- HARDING, J.S. (2003): Historic deforestation and the fate of endemic invertebrate species in streams. – *N. Z. J. Marine Freshwat. Res.* 37(2): 333-345.
- HARPER, P.P.; HARPER, F. (2003): Comparison of Nearctic and Palearctic species groups of *Leuctra*: Affinities and origin of the North American fauna (Plecoptera: Leuctridae). – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 219-223.
- HEINO, J.; MUOTKA, T.; PAAVOLA, R.; PAASIVIRTA, L. (2003): Among-taxon congruence in biodiversity patterns: can stream insect diversity be predicted using single taxonomic groups? – *Can. J. Fish. Aquat. Sci.* 60: 1039-1049.
- HELMS, B.S.; FEMINELLA, J.W.; CHANEY, P.L. (2003): Land use and stream biotic signatures: The effects of urbanization on periphyton, macroinvertebrates, and fish in watersheds of west Georgia. – *Ecol. Soc. Amer. Annual Meeting Abstracts [cd-rom]* 88: 146.
- HICKEY, C.W.; GOLDING, L.A. (2002): Response of macroinvertebrates to copper and zinc in a stream mesocosm. – *Environ. Toxic. Chem.* 21(9): 1854-1863.
- HOFFSTEN, P.-O. (2003): Effects of an extraordinarily harsh winter on macroinvertebrates and fish in boreal streams. – *Arch. Hydrobiol.* 157: 505-523.
- HULL, M.S.; CHERRY, D.S.; SOUCEK, D.J.; CURRIE, R.J.; NEVES, R.J. (2002): Comparison of Asian clam field bioassays and benthic community surveys in quantifying effects of a coal-fired power plant effluent on Clinch River biota.. – *J. Aquat. Ecosystem Stress Recovery* 9(4): 271-283.

- HUMPESCH, U.H.; ELLIOTT, J.M. (2003): Egg hatching: One mechanism for life-cycle partitioning in Plecoptera and Ephemeroptera. – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 187-188.
- HUNT, G. W.; STANLEY, E. H. (2003): Environmental factors influencing the composition and distribution of the hyporheic fauna in Oklahoma streams: Variation across ecoregions. – Arch. Hydrobiol. 158: 1-23.
- HURYN, A.D.; RILEY, R.H.; YOUNG, R.G.; ARBUCKLE, C.J.; PEACOCK, K.; LYON, G. (2001): Temporal shift in contribution of terrestrial organic matter to consumer production in a grassland river. — Freshwater Biology 46(2): 213-226.
- IANNILLI, V.; TIERNO DE FIGUEROA, J.M.; FOCHETTI, R. (2002): Life cycle of *Dinocras cephalotes* (Curtis, 1827) in Central Italy (Plecoptera, Perlidae). — Bol. S.E.A. 31: 177-179.
- JACOBSEN, D. (2003): Altitudinal changes in diversity of macroinvertebrates from small streams in the Ecuadorian Andes. – Arch. Hydrobiol. 158: 145-167.
- JEZBEROVA, M. (2003): Distribution and density of Ephemeroptera and Plecoptera of the Radíkovský brook (Czech Republic) in relation to selected environmental variables. – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 327-331.
- JOHNSON, B.R.; CROSS, W.F.; WALLACE, J.B. (2003): Long-term resource limitation reduces insect detritivore growth in a headwater stream. – J. N. Am. Benthol. Soc. 22: 565-574.
- JONSSON, M.; DANGLES, O.; MALMQVIST, B.; GUEROLD, F. (2002): Stimulating species loss following perturbation: Assessing the effects on process rates. – Proc. R. Soc. Biol. Sci. Series B 269 (1495): 1047-1052.
- JONSSON, M.; MALMQVIST, B. (2003): Mechanisms behind positive diversity effects on ecosystem functioning: Testing the facilitation and interference hypotheses. — Oecologia 134(4): 554-559.
- JOOST, W. (2002): Neufunde von *Leuctra major* Brinck 1949 (Plecoptera, Leuctridae) in Thüringen. [New finds of *Leuctra major* Brinck 1949 (Plecoptera, Leuctridae) in Thuringia]. – Abh. Ber. Mus. Natur Gotha 22: 185-186.
- KAPOOR, N.N. (2003): Fine structure of the compound eye of a stonefly nymph, *Paragnetina media* (WALKER) (Plecoptera). – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 453-456.
- KEILER, J.-A. (2002): Bedeutende Fossilvorkommen des Quartärs in Thüringen, Teil 3: Insekten. [Important occurrences of quaternary fossils in Thuringia, Part 3: Insects]. – Beitr. Geol. Thür., N.F., 9: 173-186.

- KLEMM, D.J.; BLOCKSM, K.A.; FULK, F.A.; HERLIHY, A.T.; HUGHES, R.M.; KAUFMANN, P.R.; PECK, D.V.; STODDARD, J.L.; THOENY, W.T.; GRIFFITH, M.B.; DAVIS, W.S. (2003): Development and evaluation of a Macroinvertebrate Biotic Integrity Index (MBII) for regionally assessing mid-atlantic highlands streams. – Environmental Management 31(5): 656-669.
- KOBAYASHI, S.; KAGAYA, T. (2002): Differences in litter characteristics and macroinvertebrate assemblages between litter patches in pools and riffles in a headwater stream. – Limnology 3(1): 37-42.
- KÖHLER, J.; GELBRECHT, J.; PUSCH, M. (2002): Die Spree Zustand, Probleme, Entwicklungsmöglichkeiten [River Spree Condition, problems possibilities for development]. – 1. Aufl. Stuttgart; Limnologie aktuell 10: E. Schweizerbart'sche Verlagsbuchhandlung. 384 pp.
- KONDRATIEFF, B.C.; KIRCHNER, R.F. (2003): Two new species of *Perlesta* Banks (Plecoptera: Perlidae) from eastern North America. – Proc. Entomol. Soc. Wash. 105(4): 933-939.
- KONDRATIEFF, B.C.; ZUELLIG, R.E. (2003): Stoneflies of Fort Sill, Oklahoma, U.S.A. – Perla 21: 13-17.
- KONNO, Y.; NISHIMOTO, H.; MARUYAMA, H.; TORII, T.; ISHIWATA, S. (2003): Lotic aquatic insects in the alpine zone of Daisetsuzan National Park, Hokkaido – Jap. J. Limnol. 64: 141-144
- KRNO, I. (2002): Novy nález ohrozeného druhu posvátky *Capnopsis schilleri* (Plecoptera: Capniidae) po 50 rokoch na Slovensku (Horná Orava). New report of endangered species of stonefly *Capnopsis schilleri* (Plecoptera: Capniidae) after 50 years in Slovakia (Horná Orava). — Folia Faunistica-Slovaca 7: 23-24.
- KRNO, I. (2002): Posvátky (Plecoptera) NPR Rozsutec (NP Mala Fatra). Stoneflies (Plecoptera) of the national Nature Reserve Rozsutec (National Park Mala Fatra). — Folia Faunistica Slovaca 7: 25-29.
- KRNO, I. (2003): Distribution patterns and habitats of stoneflies in Slovakia. – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 349-356.
- LENCIOMI, V.; MAIOLINI, B.; ZUCCATI, S. (2001): Temporal drift patterns in a high mountain stream system (Conca Adamello-Brenta Natural Park, Trentino, Italy). – Studi Trentini di Scienze Naturali Acta Biologica. 76: 113-123.
- LEPORI, F.; BARBIERI, A.; ORMEROD, S. J. (2003): Effects of episodic acidification on macroinvertebrate assemblages in Swiss Alpine streams. – Freshwat. Biol. 48: 1873-1885.

- LÓPEZ-RODRÍGUEZ, M.J.; LUZÓN-ORTEGA, J.M.; TIERNO DE FIGUEROA, J.M. (2003): Contribución al conocimiento de los plecópteros (Insecta, Plecoptera) del Parque Natural de las Sierras de Almijara, Tejeda y Alhama y áreas próximas (SE España), con especial referencia a sus singularidades faunísticas. – Boletín de la S.E.A., 33: 167-171.
- LOSKUTOVA, O. (2003): Fauna of stoneflies (Plecoptera) of the European North-East Russia and its change under the influence of anthropogenic factors. – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 357-362.
- LUZON-ORTEGA, J.M.; TIERNO DE FIGUEROA, J.M. (2002): Los Plecopteros (Insecta, Plecoptera) de Montes de Toledo (España). [The stoneflies (Insecta, Plecoptera) from Montes de Toledo (Spain)]. – Bol. S.E.A. 30: 175-176.
- LUZON-ORTEGA, J. M.; TIERNO DE FIGUEROA, J.M. (2003): Some aspects of behaviour of *Nemoura fulviceps* (Plecoptera: Nemouridae). – In: GAINO, E. (ed), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 47-53.
- LUZON-ORTEGA, J.M.; TIERNO DE FIGUEROA, J.M. JAIMEZ-CUELLAR, P. (2003): Prey selection pattern in *Notonecta maculata* (Hemiptera, Notonectidae). – Belgian Journal of Zoology, 2003, 133(2): 189-190.
- MacNEIL, C.; DICK, J.T.A.; BIGSBY, E.; ELWOOD, R.W.; MONTGOMERY, W.I.; GIBBINS, C.N.; KELLY, D.W. (2002): The validity of the *Gammarus:Asellus* ratio as an index of organic pollution: Abiotic and biotic influences. – Water Research 36(1): 75-84.
- MALARD, F.; GALASSI, D.; LAFONT, M.; DOLEDEC, S.; WARD, J.V. (2003): Longitudinal patterns of invertebrates in the hyporheic zone of a glacial river. – Freshwat. Biol. 48: 1709-1725.
- MALICKY-RUZICKA, H.M.; MALICKY, M. (2002): Rückkehr einer bedrohten Steinfliegenart oder Sammelartefakt - Versuch einer Interpretation am Beispiel *Brachyptera monilicornis* (Insecta: Plecoptera, Taeniopterygidae) [Return of a threatened stonefly species or collecting artifact. Attempt of an interpretation by the example *Brachyptera monilicornis* (Insecta: Plecoptera, Taeniopterygidae)]. Lauterbornia 47: 27-32.
- MARDEN, J.H.; THOMAS, M.A. (2003): Rowing locomotion by a stonefly that possesses the ancestral pterygote condition of co-occurring wings and abdominal gills.. – Biol. J. Linn. Soc. 79(2): 341-349.
- MARUYAMA, H. (2002): A new species of the genus *Cryptoperla* (Plecoptera, Peltoperlidae) from Japan. – Japanese J. Syst. Ent. 8: 1-6.

- MAUCH, E.; SCHMEDTJE, U.; MAETZE, A.; FISCHER, F. (2003): Taxaliste der Gewässerorganismen Deutschlands zur Kodierung biologischer Befunde. [Taxalist of aquatic organisms for coding biological findings]. – Informationsberichte Bayerisches Landesamt für Wasserwirtschaft 390 pp.
- MAVRI, U.; PODGORNIK, S.; SIVEC, I.; TOMAN, M.J. (2003): An approach to the study of ecology of the stonefly distribution in the Lipnica Stream, Slovenia. – In: GAINO, E. (ed.), Research on Ephemeroptera and Plecoptera; University of Perugia; pp. 363-366.
- MCINTOSH, A.R. (2002): Impact of introduced trout on *Nesameletus* mayfly populations measured using electrofishing. – Arch. Hydrobiol. 154(3): 375-392.
- MCLELLAN, I. (2003): Stewart Island stoneflies. – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 367-371.
- MCLELLAN, I. D. (2003): Six new species and a new genus of stoneflies (Plecoptera) from New Zealand. N. Z. J. Zool. 30: 101-113.
- MCNEELY, C.; POWER, M.E. (2003): Role of the armored grazer *Glossosoma penitum* in a headwater stream food web. – Ecol. Soc. Amer. Annual Meeting Abstracts [cd-rom] 88: 230-231.
- MEYER, E.I.; POEPERL, R. (2003): Secondary production of invertebrates in a Central European mountain stream (Steina, Black Forest, Germany). – Arch. Hydrobiol. 158: 25-42.
- MINAKAWA, N.; GARA, R.I. (2003): Effects of chum salmon redd excavation on benthic communities in a stream in the Pacific Northwest. – Trans. Amer. Fisheries Soc: 132(3): 598-604.
- MISERENDINO, M.L.; (2001): Length-mass relationships for macroinvertebrates in freshwater environments of Patagonia (Argentina). – Ecol. Austral. 11: 3-8.
- MORSE, C.C.; HURYN, A.D.; CRONAN, C. (2003): Impervious surface area as a predictor of the effects of urbanization on stream insect communities in Maine, U.S.A. – Environ. Monitoring and Assessment 89: 95-127.
- MUCHOW, C.L.; RICHARDSON, J.S. (2000): Unexplored diversity: macroinvertebrates in coastal British Columbia headwater streams, pp. 503-506. – In: DARLING, L.M. (ed). Proceedings of a conference on the biology and management of species and habitats at risk. Kamloops, B.C. 5-19 Feb, 1999. – Vol. 2, B.C. Ministry of Environment, Lands and Parks, Victoria and University College of the Cariboo, Kamloops.
- MURANYI, D. (2002): A *Nemoura fusca* Kis, 1963 alkeresz (Plecoptera) faj hazai elfordulása. The occurrence of *Nemoura fusca* Kis, 1963 (Plecoptera) in Hungary. — Folia Entomologica Hungarica 63: 195-196.

- MUSTOW, S.E. (2002): Biological monitoring of rivers in Thailand: use and adaptation of the BMWP score. *Hydrobiologia* 479: 191-229.
- NERBONNE, J.F.; VONDRAČEK, B. (2003): Volunteer macroinvertebrate monitoring: assessing training needs through examining error and bias in untrained volunteers. – *J. N. Am. Benthol. Soc.* 22: 152-163.
- NESTEROVITCH, A.; ZWICK, P. (2003): The development of *Nemurella picteti* Klapálek (Plecoptera: Nemouridae) in two springstreams in central Europe. – *Limnologica* 33(4): 231-243.
- NIYOGI, D.K.; SIMON, K.S.; TOWNSEND, C.R. (2003): Breakdown of tussock grass in streams along a gradient of agricultural development in New Zealand. – *Freshwat. Biol.* 48: 1698-1708.
- OBACH, M.; WAGNER, R.; WERNER, H.; SCHMIDT, H.H. (2001): Modelling population dynamics of aquatic insects with artificial neural networks. – *Ecol. Model* 146(1-3): 207-217.
- ORCE, M. (2003): A new species of *Anacroneuria* from Northwestern Argentina (Plecoptera: Perlidae). – In: GAINO, E. (ed.), Research update on Ephemeropter and Plecoptera; University of Perugia; pp. 225-228.
- ORTAZ, M. (2001): Diet seasonality of food among fishes of the upper Orituco stream, northern Venezuela. – *Revista de Biol. Tropical.* 49: 191-197.
- OTTO, A.; WESTERMANN, F. (2003): Erstnachweis von *Isoptena serricornis* (Pictet 1841) (Plecoptera, Chloroperlidae) in Rheinland-Pfalz / First record of *Isoptena serricornis* (Pictet 1841) (Plecoptera, Chloroperlidae) in Rhineland-Palatinate. – *Lauterbornia* 46: 103-106.
- PALMA, A.; FIGUEROA, R.; RUIZ, V.H.; ARAYA, E.; BERRIOS P. (2002): [Composition of the diet of *Oncorhynchus mykiss* (Walbaum 1792) (Pisces: Salmonidae) in a river system of low anthropogenic action: Nonguen stream, VIII Region, Chile.] Composicion de la dieta de *Oncorhynchus mykiss* (Walbaum 1792) (Pisces: Salmonidae) en un sistema fluvial de baja intervencion antropica: Estero nonguen, VIII Region, Chile. – *Gayana* 66(2): 129-139.
- PARK, Y.; CEREGHINO, R.; COMPIN, A.; LEK, S. (2003): Applications of artificial neural networks for patterning and predicting aquatic insect species richness in running waters. – *Ecol. Modelling* 160: 265-280.
- PENNUTO, E.M.; WOOSTER-BROWN, C.L.; BELISLE, C.A. (2002): Infestation intensity and prevalence of an ectosymbiotic midge und variable environmental and host conditions. – *Can. J. Zool.* 80(12): 2061-2071.

- PETERSEN, I.; HILDREW, A.G. (2003): Emergence of *Leuctra nigra* (Plecoptera) from a southern English stream. – Arch. Hydrobiol. 158: 185-195.
- PETERSON, J.L.; JEPSON, P.C.; JENKINS, J.J. (2001): A test system to evaluate the susceptibility of Oregon, USA, native stream invertebrates to triclopyr and carbaryl. – Environ. Toxicol. and Chem. 20: 2205-2214.
- POULTON, B.C.; WILDHABER, M.L.; CHARBONNEAU, C.S.; FAIRCHILD, J.F.; MUELLER, B.G.; SCHMITT, C.J. (2003): A longitudinal assessment of the aquatic macroinvertebrate community in the channelized lower Missouri River. – Envir. Monitoring & Assessment 85(1): 23-53.
- PUPILLI, E.; PUIG, M.A. (2003): Effects of a major flood on the mayfly and stonefly populations in a Mediterranean stream (Matarranya Stream, Ebro River basin, North East of Spain). – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 381-389.
- PUREVDORJ, S.; MÜHLENBERG.M.; SLOWIK, J. (2003): First records of the genus *Paraleuctra* Hanson (Plecoptera: Leuctridae) and *Isoperla flavescens* Zhiltzova et Potikha (Plecoptera: Perlodidae) for the stonefly fauna of Mongolia. – Mongolian J. Biol. Sci. 1: 27-30.
- QUAMME, D.L.; SLANEY, P.A. (2003): The relationship between nutrient concentration and stream insect abundance. – In: Stockner, John G.: Ed. American Fisheries Society Symposium. Nutrients in salmonid ecosystems: Sustaining production and biodiversity 34: 163-175.
- RASMUSSEN, A.K.; PESCADOR, M.L.; RICHARD, B.A. (2003): A survey of the Plecoptera of Florida. – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 391-402.
- RAVIZZA, C.; VINÇON, G. (2003): *Leuctra queirassiana orsiera*, a new subspecies of *Leuctra* from the Cottian Alps, Italy (Plecoptera Leuctridae). – Bull. Soc. entomol. ital. 135(1): 19-23.
- REICH, P.; DOWNES, B.J. (2003): The distribution of aquatic invertebrate egg masses in relation to physical characteristics of oviposition sites at two Victorian upland streams. – Freshwat. Biol. 48: 1497-1513.
- REIDELBACH, J.; CHRISTL, H. (2002): A quantitative investigation into the temporal and spatial variations in the emergence of adult blackflies (Diptera: Simuliidae) from the Breitenbach, a small upland stream in Germany. – Limnologica 32(3): 206-235.
- REYOL, Y.; COMPIN, A.; IBARRA, A.A.; LIM, P. (2003): Longitudinal diversity patterns in streams: comparing invertebrates and fish communities. – Arch. Hydrobiol. 157: 525-533.

- ROBINSON, C.T.; UEHLINGER, U.; MONAGHAN, M.T. (2003): Effects of a multi-year experimental flood regime on macroinvertebrates downstream of a reservoir. – *Aquatic Sci.* 65: 210-222.
- ROGERS, C.E.; BRABANDER, D.J.; BARBOUR, M.T.; HEMOND, H.F. (2002): Use of physical, chemical, and biological indices to assess impacts of contaminants and physical habitat alteration in urban streams. — *Environmental Toxicology and Chemistry* 21(6): 1156-1167.
- ROSCISZEWSKA, E. (2003): Ultrastructure of egg capsules of the South American stoneflies (Plecoptera: Perlidae). – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 463-467.
- RUPPRECHT, R. (2003): Drumming signals within the genus *Dinocras* (Plecoptera: Perlidae). – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 63-72.
- SÁNCHEZ ORTEGA, A.; ALBA-TERCEDOR, J.; TIERNO DE FIGUEROA, J.M. (2002): Lista faunística y bibliográfica de los Plecópteros de la Península Ibérica e Islas Baleares. – Publicaciones de la Asociación española de Limnología. N° 16. 198 pp.
- SANDBERG, J.B.; STEWART, K.W. (2003): Continued studies of drumming in North American Plecoptera; Evolutionary implications. – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 73-81.
- SANDIN, L. (2003): Benthic macroinvertebrates in Swedish streams: community structure, taxon richness, and environmental relations. – *Ecography* 26: 269-282.
- SCHLEUTER, M.; HAYBACH, A. (2003): Das Makrozoobenthos des Mains in den Jahren 1992-2001 - Eine Artenliste (The macroinvertebrate fauna from the River Main in the period 1992-2001 - list of the species). – *Lauterbornia* 48: 45-56.
- SCHUSTER, C.; BELLSTEDT, R.; GÖRING, M.; MÜLLER, U.; SAMIETZ, R. (2002): Die Flora und Fauna des Flächennaturdenkmals "Gewässerschutzgebiet Hörsel" im Landkreis Gotha mit Anmerkungen zu Pflege und Entwicklung. [Flora and fauna of the nature reserve "Gewässerschutzgebiet Hörsel" in Gotha county with comments on care and development]. – *Abh. Ber. Mus. Natur Gotha* 22: 59-80.
- SEHADOVA, H.; SAUMAN, I.; SEHNAL, F. (2003): Immunocytochemical distribution of pigment-dispersing hormone in the cephalic ganglia of polyneopteran insects. – *Cell and Tissue Res.* 312: 113-125.

- SCUDDER, G.G.E.; NEDHAM, K.M.; KENNER, R.D.; CANNINGS, R.A.; CANNINGS, S.G. (2001): Aquatic insects in British Columbia: 100 years of study. – J. Entomol. Soc. British Columbia 98: 61-81.
- SHERK, T.E.; RAU, G.H.; KRAFT, G.F. (2003): Emergence of Plecoptera from Findley Lake, Cascade Mountains, USA. – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 407-411.
- SHIEH, S.-H.; WARD, J.V.; KONDRATIEFF, B.C. (2003): Longitudinal changes in macroinvertebrate production in a stream affected by urban and agricultural activities. – Arch. Hydrobiol. 157: 483-503.
- SIVEC, I.; DIA, A. (2003): *Isoperla berthelemyi* and *Protonemura phoenicia* spp. n., two new species of stoneflies from Lebanon [Insecta: Plecoptera]. – Ephemera (2001) 3(1): 43-51.
- SIVEC, I.; GRAF, W. (2003): *Perla carantana* – a new species of the genus *Perla* (Plecoptera: Perlidae) from Austria and Slovenia. – Natura Sloveniae 4: 31-38.
- SPACEK, J.; SOLDAN, T.; PUTZ, M. (1999): *Capnopsis schilleri* (Plecoptera: Capniidae) in South Bohemia, Czech Republic. — Silva Gabreta 3: 115-121.
- SPEIRS, D.C.; GURNEY, W.S.C (2001): Population persistence in rivers and estuaries. — Ecology 82(5): 1219-1237.
- STANICZEK, A.H. (2003): Neufund der Steinfliege *Capnopsis schilleri* (Plecoptera: Capniidae) in Baden-Württemberg / New record of the stonefly *Capnopsis schilleri* (Plecoptera: Capniidae) in Baden-Württemberg. – Mitt. ent. V. Stuttgart 38: 9-12.
- STARK, B.P.; ZUNIGA DEL, M.C. (2003): The *Anacroneuria guambiana* complex of South America (Plecoptera: Perlidae). – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 229-237.
- STEPENUCK, K.F.; CRUNKILTON, R.L.; WANG, L.Z. (2002): Impacts of urban land use on macroinvertebrate communities in southeastern Wisconsin streams. – J. Amer. Water Resour. Assoc. 38(4): 1041-1051.
- STEWART, K.W. (2003): Plecoptera (Stoneflies). pp. 141-146. – In: HUTCHINS, M.; EVANS, A.V.; GARRISON, R.W.; SCHLAGER, N. (eds.). Grzimeks animal life encyclopedia, 2nd ed., Vol 3. – Gale Group Pubs., Farmington Hills, Michigan. 472 pp.
- STEWART, K.W. (2003): Plecoptera (Stoneflies). – pp. 915-919 in: RESH & CARDÉ (eds), Encyclopedia of Insects, 1266 pp., Academic Press, Elsevier Science, USA.

- STEWART, K.W.; SANDBERG, J.B. (2003): Life history of a Colorado population of *Kogotus modestus* (Plecoptera: Perlodidae). – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 195-200.
- STEWART, T.W.; SHUMAKER, T.L.; RADZIO, T.A. (2003): Linear and nonlinear effects of habitat structure on composition and abundance in the macroinvertebrate community of a large river. – American Midland Naturalist 149(2): 293-305.
- STLOUKAL, E. (2001): Databanka fauny Slovenska - stav a nove prirastky databaz v roku 2001. Current status and new data in Databank of Slovak fauna in year 2001. — Folia Faunistica Slovaca 6: 145-168.
- TERRY, M.D.; WHITING, M.F. (2003): Phylogeny of Plecoptera: Molecular evidence and evolutionary trends. – Entomologische Abhandlungen 61(3): 130-131.
- TESLENKO, V. (2003): Zoogeography of the stoneflies (Plecoptera) of the Kuril Archipelago. – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 421-427.
- TESLENKO, V.A.; ZHILTSOVA, L.A. (2003): Nymphs of the genus *Isocapnia* (Plecoptera, Capniidae) from Palaearctic. – Zoologicheskii Zhurnal 82(3): 354-365.
- THOMPSON, R.M.; TOWNSEND, C.R. (2003): Impacts on stream food webs of native and exotic forest: an intercontinental comparison. – Ecology 81: 145-161.
- TIERNO DE FIGUEROA, J.M. (2003): Mate guarding and displacement attempts in stoneflies (Plecoptera). – Biologia, Bratislava 58(5): 925-928.
- TIERNO DE FIGUEROA, J.M.; DERKA, T. (2003): Egg description of *Isoptena serricornis* (Plecoptera: Chloroperlidae). – Entomological Problems 33(1-2): 55-57.
- TIERNO DE FIGUEROA, J.M.; LUZON-ORTEGA, J.M. (2002): The mating call of *Isoperla bipartita* Aubert, 1962 (Plecoptera, Perlodidae). – Aquat. Insects 24(2): 87-90.
- TIERNO DE FIGUEROA, J.M.; LUZON-ORTEGA, J.M.; SANCHEZ-ORTEGA, A. (2003): Protandry and its relationship with adult size in some Spanish stoneflies species (Plecoptera). – Annals Entomol. Soc. Amer. 96(4): 560-562.
- TIERNO DE FIGUEROA, J.M.; SÁNCHEZ-ORTEGA, A.; MEMBIELA-IGLESIAS, P.; LUZÓN-ORTEGA, J.M. (2003): Plecoptera. – In: Fauna Ibérica, vol. 22. RAMOS, M.A. et al. (Eds.). Museo Nacional de Ciencias Naturales. CSIC. Madrid. 2003. 404 pp.
- TIERNO DE FIGUEROA, J.M.; SEZZI, E.; FOCHETTI, R. (2003): Feeding in the genus *Tyrrhenoleuctra* (Plecoptera, Leuctridae). – Bollettino de la Società Entomologica Italiana, 134(3): 207-210.

- VALLADARES, L.F.; VEGA, F.J.; MAZE, R.A.; REGIL, J.A.; GARCIA-CRIADO, F. (2002): Biodiversity of aquatic macroinvertebrates in the Natural Park of Valderejo (Alava): Implications for conservation. – Bol. de la Assn. Espanola de Entomol. 26: 37-55.
- VAN DER GEEST, H.G.; SOPPE, W.J.; GREVE, G.D.; KROON, A.; KRAAK, M.H.S. (2002): Combined effects of lowered oxygen and toxicants (cooper and diazinon) on the mayfly *Ephoron virgo*. – Environ. Toxicol. Chem. 21(2): 431-436.
- VELASQUEZ, S.M.; MISERENDINO, M.L. (2003): Habitat type and macroinvertebrate assemblages in low order Patagonian streams. – Arch. Hydrobiol. 158: 461-483.
- VINÇON, G.; PARDO, I. (2003): Two new *Nemoura* species from the northwestern Iberian Peninsula and the Pyrenees (Plecoptera, Nemouridae). – Nouv. Rev. Entomol. 20(1): 29-38.
- VINSON, M.R.; HAWKINS, C.P. (2003): Broad-scale geographical patterns in local stream insect genera richness. – Ecography 26: 751-767.
- VRBA, J.; KOPACEK, J.; FOTT, J.; KOHOUT, L.; NEDBALOVA, L.; PRAZAKOVA, M.; SOLDAN, T.; SCHAUUMBURG, J. (2003): Long-term studies (1871-2000) on acidification and recovery of lakes in the Bohemian Forest (central Europe). – Sci. Total Environment 310(1-3): 73-85.
- WAGNER, F. (2003): Flight behaviour of merolimnic insects from the Leutra River (Thuringia, Germany). – Aquatic Insects 25(1): 51-62.
- WALKER, I. (2003): The benthic insect fauna of the blackwater forest stream Rio Tarumã-Mirim (Manaus, Amazonas): patterns of population dynamics and their implications for ecosystem stability. – Amazoniana 17: 471-480.
- WANTZEN, K. M. (2003): Cerrado streams - characteristics of a threatened freshwater ecosystzem type on the Tertiary Shields of Central South America. – Amazoniana 17: 481-502.
- WEATHERHEAD, M.A.; JAMES, M.R. (2001): Distribution of macroinvertebrates in relation to physical and biological variables in the littoral zone of nine New Zealand lakes. – Hydrobiologia. 462: 115-129.
- WEIGEL, B. M. (2003): Development of stream macroinvertebrate models that predict watershed and local stressors in Wisconsin. – J. N. Am. Benthol. Soc. 22: 123-142.
- WEIGEL, B.M.; WANG, L.; RASMUSSEN, P.W.; BUTCHER, J.T.; STEWART, P.M.; SIMON, T.P.; WILEY, M.J. (2003): Relative influence of variables at multiple spatial scales on stream macroinvertebrates in the Northern Lakes and Forest ecoregion, U.S.A. – Freshwat. Biol. 48(8): 1440-1461.

- WEIGELHOFER, G.; WARINGER, J. (2003): Vertical distribution of benthic macroinvertebrates in riffles versus deep runs with differing contents of fine sediments (Weidlingbach, Austria). – Internat. Rev. Hydrobiol. 88: 304-313.
- WEIGELHOFER, G.; WARINGER, J.A. (2003): Response of macroinvertebrates to fine sediment accumulations within the hyporheic zone of a calcareous sandstone stream (Weidlingbach, Austria). – Arch. Hydrobiol. Suppl. 147: 327-346.
- WESTERMANN, F. (2003): *Capnia bifrons* (Insecta, Plecoptera) als Leitart sommertrockener Bäche in Rheinland-Pfalz [*Capnia bifrons* (Insecta, Plecoptera) as indicator-species of intermittent small streams in Rhineland-Palatinate (Germany)]. – Lauterbornia 46: 107-119.
- WINKELMANN, C.; KNOOP, J.H.E.; BENNDORF, J. (2003): Abiotic features and macroinvertebrate colonization of the hyporheic zone of two tributaries of the river Elbe (Germany). – Limnologica 33: 112-121.
- WINTER, J.G.; SOMERS, K.M.; DILLON, P.J.; PATERSON, C.; REID, R.A. (2002): Impacts of golf courses on macroinvertebrate community structure in Precambrian Shield streams. – J. Environ. Qual. 31(6): 2015-2025.
- WOODWARD, G.; HILDREW, A.G. (2002): The impact of a sit-and-wait predator: separating consumption and prey emigration. – Oikos 99(3): 409-418.
- YASICK, A.L.; SIMMONS, T.W.; EARLE, J.I. (2003): Parasitic water mite larvae (Acari: Stygothrombidiidae) associated with stonefly adults (Insecta: Plecoptera) from an Allegheny mountain stream. – Pp. 323-330 in: SMITH, I.M. (ed.): An acarological tribute to David R.Cook: From Yankee Springs to Wheeny Creek. Indira Publishing House, West Bloomfield, MI. 331 pp.
- YOSHIMURA, M. (2003): Difference of ovariole maturation depending on emergence date in *Isoperla aizuna* (Perlodidae: Plecoptera). – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 211-215.
- YOSHIMURA, M. (2003): Relations of intraspecific variations in fecundity, clutch size, and oviposition frequency to the body size in three species of stoneflies, *Sweltsa* sp., *Isoperla aizuna*, and *Stavsolus japonicus*. – Limnology 4(2): 109-112.
- YOSHIMURA, M.; ISOBE, Y.; OISHI, T. (2003): The relation among emergence date, days until first effective mating, fecundity and adult longevity in *Isoperla aizuna* and *Sweltsa* sp. (Plecoptera). – Zool. Sci. (Tokyo) 20(4): 471-479.

- ZAVODSKA, R.; SAUMAN, I.; SEHNAI, F. (2003): Distribution of PER protein, pigment-dispersing hormone, prothoracitropic hormone, and eclosion hormone in the cephalic nervous system of insects. – J. Biol. Rhythms 18: 106-122.
- ZHILTZOVA, L (2003): The Leuctridae (Plecoptera) of Russia and adjacent territories (within the limits of the former USSR) . – In: GAINO, E. (ed.), Research update on ephemeroptera and Plecoptera; University of Perugia; pp. 239-244.
- ZHILTZOVA, L.A. (2003): Plecoptera Gruppe Euholognatha. – Fauna of Russia and Neighbouring Countries, Insecta Plecoptera, 1(1): 537 + 7 unnumbered pages; Russian Academy of Sciences, Zoological Institute, St. Petersburg, Nauka; ISBN 5-02-02613-7.
- ZHILTZOVA, L.A.; CHERCHESOVA, S.K. (2003): Descriptions of the larvae of stoneflies of the genus *Perla* Geoffr. (Plecoptera, Perlidae) from the Caucasus. — Entomol. Obozr. 82(2): 321-326, 530.
- ZHU, F.; YANG, D. (2003): Two new species of *Nemoura* from China (Plecoptera, Nemouridae). – Acta Zootaxonomica Sinica 28(3): 474-477. Chinese, English abstract.
- ZHU, F.; YANG, D.; YANG, C. (2002): A new species of the genus *Indonemoura* Baumann (Plecoptera; Nemouridae) from Tibet, China. – Entomol. Sci. 5: 317-320. (correction of this citation in PERLA 21).
- ZHU, F.; YANG, D. (2003): Three new species of *Amphinemura* (Plecoptera: Nemouridae) from Tibet, China [sic!]. — Entomologia Sinica 10(1): 51-56.
- ZHU, F.; YANG, D.; YANG, C.K. (2003): Two New Species of *Mesonemoura* from Tibet, China (Plecoptera: Nemouridae). — Aquatic-Insects 25(1): 1-8.
- ZIEMANN, H. (2002): Veränderungen der Besiedelungsstrukturen in einem Flussabschnitt der Gera nach der Abwasserlastsenkung durch den Neubau der Verbandskläranlage Arnstadt. [Changes in colonisation structure in a section of River Gera after reduced waste water impact by installation of the Arnstadt waste water treatment plant]. – Abh. Ber. Mus. Nat. Gotha 22: 81-88.
- ZUELLIG, R.E.; KONDRATIEFF, B.C.; RHODES, H.A. (2002): Benthos recovery after an episode sediment release into a Colorado Rocky Mountain river. – Western N. Amer. Natur. 62: 59-72. (correction of this citation in PERLA 21 p. 39).
- ZWICK, P. (2003): Morphological support of the major clades of Plecoptera. – Entomologische Abhandlungen 61(2): 128-130.
- ZWICK, P. (2003): 8. Ordnung Plecoptera, Steinfliegen, Frühlingsfliegen, Uferfliegen. – Kaestner, Lehrb. Zool. 1(5), 2. Aufl.: 144-154; Spektrum Akademischer Verlag, Heidelberg.

- ZWICK, P. (2003): Plecoptera research today: questions to be asked in the new millennium. – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 245-251.
- ZWICK, P. (2003): Shapes and patterns of wingpad development in the Plecoptera. – In: GAINO, E. (ed.), Research update on Ephemeroptera and Plecoptera; University of Perugia; pp. 477-483.
- ZWICK, P. (2003): The stonefly (Insecta: Plecoptera) seed bank theory: new experimental data. - Verh. Intern. Verein. Limnol. 28: 1317-1323.
- ZWICK, P.; HOHMANN, M. (2003): Direct development, no diapause, in *Taeniopteryx nebulosa* (Plecoptera, Taeniopterygidae). – Lauterbornia 47: 141-151.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Perla](#)

Jahr/Year: 2004

Band/Volume: [22](#)

Autor(en)/Author(s): Redaktion

Artikel/Article: [Recent Plecoptera Literature 25-45](#)